

# ALB IANA

*The 'crowning' of the erathem boundary*



*from left: Aymon Baud, Yuri Zacharov, Mike Orchard, Rich Lane, Yin Hongfu*

## Contents

	<i>Pages</i>
<b>Executive Notes</b>	<b>3</b>
<b>Reports</b>	
The new GSSP, base of the Triassic: some consequences. <i>Baud A.</i>	6
Functional Importance of New Skeletal Elements («Eye Capsules») of Euconodonts <i>Buryi G. I., and Kasatkina A. P.</i>	9
Age and Correlation of Triassic Tetrapod Assemblages from Brazil. <i>Lucas S. G.</i>	13
Triassic marine reptiles from China <i>Li J.-L., Liu J., Li C.</i>	21
A proposed area for the study of the Accessory Section and Point of the Terrestrial Permian-Triassic Boundary <i>Peng Y., Yin H., Yang F. and Wang S.</i>	26
<b>Meeting Reports</b>	
Oman Pangea Symposium and Fieldmeeting, Muscat, Sultanate of Oman, 7-20 January, 2001. <i>Baud A., Marcoux J., Beauchamp B.</i>	32
The Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events, Changxing, China, August 10-13, 2001. <i>Hongfu Y., Jinnan T.</i>	34
Report on the field excursion in western Guizhou and eastern Yunnan <i>Yuanqiao P., Shangyan W. and Songzhu G.</i>	42
<b>New Triassic Literature</b>	
Triassic Bibliography <i>Warrington G., Kerp H., Kürschner W. M., Lucas S. G., Visscher H.</i>	46
British Triassic Literature: Supplement 26 <i>Warrington G.</i>	59
New Books	60
<b>Future Meetings</b>	62
<b>STS Members List 2001</b>	64
<b>Guidlines for the submission of manuscripts</b>	72

---

The primary aim of ALBERTIANA is to promote the interdisciplinary collaboration and understanding among members of the I.U.G.S. Subcommittee on Triassic stratigraphy. Within this scope ALBERTIANA serves as the newsletter for the announcement of general information and as a platform for discussion of developments in the field of Triassic stratigraphy. ALBERTIANA thus encourages the publication of announcements, literature reviews, progress reports, preliminary notes etc. - i. e. those contributions in which information is presented relevant to current interdisciplinary Triassic research.

---

**Editor**

Dr. Wolfram M. Kürschner, Laboratory of Palaeobotany and Palynology, Utrecht University,  
Budapestlaan 4, 3584 CD Utrecht, The Netherlands, w.m.kurschner@bio.uu.nl;

**Editorial Committee**

Dr. Aymon Baud, Musée de Géologie, BFSH2-UNIL, 1015 Lausanne, Switzerland,  
aymon.baud@sst.unil.ch;

Prof. Dr. Hans Kerp, WWU, Abt. Palaeobotanik, Hindenburgplatz 57, 48143 Münster, Germany,  
kerp@uni-muenster.de;

Dr. Spencer G. Lucas, New Mexico Museum of Natural History, 1801 Mountain Road N. W., Albuquerque,  
NM 87104, USA, slucas@nmmnh.state.nm.us;

Dr. Mike Orchard, Geological Survey of Canada, 101-605 Robson Street, Vancouver, British Columbia,  
V6B 5J3, Canada, morchard@nrcan.gc.ca;

Dr. E. T. Tozer, Geological Survey of Canada, 101-605 Robson Street, Vancouver, British Columbia,  
V6B 5J3, Canada, etozer@nrcan.gc.ca;

Prof. Dr. Henk Visscher, Laboratory of Palaeobotany and Palynology, Utrecht University,  
Budapestlaan 4, 3584 CD Utrecht, The Netherlands, h.visscher@bio.uu.nl.

---

Cover: *Magnetostratigraphy of the PTB GSSP Meishan Section D (Episodes, 24 (2), p. 110, Yin, Zhang, Tong, Yang & Wu).*

## Executive Notes

### From the Chair

As this issue was going to press, I was informed that **the Shallow Tethys Conference is postponed** until next year. The STS meeting that was to be held in conjunction with this conference therefore no longer has a parent meeting through which proceedings, accommodations, meeting rooms etc. would have been arranged. Nevertheless, the STS executive and our Hungarian hosts have decided to proceed with an STS meeting in Hungary in early September. This will include field trips to potential GSSP sites in the Balaton Highlands and a full day of presentations and discussions, as announced below. As stated elsewhere in this issue, it was planned to make the session on 'Middle Triassic boundaries' the venue for comprehensive discussion of, and scheduling of decisions on, the O/A, A/L, and L/C boundaries. I therefore encourage all those with an interest in, or a contribution to make on these boundaries to attend the meeting. Under pressure from the IUGS/ICS, I have set a goal to come to a final vote on GSSP placement for each of these three boundaries by the end of 2003, starting with the A/L this year.

Because time is getting short and the organizers need as much lead time as possible to make suitable arrangements, **I ask that STS members immediately communicate their interest in attending the STS meeting. Please respond to [haas@ludens.elte.hu](mailto:haas@ludens.elte.hu) and copy to [morchard@nrcan.gc.ca](mailto:morchard@nrcan.gc.ca).** Limited funds may be available for those people whose attendance depends on a subsidy if they propose to contribute to the proceedings of the meeting. Those who seek help with registration, accommodation, or exceptionally with travel should apply to the Chair (M.J. Orchard) stating their exact needs.

Many of you will have learnt that my proposal for an IGCP project on **Triassic time and trans-Panthalassan correlation** has been accepted. This project, IGCP 467, will become a further vehicle for global Triassic studies and will augment STS activities as we push forward toward the establishment of an unambiguous global time scale. The project will focus on establishing linkages between faunal provinces ('trans-Panthalassa'), as well as between the marine and continental realms, leading to the development of a broadly applicable temporal framework and its application in reconstructing the Triassic world. It is therefore

anticipated that IGCP workshops and sponsorship will supplement future STS meetings, as will be the case with those announced herein.

**The GAC meeting in Vancouver, 26-28th, May 2003** will feature a full day session on "Extinction events, faunal turnovers, and natural boundaries within and around the Late Triassic." It will be co-sponsored by the STS and by both IGCP projects 458 (T-J boundary events) and 467, and is convened in order to set the stage for C/N and N/R boundary resolution. The subsequent field trip to the classic area of Williston Lake will take place if enough support is shown. We must quickly establish the number of likely participants on this field trip which first involves an hour flight to NE BC, followed by road and boat travel; costs are being worked out. **Please email the Chair ASAP if you are interested in participating in the Williston Lake field trip.** The Chinle field trip is now planned in conjunction with the **GSA meeting, Seattle, November 2003.**

We have recently learned that the **IGC 2004** organizing committee has formally accepted both a session on "Triassic in the Tethys Realm" and a workshop on "The Upper Triassic: definition, subdivision, and correlation" at the Florence meeting in August 2004. We hope to organize field trips before hand to view some Upper Triassic boundary candidates, and make this meeting the final forum for discussion on Upper Triassic GSSPs. **Plan to be there!**

#### Joint STS / IGCP 467 field meeting on Middle Triassic stage boundaries Veszprém, Hungary, September 5-8, 2002

##### Program

*September 5, Thursday*

Arrival at Budapest

Assembly of participants at the Hungarian Geological Institute (14 Stefánia út) (with opportunity to visit the building and its Geological Museum, and meet colleagues)

Departure for Veszprém at 6 pm.

*September 6, Friday*

Field trip to the Balaton Highland

Localities to be visited: Felsörs (Anisian/Ladinian GSSP candidate), Nosztori Valley (Ladinian/Carnian boundary section), Köveskál (Ladinian/Carnian boundary section)

*September 7, Saturday*

Scientific session

Business meeting of STS with discussion of GSSP issues

Business meeting of IGCP 467

*September 8, Sunday*

Half-day meeting continuing from previous day, if warranted

Departure for Budapest after lunch

### **Transportation**

Participants are responsible for making their own travel arrangements from their home country to and from Budapest. Transfer from Budapest to Veszprém (and back to Budapest) will be provided by rental bus or minibuses.

### **Conference venue**

Regional Center of the Hungarian Academy of Sciences, Veszprém (The building is located in the historic Veszprém Castle. The building has a large conference hall and a smaller meeting room, its own cafeteria, and guest rooms).

### **Accommodation**

There is a limited number of guest rooms available at the conference venue (double rooms). Additional accommodation as needed will be arranged in the dormitory of the Veszprém University, across the street from the conference venue.

### **Costs**

The following estimates are subject to change depending on the number of participants and final arrangements. Accommodation (3 nights): EURO 75 (per person, double occupancy, breakfast included) Registration fee (including transportation, field trip costs, guide books, meals starting with dinner on September 5 and ending with lunch on September 8): EURO 120

### **Organization**

The local organizing committee consists of J. Haas (Chair), T. Budai, S. Kovács, J. Pálffy, and A. Vörös. The meeting is organized by the Triassic Subcommittee of the Hungarian Stratigraphic Commission, the Hungarian Academy of Sciences, and the Hungarian Geological Society.

### **Pre-registration**

Please send an e-mail or fax expressing your interest and a preliminary title if you intend to give a presentation at the scientific session to the following address:

Dr. János Haas

Geological Research Group, Hungarian Academy of Sciences

Pázmány sétány 1/c, Budapest, H-1117 Hungary

Fax: +36 1 381-2128

Email: [haas@ludens.elte.hu](mailto:haas@ludens.elte.hu)

After canvassing the membership, both Maurizio and I have concluded that there is insufficient interest in a visit to the Italian Middle Triassic candidate sections and it is therefore not useful to organize a trip for the few who would participate; fortunately most of us have seen them. Similarly, an earlier meeting in June is unlikely to attract many who have already made plans for the summer. On the other hand, I assume that those STS members who were interested in participating in the original Shallow Tethys meeting are more likely to attend a meeting around the original date.

I am therefore proposing, as originally conceived, an STS meeting in Budapest with the focus on Middle Triassic boundaries, to include field excursions to Felsőors and also to Ladinian-Carnian localities as suggested by Sandor. There should be a full day of key presentations and discussion. Maurizio has suggested the itinerary given below, which appears good to me. Please consider it and let me know ASAP your view. Suggest modifications if you wish and let me have information about accommodation, meeting place, costs of field trip etc. so an announcement can be included in *Albertiana* (this must be given to Wolfram Kurschner by the 9th March). I will also distribute an email and alert the STS membership. You could in fact draft your own circular with a return-by date so that you have immediate notification of attendees for organizational purposes. Suggested itinerary:

September 5, Thu.: Arrival in Budapest.

September 6 Fri: visit to Felsőors, Koveskal and Nostory valley.

September 7 Sat: Full day of discussion on the two boundaries.

Sept. 8 Sun: Departure. (It is important to have the night Sat/Sun for the week end fare on the flights).

As far as the program is concerned, I would like to see time set aside for a review of each boundary (O-A/A-L/L-C) and pertinent presentations thereon. I hope that this can be accommodated in a single day, but I am not adverse to extending the presentations/discussions if it appears necessary (½ day Sunday?). The meeting announcement should include an invitation to present summary and/or new data from key areas. We know these would include at least Jozsef Palfy on the new radiometric dates from the A-L at Felsőors, Marco Balini on the L-C in Spiti, and I think Alda Nicora and I would be ready to summarize the conodonts from the O-A at Desli Caira. I would also like to summarize the conodont biostratigraphy and correlation potential about each boundary in North America. We may not fully resolve these boundaries

but I wish the status of all of them to be discussed and a schedule for decisions formulated.

Now that IGCP 467 is a reality, I would like to propose to make this meeting a formal workshop both for STS and IGCP. Some funds may be available from both sources to offset costs.

Looking forward to your final comments.

Mike Orchard, Chair STS (ICS, IUGS)

### **From the Secretary**

J. D. Campbell (1927-2001)

It was with great sadness that I learnt of the death of John Douglas ('Doug') Campbell, at Warrington, New Zealand, on 27 July 2001. Doug was a former member of the Subcommission on Triassic Stratigraphy (Albertiana 10 and 22) who was well known for his studies on brachiopod faunas and Triassic biostratigraphy. An Obituary and an appreciation of his work appeared in Newsletter 126 of the Geological Society of New Zealand (November 2001).

**Geoffrey Warrington, STS Secretary**

## Reports

### The new GSSP, base of the Triassic: some consequences

Aymon BAUD

Musee de Geologie, BFSH2-UNIL, 1015  
Lausanne Switzerland  
aymond.baud@sst.unil.ch

The base of the Triassic was proposed by Yin et al. (1996a) at Meishan (S China) section D, bed 27c with the FAD of the conodont *H. parvus*. This Global Stratotype Section and Point (GSSP) has been adopted by the International Commission on Stratigraphy (Yin et al., 2001). It has been unveiled by an opening Ceremony of the GSSP Monument, August 11 2001, in the Meishan Quarry, during the International Symposium on the Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events (10-13 August 2001, Changxing, China). Thanks to our Chinese colleagues for an impressive organization.

According to Yin et al. (1996b) in their proposals for global correlation based on ammonoids and conodonts, the Permian-Triassic boundary interval can be subdivided in 6 units (PTBU) from the lower Changhsingian (PTBU 1) to the upper Griesbachian (PTBU 6). The base of the Triassic corresponds to the base of PTBU 5.

The PTBU 4 corresponds to the interval separating the base of the boundary clay to the FAD of *parvus* (beds 25 to 27b in Meishan).

The upward shift of the boundary from the base of bed 25 proposed twenty years ago to the base of the bed 27c formally adopted in Meishan Quarry (shift which corresponds strictly to the PTBU4), plus recent discoveries, new datations and correlations bear numerous consequences. We will set out below some of them.

1. Following the discovery by C. Henderson of key conodonts in the Otto Fiord area (Ellesmere Island, Arctic Canada) and according to Henderson & Baud (1997), there is an overlap between the upper Changhsingian and the lower Griesbachian (*sensu* Tozer, 1967). It is why A. Baud, B. Beauchamp and C. Henderson are preparing a redefinition of the Griesbachian substage (abstract: Baud & Beauchamp, 2001).

2. Consequently, the *Otoceras concavum* and *borale* fauna of the Arctic ranging below the *Ophiceras* fauna, are latest Permian in age (partly PTBU 4) and older than the *Otoceras woodwardi* fauna of the Himalayas which co-occurs with *Ophiceras*(PTBU 5).

3. Consequently, due to the normal up to high sedimentation rate in the boreal area (Sverdrup Basin, E. Greenland, Barents Sea, Spitzberg, Verkoyansk area), the base of the Triassic has to be shifted upward by several to tens of meters. The Permian-Triassic boundary (Erathem boundary) occurs within a monotonous shaly facies of great thickness containing impoverished biota with ammonoids (*Otoceas*, *Glyptophiceras*, *Hypophiceras*, *Tompophiceras* and *Ophiceras*), bivalves (*Claraia*) and conodonts (*Hindeodus*, *Neogondolella*). The palynological boundary, with the appearance of the Triassic key palynospecies occurs in the latest Permian (Cirilli et al. 2001).

4. According to Chen and Komatsu (2001), the first *Claraia* zone or assemblage with *C. baoqingensis*, *C. griesbachi* and *C. bioni* appear in the latest Permian (PTBU 4).

5. The main extinction pulse (top of bed 24 in Meishan, Jin et al. 2000) with an age > 254My, occurs more than 1 million year before the GSSP (= 253My, new data from Mundil et al. 2001).

6. The so called « boundary clay » and the associated volcanic ashes event in Meishan do not occur at the boundary (GSSP) but below and before, in the Late Changhsingian (PTBU 4).

7. Except for the Tethys Himalaya (Spiti, Central Himalaya, Nepal), the well-known base of Triassic transgression is starting in the Late Changhsingian (PTBU 3 to PTBU 4).

8. The sharp negative shift of the carbon isotope profiles (Baud et al. 1989, 1996) occurs in the late Changhsingian (upper part of the PTBU 3 to the lower part of the PTBU4), before the boundary and not at the boundary. The GSSP occurs within a positive rebound following the main negative shift.

9. According to Zhu et al. (1999), the GSSP occurs within a reversal magnetic zone and does not coincide with the base or within a normal magnetic zone as largely accepted (review in Jin et al., 2000).

Part of the misinterpretation or wrong correlations of the boundary is due to the highly condensed nature of the Stratotype (see Baud, 1996) and the apparent short interval deduced from the nearness

of the Sequence boundary, the event boundary and the biochronological boundary (GSSP). But we have to remind that proximity (few centimeters) does not mean short time interval.

Measuring the thickness, the PTBU 4 is only twelve centimeters in Meishan, section D. Its duration is more than one million years (see point 5). The PTBU 4 is about 4m thick in Shangsi, 2,8m in Guryul Ravine (Kashmir), one meter in Salt Ranges (Pakistan), 0.6m in Abadeh and 0.4 to 4-6m in the Southern Alps. In the Arctic, with high sedimentation rates, the PTBU 4 has up to some hundred times the thickness of the Meishan one and up to ten times the thickness of the Shangsi one (Figure 1) and values are between 10 and 50m according to the area and sections.

High rate of sedimentation during this time interval can occur in continental area. We have to take care about this when fixing the boundary, which is not an event boundary as mass extinction nor a collapse of terrestrial ecosystems or a so called "fungal spike", nor a physical boundary as a reversal magnetic zone or abrupt facies change.

### References

- Baud, A. 1996: The Permian-Triassic Boundary: recent developments, discussion and proposals. *Albertiana* 18, p. 6-9.
- Baud, A., Atudorei, V. and Sharp, Z., 1996a. Late Permian and Early Triassic Evolution of the Northern Indian Margin: Carbon Isotope and Sequence Stratigraphy. *Geodinamica Acta* 9, p. 57-77.
- Baud, A., and Beauchamp, B., 2001, Proposals for the redefinition of the Griesbachian substage and for the base of the Triassic in the Arctic regions, in Yan, J. and Peng, Y. (eds), Proceedings of the International Symposium on the Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events, Changxing (China), University of Geosciences Press, p. 26-28.
- Baud, A., Holser, W.T. and Magaritz, M., 1989. Permian-Triassic of the Tethys: Carbon isotope studies. *Geol. Rundschau* 78, p. 1-25.
- Chen, J., and Komatsu, T., 2001, Bivalves from the beds near the Permian-Triassic Boundary at Meishan section of Changxing, south China and the correlation with shallow marine bivalve fauna, in Yan, J., and Peng, Y., editors, Proceedings of the International Symposium on the Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events: Changxing (China), China University of Geosciences Press, p. 37-39.
- Cirilli, S., Spina, M. and Baud, A., 2001. Palynology of the uppermost Permian - basal Triassic successions in the High Arctic (Canada) and comparison with some PTB Gondwanian localities, Conference on the Geology of Oman, Abstract Book, Muscat, Oman, p.28.
- Henderson, C., and Baud, A., 1997, Correlation of the Permian-Triassic boundary in Arctic Canada and comparison with Meishan, China, in W. Naiwen, and J. Remane, (eds.), *Stratigraphy: Proceedings of the 30th IGC*, v. 11: Beijing, VSP, p. 143-152.
- Jin, Y., Shang, Q. and Cao, C., 2000. Late Permian magnetostratigraphy and its global correlation. *Chinese Science Bulletin*, 45(8): p. 698-705.
- Jin, Y. G., Wang, Y., Wang, W., Shang, Q. H., Cao, C. Q., and Erwin, D. H., 2000, Pattern of marine mass extinction near the Permian-Triassic boundary in South China: *Science*, 289, p. 432-436.
- Mundil, R., Metcalfe, I., Ludwig, K. R., Renne, P. R., Oberli, F., and Nicoll, R. S., 2001, Timing of the Permian-Triassic biotic crisis: implications from new zircon U/Pb age data (and their limitations): *Earth & Planetary Science Letters*, 187, p. 131-145.
- Tozer, E. T., 1967, A standard for Triassic time. *Bulletin - Geological Survey of Canada*, 156, 103pp.
- Yin, H., Sweet, W. C., Glenister, B. F., Kotlyar, G. V., Kozur, H., Newell, N. D., Sheng J., Yang, Z., and Zakharov, Y., 1996a, Recommendation of the Meishan section as the global stratotype and point (GSSP) for the basal Boundary of Triassic System. *Newsletter in Stratigraphy*, 34, p. 81-108.
- Yin, H., Wu, S., Ding, M., Zhang, K., Tong, J., Yang, F. and Dulong, L., 1996b, The Meishan section. Candidate of the global stratotype and point (GSSP) of the Permian-Triassic boundary (PTB), in H. Yin, (eds.), *The Paleozoic-Mesozoic Boundary. Candidates of the Global Stratotype Section and Point of the Permian-Triassic Boundary: Wuhan, China University of Geosciences Press*, p. 31-47.
- Yin, H., Zhang, K., Tong, J., Yang, Z. and Wu, S., 2001. The Global Stratotype Section and Point (GSSP) of the Permian-Triassic Boundary. *Epi-sodes*, 24(2): 102-114.

Zhu, Y. and Liu, Y., 1999. Magnetostratigraphy of the Permo-Triassic Boundary Section at Meishan, Changxing, Zhejiang Province. In: H. Yin and J. Tong (eds.), Proceedings of the International Conference on Pangea and the Paleozoic-Mesozoic Transition. China University of Geosciences Press, Wuhan (China), p. 79-84.



## Functional Importance of New Skeletal Elements («Eye Capsules») of Euconodonts

<sup>1,3</sup>Galina I. Buryi, and <sup>2</sup>Alla P. Kasatkina

<sup>1</sup>Federal Far Eastern Geological Institute, Far Eastern Branch, Russian Academy of Sciences, Stoletiya Prospect 159, Vladivostok-22, 690022; <sup>2</sup>Pacific Oceanological Institute, Far Eastern Branch, Russian Academy of Sciences, 43 Baltiyskaya st., Vladivostok, 690041, Russia  
<sup>3</sup>corresp. author: buryi@mail.ru

New skeletal elements of euconodonts have been found in the Lower Triassic deposits of South Primorye. The finding is confined to the siltstone mass (50 m) referred to the *Anasibirites nevolini* ammonite Zone (Buryi, 1979, p. 91) cropped out in the Kamenushka river basin (Perevalnyi creek) (Fig. 1).


In the upper parts of this mass, within the lens of calcareous sandstone, overfull with ammonites *Anasibirites nevolini* Buriy and Zharnikova, *Meekoceras subcratum* Kiparisova, *M. boreale* Diener, *Arctoceras robinsoni* (Kiparisova),

*Prospingitoides ovalis* (Kiparisova), *Juvenites sinuosus* (Kiparisova), abundant conodonts were found (sample 4035): platform - elements of *Neogondolella milleri* (Muller) - 22 specimens, segminate - elements of *Neospathodus lanceolatus* Mosher - 1 specimen, *Smithodus discreta* (Muller) - 19 specimens, *S. csritagali* (Huckriede) - 1 specimen, *S. concervativa* (Muller) - 4 specimens and ramiform elements of *Furnishius triserratus* Clark - 170 specimen, *Ellissonia triassica* Muller - 9 specimens, *E. meissneri* (Tatge) - 3 specimens, *E. magnidentata* (Tatge) - 4 specimens, *E. nevadensis* Muller - 3 specimens, *Hindeodella triassica* Muller - 36 specimens, *H. nevadensis* Muller - 24 specimens, *H. raridenticulata* Muller - 15 specimens, *H. budurovi* Buryi - 7 specimens, *Parachirognathus symmetrica* (Staesche) - 4 specimens, *P. inclinata* Staesche - *Hadrodentina adunca* Staesche - 13 specimens, *H. subsymmetrica* (Muller) - 36 specimens, *H. symmetrica* (Staesche) - 30 specimens, *Xaniognathus curvatus* Sweet - 2 specimens.

Together with these conodont elements, 3 specimens of new skeletal elements analogous to them in color (light brown transparent to deep brown) and in the surface structure. By analogy with P, M, and S conodont elements, we propose to identify new skeletal elements as H elements.

### Description of skeletal elements

Several varieties of H elements (Fig. 2 - 4) have



**Figure 1.** Location of Lower Triassic H elements in the Kamenushka River basin (South Primorye).

been found: H1, H2 and H3 that occur jointly. The tern H elements are proposed for the new skeletal elements by analogy with P, M, and S conodont elements.

**H1 elements** («bagel»-shaped [Russian baranka] (Fig. 2). The flattened plate shape with a hole on the center, resembling a bagel, is irregular-rounded. Its longer diameter is 0.72 mm and its shorter one is 0.57 mm. The thickness of the plate along the outer margin is about 0.1 mm. The outer margin of the plate is uneven and has abundant projections. At a distance of 0,2 mm from the outer margin, of this plate, nearly parallel to it, an inner margin extends. It is observed to project its surface like a border, a narrow crest bearing hardly visible projections or papillae. We suggest that the muscles, maintaining and governing the conodont elements of the mouth apparatus, were attached that to these papillae on the inner margin, as well as to the projections of the margins of the plate. The space between the outer and inner margins of the plate forms a sort of outer ring characterizing the H1 element.

From the inner margin towards the center, we distinguish its inner ring. In the H1 element, the surface of the outer ring is smooth and lustrous. The element has a crystalline structure similar to that of most of the conodont elements. The surface of the inner ring is composed of uneven, structureless, material, the thickness of which decreases gradually down to the nearly triangular hole in the center. The hole is about 0.3 mm long and its most widened part is 0.09 mm. The opposite lower surface of the plate is smooth. We suggest that it was faced inward the pharynx and had no attaching function.

**H2 element** (Fig. 3). H2 element is very similar to H1 element, only its outer ring is narrower and the inner ring is absent. The outer ring is hollow and looks like a hollow semisphere, the walls of which are very thin and almost transparent. Tubercular projections of the outer margin, are pronounced (the specimen was broken with a needle during inspection and so it consists of two semi rings).

**H3 element** (Fig. 4). The third variety of H element is plate-like with no hole in the center. The oval-leaf plate has a maximal diameter 0.87 mm and a shorter one of 0.65 mm. The thickness of this plate is about 0.1 mm.

It has an outer ring, but the projections of its outer margin are less pronounced than in H1 element. The space of the inner ring, having no hole, is filled with four prominent elongated projections, which

start from a common pain on the inner margin of the outer ring. Two central projections extend farther to the opposite margin of the outer ring, and two lateral projections come to the middle of the outer ring area. Such arrangement of the projections of the inner ring in H3 element resembles the arrangement of veins in a tree leaf. In contrast to H1 and H2 elements, the surface of the outer ring in H3 element is rough whereas the surface of the projections of the inner ring is smooth and listrous.

## Discussion

We suggest that H skeletal elements, described above, were the parts of the euconodont head complex, which is composed of a mouth apparatus and pair skeletal elements of a head.

The mouth apparatus of euconodont is known to be of different modifications of P, M, and S conodont elements (Aldridge et al., 1995; Purnell et al., 1995; Purnell and Donoghue, 1997) in different proportions. Using them, the mouth apparatus provides catching of a prey and feeding of an animal and favours water filtering through the branched tooth net as well as crushing of the filtered prey with the help of more massive platform elements. The mouth apparatus appears to be a rather strong instrument, the work of which needs a strong muscular system. In our opinion, the place of attachment of muscles, joining isolated conodont elements into the mouth apparatus and governing these elements, was a pair of the head attaching plates or H elements.

Thus, we think that in the head part of euconodont there was a single mouth complex composed, probably, of a pharynx, about 15 P, M, and S conodont elements, muscles, and two head attaching plates (H elements) necessary for joining and governing the conodont elements in the mouth apparatus.

In Recent Chaetognatha animals, similar to euconodonts, chitinous setae and crenations of the prehensile apparatus are attached to the body also with two prolate semi-transparent plates arranged on the ventral and dorso-ventral sides of the head (Kasatkina, 1982; Kasatkina and Buryi, 1996, 1997).

There is another interpretation of the functional importance of these structures. In the head parts of the euconodont animal imprints reported from the Carboniferous Granton bed of Edinburg, Scotland, and the Upper Ordovician Soom Shale of South Africa, there were observed lobate structures, similar to H elements, which were suggested to be «eye capsules» (Aldridge et al., 1993; Aldridge

and Theron, 1993).

Thus, in spite of the fact that Kasatkina found a now-living conodont animal - "alive fossil" At the depth of 1992-1993 m in the sea of Laptevkh of Arctic Siberia (Kasatkina, 2000), the physiological function of these skeletal plates remains still unclear.

In conclusion we'd like to ask all specialists, who study conodont elements to pay attention to these H elements in their samples.

### Acknowledgements

We thank Dr. Kirill Budurov, who in 1974, looking through our collection of conodont elements from South Primorye, found H2 element described in this paper and attract our attention to a great importance of this finding for understanding the structure of a conodont animal. The work was made under the financial support of RFBR grant and 01-04-49416 (Revision of Euconodonta (Conodonta) mouth apparatus features on the basis of the recent euconodont organism original findings and Chaetognatha morphological peculiarities).01-05-64599 ("Palaeontological basis of the *Anasibirites nevolini* and *Tirolites* Zones of the Olenekian in South Primorye and global correlation of the Lower Triassic".

### References

Aldridge, R.J., Briggs, D.E.G., Smith, M.P., Clarkson, E.N.K. & Clark, N.D.L., 1993. The anatomy of conodonts. *Philosophic Transactions of Royal Society, B* 340: 405-21, London.

Aldridge, R.J. & Theron, J.N., 1993. Conodonts with preserved soft tissue from a new Ordovician Konservat-Lagerstätte. *Journal of Micropalaeontol.*, 12: 113-7.

Aldridge, R.J., Purnell, M.A., Gabbot, S.E. & Theron, N.N., 1995. The apparatus architecture and function of *Promissium pulchrum* Kovacs-Endrody (Conodonts, Upper Ordovician) and the prionodontid plan. *Philosophic Transactions of Royal Society, B* 347: 275-91, London.

Buryi, G.I., 1979. Lower Triassic conodonts in the Southan Primorye. Nauka, Moscow, 144 pp. (in Russian).

Kasatkina, A.P., 1982. Chaetognaths in the of the Sea USSR and adjacent Waters. Nauka, Leningrad 136 pp. (in Russian).


Kasatkina, A.P., 2000. Alive fossils of conodonta (Euconodontophyles). *Doklady Akademii Nauk. General Biology*, 375(5): 710-713 (in Russian).

Kasatkina, A.P. & Buryi, G.I., 1996. On the relation of chaetognaths and conodonts. *Albertiana*. 18: 21-23, Münster.



Kasatkina, A.P. & Buryi, G.I., 1997. Chaetodonta: a new animal superphylum and its position in animal systematics, *Doklady Biological Sciences, Moscow*, 356: 503-505.

Purnell, M.A., Aldridge, R.J., Donoghue, P.C.J. & Gabbot, S.E., 1995. Conodonts and the first vertebrates, *Endeavour, New Series*, 19(1): 20-27, Oxford.


Purnell, M.A. & Donoghue, P.C.J., 1997. Architecture and functional morphology of the skeletal apparatus of ozarkodinid conodonts. *Philosophic Transactions of Royal Society, B* 352: 1545-64, London.



**Figure 2.** Euconodont  $H_1$  element (bagel) - DVGI 3B-1 x 300. Lower Triassic, Olenekian, *Anasibirites nevolini* Zone. Om - outer margin; p - projection; im - inner margin (border with papillae); or - outer ring; ir - inner ring; h - hole.



**Figure 3a,b** Euconodont  $H_2$  element (two semi-rings); a - DVGI 3B-2, x 300. b - DVGI 3B-3, x 300. Lower Triassic, Olenekian, *Anasibirites nevolini* Zone.



**Figure 4.** Euconodont  $H_3$  element - DVGI 3 B-4. Lower Triassic, Olenekian *Anasibirites nevolini* Zone.

## Age and Correlation of Triassic Tetrapod Assemblages from Brazil

Spencer G. Lucas

*New Mexico Museum of Natural History,  
1801 Mountain Road N.W., Albuquerque,  
New Mexico 87104-1375, USA  
slucas@nmmnh.state.nm.us*

Triassic tetrapod fossils from the state of Rio Grande do Sul, Brazil, come from three lithostratigraphic units—Sanga do Cabral, Santa Maria and Caturrita formations—in the Rosário do Sul Group. No detailed lithostratigraphic framework of the Brazilian Triassic tetrapod localities has been developed, and may be impossible, given the heavily vegetated landscape of Rio Grande do Sul, where outcrops are intermittent, many are man made (road cuts, quarries, etc.), and few exposures encompass more than several meters of stratigraphic section. Three distinct biostratigraphic assemblages of Triassic tetrapods can be recognized: (1) Lootsbergian tetrapods from the Sanga do Cabral Formation; (2) the Berdyankian (Chanarian) *Dinodontosaurus* Assemblage Zone of the middle part of the Alemoa Member of the Santa Maria Formation; and (3) the Adamanian (Ischigualastian) *Hyperodapedon* Assemblage Zone of the upper part of the Alemoa Member and the Caturrita Formation. Recent attempts to refine this biostratigraphy by recognizing more subdivisions are laudable but lack convincing support from lithostratigraphic or biostratigraphic data.

### Introduction

Triassic tetrapod fossils from Brazil come from the state of Rio Grande do Sul (Fig. 1) in strata exposed on the southern flank of the Paraná basin and have been collected for nearly a century (e.g., Woodward, 1907; Huene, 1935-1942; Beltrão, 1965; Barberena, 1977; Barberena et al., 1985a; Schultz et al., 2000; Abdala et al., 2001). These tetrapod fossils come from three lithostratigraphic units—Sanga do Cabral, Santa Maria and Caturrita formations—in the Rosário do Sul Group (e.g., Andreis et al., 1980; Barberena et al., 1985a; Faccini et al., 1995, 2000) (Fig. 1). Much has been written about the age and correlation of these tetrapods, especially since the 1970s (Barberena, 1977; Barberena et al., 1985a, b; Schultz et al., 1994, 1998, 2000; Scherer et al., 1995; Schultz, 1995; Abdala et al., 2001). Based on my studies of collections in Brazil, and an examination in the field

of some key localities, I present here an assessment of the age and correlation of the Brazilian Triassic tetrapods.

### Stratigraphic Context

As Faccini et al. (2000) state, the Sanga do Cabral, Santa Maria and Caturrita formations (Fig. 1) are mostly fluvial conglomerates, sandstones and mudstones that were deposited in a large intracratonic basin in what was southwestern Gondwana, the Paraná basin. The relatively thin (up to 20 m thick) Sanga do Cabral Formation is fluvial and eolian sandstones disconformably overlain by fluvial sandstone and conglomerate of the Passo das Tropas Member (~20-30 m thick) of the Santa Maria Formation. The overlying Alemoa Member of the Santa Maria Formation (~100-120 m thick) is mostly floodplain mudrocks and minor fluvial channel sandstones. The Caturrita Formation (~60 m thick) is mostly sandy and gravelly river channel deposits. Triassic tetrapod assemblages of biostratigraphic significance come from the middle part of the Sanga do Cabral Formation, the middle and upper part of the Alemoa Member of the Santa Maria Formation and from the Caturrita Formation (Fig. 2).

### Previous Schemes of Tetrapod Biostratigraphy

Various schemes of tetrapod biostratigraphy have been proposed for the Brazilian Triassic (Fig. 2). It is essential to understand that no detailed lithostratigraphic framework of the Brazilian Triassic tetrapod localities has been developed. Indeed, such stratigraphic organization may be impossible, given the heavily vegetated landscape of Rio Grande do Sul, where outcrops are intermittent, many are man made (road cuts, quarries, etc.), and few exposures encompass more than several meters of stratigraphic section. Of course, it is possible to order grossly the Triassic tetrapod localities in Rio Grande do Sul by assigning them to formations and members, but a more detailed stratigraphic ordering remains to be demonstrated. This, however, has not hindered some workers (e.g., Schultz, 1995; Abdala et al., 2001) from proposing a temporal ordering of fossil assemblages that lacks detailed support from stratigraphic data.

Also critical to tetrapod biostratigraphy in the Brazilian Triassic is its correlation to the Argentinian Triassic tetrapod assemblages, which can be ordered in a precise lithostratigraphic framework (e.g., Bonaparte, 1966, 1967, 1969, 1970, 1982; Rogers et al., 1993). The Middle-Late Triassic Argentinian assemblages are the basis of provincial




Figure 1: Location of Rio Grande do Sul and its Paleozoic-Mesozoic (Gondwana) strata on the southern flank of the Paraná basin, and summary of Triassic stratigraphy in that basin.

land-vertebrate faunachrons (lvfs), the Chanarian, Ischigualastian and Coloradan, and can be correlated to a global tetrapod biochronology (Lucas, 1998).

Barberena (1977) is the point of departure for all detailed correlation of the Brazilian Triassic tetrapod assemblages. He recognized two tetrapod assemblage zones in the Santa Maria Formation (Fig. 2), correlating the lower, Therapsida Assemblage Zone to the Argentinian Chanarian and the younger, Rhynchocephalia Assemblage Zone to the Ischigualastian. Barberena et al. (1985a) renamed these zones (Fig. 2), but advocated the same correlations. Barberena et al. (1985b; also see Holz & Barberena, 1994) followed suit, but also stressed the identification of four, temporally successive local faunas (Fig. 2). However, as noted by Lucas (1993), no stratigraphic data exist to indicate the Pinheiros and Chiniquá local faunas (= tetrapod assemblages from those areas) are actually temporally successive.

Schwanke (1998) suggested that there are three successive dicynodont assemblages in the Santa Maria Formation, a lower assemblage with *Chanaria*, *Barysoma*, *Dinodontosaurus* and *Ischigualastia* immediately overlain by an assemblage with *Stahleckeria* and *Dinodontosaurus*, and a third, much younger assemblage with *Jachaleria*. Schultz et al. (1994, 1998, 2000) and Scherer et al. (1995) essentially reiterated the zonation of Barberena and co-workers but renamed the zones

as “cenozones” (Fig. 2). However, they did distinguish the stratigraphically highest tetrapod assemblage, from the Caturrita Formation, as the “*Jachaleria* interval.” At the same time, though, Schultz (1995) proposed a succession of “associations” (Fig. 2) that, nevertheless, lacked any detailed stratigraphic data to support its temporal ordering. Most recently, Abdala et al. (2001) proposed a slightly modified zonation, adding a Traversodontid Biozone between the *Dinodontosaurus* and Rhynchosaur Biozones (Fig. 2). Nevertheless, the “Traversodontid Biozone” is based on a single locality (Santa Cruz do Sul) that cannot be ordered stratigraphically with respect to other localities.

### Sanga do Cabral Formation

The oldest Triassic tetrapods from Brazil are from the middle part of the Sanga do Cabral Formation, especially in the vicinity of Catuçaba. The assemblage includes a rhytidosteid amphibian, *Procolophon*, other procolophonid specimens, a protorosaurid vertebra, skull material and an ulna of a pareiasaur and a dicynodont stapes (Lavina, 1983; Barberena et al., 1985a; Dias-da-Silva, 1997, 1998; Dias-da-Silva & Schultz, 1999; Schultz & Dias-da-Silva, 1999; Schwanke & Kellner, 1999). Brazilian workers (e.g., Andreis et al., 1980; Barberena et al., 1985a; Faccini et al., 1995; Kellner & Campos, 1999) have long, and, I believe correctly, assigned these tetrapods an Early Triassic age; they are most reasonably (based primarily on




Figure 2: Previous schemes of Triassic tetrapod biostratigraphy of the Santa Maria and Caturrita formations compared with the scheme advocated here.

the presence of *Procolophon*) correlated to the *Lystrosaurus* “zone” of the South African Karoo and thus are of Lootsbergian age *sensu* Lucas (1998). The pareiasaur record here is particularly interest-

ing, as it joins *Sclerosaurus* from the Buntsandstein of the German-Swiss borderland as the only possible Triassic records of pareiasaurs.

## Santa Maria Formation

The most extensive Triassic tetrapod assemblages from Brazil come from the Santa Maria Formation and include fossils of procolophonids, rhynchosaurs, proterochampsids, raiisuchids, aetosaurs, crocodylomorphs, dinosaurs, dicynodonts and cynodonts (e.g., Woodward, 1907; Huene, 1935-1942; Romer & Price, 1944; Price, 1947; Cox, 1965; Colbert, 1970; Barberena & Daemon, 1974; Bonaparte & Barberena, 1975; Araújo & Gonzaga, 1980; Araújo, 1981; Barberena, 1982; Zacarias, 1982; Texeira, 1982; Barberena et al., 1985a; Mattar, 1987; Azevedo et al., 1990; Azevedo, 1995; Kellner, 1998; Kellner & Campos, 1999, 2000; Abdala & Ribeiro, 2000; Langer & Schultz, 2000; Abdala et al., 2001; Lucas & Heckert, 2001). These assemblages are usually, and I believe correctly, considered to be Middle to Late Triassic age and correlated to the Chanarian and Ischigualastian lvs in Argentina (e.g., Barberena, 1977; Barberena et al., 1985a, b; Schultz et al., 1994, 1998, 2000; Scherer et al., 1995; Schultz, 1995; Abdala et al., 2001).

The stratigraphic data (or lack thereof) and the composition of the tetrapod assemblages of the Santa Maria Formation support recognition of two, temporally successive tetrapod faunas separated by a substantial hiatus (Fig. 2). This was the conclusion advocated first by Barberena (1977), and followed by those who have attempted a global correlation of Triassic tetrapod assemblages (e.g., Ochev & Shishkin, 1989; Lucas, 1998, 1999). Recent attempts to refine this correlation by recognizing more subdivisions (e.g., Schultz, 1995; Abdala et al., 2001) are laudable, but they lack support from lithostratigraphic or biostratigraphic data.

### Dinodontosaurus Assemblage Zone

The middle part of the Alemoa Member of the Santa Maria Formation yields a tetrapod assemblage dominated by the dicynodont *Dinodontosaurus* (Candelaria and Chiniquã local faunas of Barberena, 1977). Besides *Dinodontosaurus* (= *Chanaria*), key elements of the assemblage include the dicynodont *Stahleckeria* (= *Barysoma*) and the traversodontid cynodonts *Traversodon*, *Belesodon* and *Massetognathus*. The assemblage of tetrapods from the middle part of the Alemoa Member is readily correlated with the Chañares local fauna of the Ischichuca Formation in Argentina. These assemblages of the Chanarian lvs are generally considered to be of Ladinian age, although the basis of this correlation is tenuous (e.g., Lucas & Harris, 1996). The cynodont-dominated fossil lo-

cality in the Alemoa Member at Santa Cruz do Sul (Abdala et al., 2001) is not demonstrably younger than other localities in this assemblage zone, so I see no basis for a "Traversodontid Biozone" younger than the *Dinodontosaurus* Assemblage Zone.

Schwanke Peruzzo (1990) and Schwanke Peruzzo & Araújo-Barberena (1995) assigned a partial skull from near Candelaria, from part of the Pinheiros local fauna, to the dicynodont *Ischigualastia*. This created an apparent range extension of the genus *Ischigualastia*, known only from Ischigualastian age strata in Argentina, into Chanarian age strata in Brazil. However, I have restudied this specimen, and it is *Stahleckeria*.

### Hyperodapedon Assemblage

The Upper Triassic vertebrate assemblage from the upper part of the Alemoa Member of the Santa Maria Formation is mostly from the vicinity of Santa Maria City. This is the Rhynchocephalia Assemblage Zone of Barberena (1977) or the *Scaphonyx* Zone of Barberena et al. (1985a). As the rhynchosaurs in this assemblage are dominantly *Hyperodapedon*, not *Scaphonyx* (Langer & Schultz, 2000), I rename the zone the *Hyperodapedon* Assemblage Zone (Fig. 2).

The *Hyperodapedon* Assemblage Zone in the Alemoa Member includes abundant fossils of the rhynchosaur *Hyperodapedon* (formerly *Scaphonyx*); the aetosaur *Stagonolepis* (formerly *Aetosauroides*); less abundant traversodontids and other cynodonts, including *Charrudon*, *Therioherpeton* and *Gomphodontosuchus*; the proterochampsids *Cerritosaurus binsfeldi*, *Rhadinosuchus gracilis*, and *Hoplitosuchus raii*; and the archetypal raiisuchian *Raiisuchus tiradentes*. Alemoa Member dinosaurs are the theropod *Staurikosaurus pricei* Colbert, 1970, the prosauropod *Saturnalia tupiniquim* Langer, Abdala, Richter & Benton, 1999 (also see Kellner & Campos, 2000), and the theropod *Teyuwasu barberenai* Kischlat, 1999.

In this assemblage zone most, if not all, of the rhynchosaurs, long referred to *Scaphonyx*, are now assigned to *Hyperodapedon*, a genus known from Otischalkian-Adamanian strata in India and Great Britain (Hunt & Lucas, 1991; Langer & Schultz, 2000; Langer et al., 2000). Argentinian material of *Hyperodapedon* comes from the Ischigualasto Formation (e.g., Contreras, 1999). The aetosaur *Stagonolepis* is also of well documented Adamanian age in the USA and Europe and is abundant in the Ischigualasto Formation in Argen-



tina (Heckert & Lucas, 2000; Lucas & Heckert, 2001). Clearly, the presence of *Hyperodapedon* and *Stagonolepis* supports correlation of the *Hyperodapedon* Assemblage Zone of the Santa Maria Formation with the vertebrates of the Ischigualasto Formation in Argentina, and therefore an Ischigualastian (Adamian) age assignment (Lucas, 1998; Lucas & Heckert, 2001).

### Caturrita Formation

The tetrapod assemblage from the Caturrita Formation includes a sphenodont skull and postcrania, the proterochampsid *Proterochampsia nodosa*, archosaur teeth, phytosaur teeth and jaw fragments, the cynodont *Exaeretodon*, the rhynchosaur *Hyperodapedon*, the dicynodont *Ischigualastia* (= *Jachaleria candelariensis* Araújo & Gonzaga) and a supposed *Erythrotherium*-like mammalian mandible fragment (Araújo & Gonzaga, 1980; Barberena et al., 1985a; Dornelles, 1990; Ferigolo, 1999; Bonaparte & Ribeiro, 1999; Faccini et al., 2000). Caturrita Formation dinosaurs are the theropod? *Guaibasaurus candelarai* Bonaparte, Ferigolo & Ribeiro, 1999 and a new, undescribed prosauropod (Azevedo, 1993; Azevedo et al., 1998, 1999). Lucas (1993) suggested that the skull assigned to *Jachaleria* belongs to *Ischigualastia*, and I maintain this conclusion, having now studied the Brazilian material firsthand.

The Caturrita assemblage thus shares index taxa with the Ischigualasto Formation of Argentina (*Ischigualastia*, *Exaeretodon* and *Hyperodapedon*), so I also assign it an Ischigualastian age. I therefore reject correlations, such as Bonaparte (1982), Barberena et al. (1985a) and Schultz et al. (2000), that indicate that at least part of the Caturrita Formation is younger than the Ischigualastian. All Late Triassic tetrapods known from Brazil are of Ischigualastian age (Fig. 2).

Certainly the Caturrita Formation tetrapods are stratigraphically above those from the upper part of the Alemoa Member of the Santa Maria Formation. However, the two assemblages are not, at present, biochronologically separable. Thus, both are of Ischigualastian age, and I include both in the *Hyperodapedon* Assemblage Zone (Fig. 2). More collecting and study of Caturrita Formation tetrapods are needed to provide a basis for recognizing them as a biochronologically distinct assemblage.

### Acknowledgments

I am grateful to numerous colleagues in Brazil for access to collections, other data and helpful discussions, especially Fernando Abdala, Sergio

Dias-da-Silva, Alex Kellner, Edio Kischlat, Martha Richter, Cesar Schultz, Cibele Schwanke and Cristina Vega. Andrew Heckert and Kate Zeigler provided helpful comments on the manuscript.

### References

- Abdala, F. & Ribeiro, A. M. 2000. A new therioherpetid cynodont from the Santa Maria Formation (middle Late Triassic), southern Brazil. *Geodiversitas*, 22: 589-596.
- Abdala, F. Ribeiro, A. M. & Schultz, C. L. 2001. A rich cynodont fauna of Santa Cruz do Sul, Santa Maria Formation (Middle-Late Triassic), southern Brazil. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, in press.
- Andreis, R. R., Bossi, G. E. & Montardo, D. K. 1980. O Grupo Rosário do Sul (Triássico) no Rio Grande do Sul. *Anais do 31 Congresso de Sociedade Brasileira de Geologie*, 2: 659-673.
- Araújo, D. C. 1981. Sobre a presença de *Chanaria* na Formação Santa Maria (Triássico do Brasil). *Anais II Congresso Latino-Americano de Paleontologia*, Porto Alegre: 307-319.
- Araújo, D. C. & Gonzaga, T. D. 1980. Uma nova espécie de *Jachaleria* (Therapsida, Dicynodonta) do Triássico de Brasil. *Actas II Congreso de Paleontología y Bioestratigrafía y I Congreso Latinoamericano de Paleontología*, Buenos Aires, 1: 159-174.
- Azevedo, S. A. K. 1993. Os dinossauros Triássicos no sul do Brasil: Dados e perspectivas. *Acta Geologica Leopoldensia*, 16: 31-40.
- Azevedo, S. A. K. 1995. Paleoeecologia de *Prestosuchus chinquensis* Huene, 1942. Inter-relações paleoecológicas entre as faunas reptilianas do Triássico Sul-Brasileiro. *Anais da Academia Brasileira de Ciências*, 67: 55-59.
- Azevedo, S. A. K., Schultz, C. L. & Barberena, M. C. 1990. Novas evidências bioestratigráficas e paleoecológicas na análise da evolução explosiva dos rincossauros do Triássico. *Paulo-Coutiana*, 4: 23-33.
- Azevedo, S. A. K., Da Rosa, A. A., Boelter, R. A. & Leal, L. A. 1998. A prosauropod dinosaur from the Late Triassic of southern Brazil. *Simpósio Brasileiro de Paleontologia de Vertebratas*, *Boletim de Resumos*: 14.

- Azevedo, S. A. K., Da Rosa, A. A., Boelter, R. A. & Leal, L. A. 1999. A prosauropod dinosaur from the Late Triassic of southern Brazil. *Paleontologia em Destaque*, 26: 59.
- Barberena, M. C. 1977. Bioestratigrafia preliminar da Formação Santa Maria. *Pesquisas, Porto Alegre*, 7: 111-129.
- Barberena, M. C. 1982. Uma nova espécie de *Proterochampsia* (*P. nodosa*, sp. nov.) do Triássico do Brasil. *Anais da Academia Brasileira de Ciências*, 54: 127-141.
- Barberena, M. C. & Daemon, R. F. 1974. A primeira ocorrência de Amphibia (Labyrinthodontia) na Formação Rio do Rasto. Implicações geocronológicas e estratigráficas. *Anais do XXVII Congresso Brasileiro de Geologia, Porto Alegre*, 2: 251-261.
- Barberena, M. C., Araújo, D. C. & Lavina, E. L. 1985a. Late Permian and Triassic tetrapods of southern Brazil. *National Geographic Research*, 1: 5-20.
- Barberena, M. C., Araújo, D. C., Lavina, E. L. & Azevedo, S. A. 1985b. O estado atual do conhecimento sobre os tetrápodes Permianos e Triássicos do Brasil Meridional. *Coletânea de Trabalhos Paleontológicos, MME-DNPM, Série Geologia, Seção de Paleontologia e Estratigrafia*, 27: 21-28.
- Beltrão, R. 1965. Paleontologia de Santa Maria e São Pedro do Sul, Rio Grande do Sul, Brasil. *Boletim do Instituto de Ciências Naturais da Universidade Federal de Santa Maria*, 2: 1-156.
- Bonaparte, J. F. 1966. Chronological survey of the tetrapod-bearing Triassic of Argentina. *Breviora*, 251: 1-13.
- Bonaparte, J. F. 1967. Cronología de algunas formaciones Triásicas de Argentina: Basada en restos de tetrápodos. *Asociación Geológica de Argentina Revista*, 21: 20-38.
- Bonaparte, J. F. 1969. Datos sobre la evolución paleoecología en las formaciones Triásicas de Ischigualasto-Villa Unión (San Juan-La Rioja). *Acta Geologica Lilloana*, 10: 189-206.
- Bonaparte, J. F. 1970. Annotated list of the South American Triassic tetrapods. *Second Gondwana Symposium Proceedings and Papers*: 665-682.
- Bonaparte, J. F. 1982. Faunal replacement in the Triassic of South America. *Journal of Vertebrate Paleontology*, 2: 362-371.
- Bonaparte, J. F. & Barberena, M. C. 1975. A possible mammalian ancestor from the Middle Triassic of Brazil (Therapsida-Cynodontia). *Journal of Paleontology*, 49: 931-936.
- Bonaparte, J. F. & Ribeiro, A. M. 1999. Novos vertebrados no Triássico Superior do RS, Brasil. *Paleontologia em Destaque*, 26: 60.
- Bonaparte, J. F., Ferigolo, J. & Ribeiro, A. M. 1999. A new early Late Triassic saurischian dinosaur from Rio Grande do Sul State, Brazil. *National Science Museum Tokyo Monograph*, 15: 89-109.
- Colbert, E. H. 1970. A saurischian dinosaur from the Late Triassic of Brazil. *American Museum Novitates*, 2405: 1-39.
- Contreras, V. H. 1999. Rhynchosaurs from the Ischigualasto Formation (Upper Triassic, late Carnian), San Juan, Argentina: VII International Symposium on Mesozoic Terrestrial Ecosystems, Buenos Aires, Argentina, Abstracts: 19.
- Cox, C. B. 1965. New Triassic dicynodonts from South America, their origins and relationships. *Philosophical Transactions of the Royal Society of London*, B248: 457-516.
- Dias-da-Silva, S. 1997. Novos achados de vertebrados fósseis na formação Sanga do Cabral. 15 Congresso Brasileiro de Paleontologia, São Pedro, 1997, *Boletim de Resumos*: 82.
- Dias-da-Silva, S. 1998. Novos achados de vertebrados fósseis na Formação Sanga do Cabral (Eotriássico de Bacia do Paraná). *Acta Geologica Leopoldensia*, 21: 101-108.
- Dias-da-Silva, S. & Schultz, C. L. 1999. New procolophonids (Parareptilia, Procolophonoidea) and rhytidosteids (Amphibia, Temnospondyli) of the Triassic of southern Brazil. *Ameghiniana*, 36: 99.
- Dornelles, J. E. F. 1990. Registro sobre a ocorrência de dentes de um arcossáurio para a formação Caturrita, Triássico Superior do Rio Grande do Sul, Brasil. *Ciência e Natura, Santa Maria*, 12: 99-101.
- Faccini, U. F., Scherer, C. M. & Nowatzki, C. H. 1995. Breve panorama sobre a estratigrafia do Permo-Triássico da borda sudeste de Bacia do

- Paraná: Conjeturas e refutações. Comunicações do Museu de Ciências e Tecnologia UBEA/PUCRS Serie Ciências da Terra, 1: 13-18.
- Faccini, U. F., Lavina, E. L., Lopes, R. da C., Schultz, C. L. & Dutra, G. 2000. Gondwana sequences (Early Permian to Early Cretaceous) in southern border of Paraná basin—stratigraphy and paleontology (Rio Grande do Sul, southern Brazil). 31 International Geological Congress, Pre-Congress Field Trip-Bft 02, Guidebook, 37 p.
- Ferigolo, J. 1999. South American first record of a Sphenodontia (Lepidosauria, Rhynchocephalia) from Late Triassic-early Jurassic of Rio Grande do Sul State, Brazil. VII International Symposium on Mesozoic Terrestrial Ecosystems, Buenos Aires, Argentina, Abstracts:25
- Heckert, A. B. & Lucas, S.G. 2000. Taxonomy, phylogeny, biostratigraphy, biochronology, paleobiogeography, and evolution of the Late Triassic Aetosauria (Archosauria: Crurotarsi). Zentralblatt für Geologie und Paläontologie Teil 1 (Heft 11-12): 1539-1587.
- Holz, M. & Barberena, M. C. 1994. Taphonomy of the south Brazilian Triassic paleoherpetofauna: Pattern of death, transport and burial. Palaeogeography, Palaeoclimatology, Palaeoecology, 107: 179-197.
- Huene, F.v. 1935-1942. Die fossilen Reptilien des südamerikanischen Gondwanalandes, ergebnisse der Sauriergrabungen in Südbrasilien 1928/29. C.H. Beck'sche Verlagsbuchhandlung, München.
- Hunt, A. P. & Lucas, S. G. 1991. A new rhynchosaur from West Texas (USA) and the biochronology of Late Triassic rhynchosaur. Palaeontology, 34:191-198.
- Kellner, A. W. A. 1998. Panorama e perspectiva do estudo de répteis fósseis no Brasil. Anais da Academia Brasileira de Ciências, 70: 647-676.
- Kellner, A. W. A. & Campos, D. de A. 1999. Vertebrate paleontology in Brazil—a review. Episodes, 22: 238-251.
- Kellner, A. W. A. & Campos, D. de A. 2000. Brief review of dinosaur studies and perspectives in Brazil. Anais da Academia Brasileira de Ciências, 72: 509-538.
- Kischlat, E. 1999. A new dinosaurian “rescued” from the Brazilian Triassic: *Teyuwasu barberenai*, new taxon. Paleontologia em Destaque, 26: 58.
- Langer, M. C. & Schultz, C. L. 2000. A new species of the Late Triassic rhynchosaur *Hyperodapedon* from the Santa Maria Formation of south Brazil. Palaeontology, 43: 633-652.
- Langer, M. C., Ferigolo, J. & Schultz, C. L. 2000. Heterochrony and tooth evolution in hyperodapedontine rhynchosaur (Reptilia, Diapsida). Lethaia, 33: 119-128.
- Langer, M. C., Abdala, F., Richter, M. & Benton, 1999. A sauropodomorph dinosaur from the Upper Triassic (Carnian) of southern Brazil. Comptes Rendus Academie Science Paris, 329: 511-517.
- Lavina, E. L. 1983. *Procolophon pricei*, sp. n., um novo procolofonídeo do Triássico do Rio Grande do Sul. Iheringia, 2: 51-78.
- Lucas, S. G. 1993. *Barysoma lenzii* (Synapsida: Dicynodontia) from the Middle Triassic of Brazil, a synonym of *Stahleckeria potens*. Journal of Paleontology, 67: 318-321.
- Lucas, S. G. 1998. Global Triassic tetrapod biostratigraphy and biochronology. Palaeogeography, Palaeoclimatology, Palaeoecology, 143:347-384.
- Lucas, S. G. 1999. Tetrapod-based correlation of the nonmarine Triassic. Zentralblatt für Geologie und Paläontologie Teil 1 ( Heft 7-8): 497-521.
- Lucas, S. G. & Harris, S. K. 1996. Taxonomic and biochronological significance of specimens of the Triassic dicynodont *Dinodontosaurus* Romer, 1943 in the Tübingen collection. Paläontologische Zeitschrift, 70: 603-622.
- Lucas, S. G. & Heckert, A. B. 2001. The aetosaur *Stagonolepis* from the Upper Triassic of Brazil and its biochronological significance. Neues Jahrbuch für Geologie und Paläontologie Monatshefte, in press.
- Mattar, L. C. B. 1987. Sobre *Barberenasuchus brasiliensis* gen et sp. nov., do mesotriássico do Rio Grande do Sul, Brasil. I. Diagnoses genérica e específica. 10 Congresso Brasileiro de Paleontologia, Rio de Janeiro, 1987, Anais, 1: 77-98.
- Ochev, V. G. & Shishkin, M. A. 1989. On the principles of global correlation of the continental Tri-

- assic on the tetrapods. *Acta Palaeontologica Polonica*, 34: 149-173.
- Price, L. I. 1947. Um procolofonídeo do Triássico do Rio Grande do Sul. *Boletim da Divisão de Geologia e Mineralogia*, 122: 1-27.
- Rogers, R. R., Swisher, C. C. III, Sereno, P. C., Monetta, A. M., Forster, C. A. & Martínez, R. N. 1993. The Ischigualasto tetrapod assemblage (Late Triassic, Argentina) and  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of dinosaur origins. *Science*, 260:794-797.
- Romer, A. S. & Price, L. I. 1944. *Stahleckeria lenzii*, a giant Triassic Brazilian dicynodont. *Bulletin of the Museum of Comparative Zoology*, 92: 465-491.
- Scherer, C. M., Faccini, U. F., Barberena, M. C., Schultz, C. L. & Lavina, E. 1995. Bioestratigrafia da Formação Santa Maria: Utilização das cenozonas como horizontes de correlação. *Comunicações do Museu de Ciências e Tecnologia UBEA/PUCRS Serie Ciencias da Terra*, 1: 43-50.
- Schultz, C. L. 1995. Subdivisão do Triássico do RS com base em macrofósseis: Problemas e perspectivas. *Comunicações do Museu de Ciências e Tecnologia UBEA/PUCRS Serie Ciencias da Terra*, 1: 25-32.
- Schultz, C. L. & Dias-da-Silva, S. 1999. A possible new pareiasaurid in the Sanga do Cabral Formation, Lower Triassic(?) of southern Brazil. *Paleontologia em Destaque*, 26: 49.
- Schultz, C. L., Scherer, C. M. & Barberena, M. C. 1994. Uma nova proposta de zoneamento bioestratigrafico para o Triássico Superior Sul-Rio-Grandense. 38 Congresso Brasileiro Geologie, *Boletim de Resumos Expandidos*, 1: 107-108.
- Schultz, C. L., Scherer, C. M. & Barberena, M. C. 1998. Biostratigraphy of southern Brazilian Upper Triassic. *Epicontinental Triassic Symposium Halle, Abstracts*: 159-160.
- Schultz, C. L., Scherer, C. M. & Barberena, M. C. 2000. Biostratigraphy of southern Brazilian Middle-Upper Triassic. *Revista Brasileira de Geociências*, 30: 491-494.
- Schwanke, C. 1998. Estabelecimento de rotas de dispersão mediante o estudo da herbivoria em dicinodontes. Unpublished Ph. D. dissertation, Porto Alegre, Universidade Federal do Rio Grande do Sul, 256 p. (volume 1) and figures (volume 2).
- Schwanke, C. & Kellner, A. W. A. 1999. Sobre o primeiro registro de Synapsida no Triássico basal do Brasil. 16 Congresso Brasileiro de Paleontologia, Crato, 1999, *Boletim de Resumos*: 101.
- Schwanke Perruzo, C. 1990. A presença do gênero *Ischigualastia* Cox, 1962 (Reptilia, Synapsida, Therapsida, Anomodontia, Dicynodontia) na formação Santa Maria, Rio Grande do Sul, Brasil. Unpublished M. S. thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, 186 p.
- Schwanke Peruzzo, C. & Araújo-Barberena, D. C. 1995. Sobre a ocorrência do gênero *Ischigualastia* Cox, 1962 na Formação Santa Maria, Triássico do Rio Grande do Sul. *Anais da Academia Brasileira de Ciências*, 67: 175-181.
- Teixeira, A. M. S. 1982. Um novo cinodonte carnívoro (*Probelesodon kitchingi*, sp. nov.) do Triássico do Rio Grande do Sul. *Comunicações do Museu de Ciências da PUCRGS*, 24: 1-31.
- Woodward, A. S. 1907. On some fossil reptilian bones from the state of Rio Grande do Sul. *Revista do Museu Paulista*, 7: 46-57.
- Zacarias, J.D. 1982. Una nova especie de tecodonte aetosaurio (*Aetosauroides subsulcatus*, sp. nov.) de Formação Santa Maria, Triássico de Rio Grande do Sul, Brasil. Unpublished M.S. thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, 69 p.

## Triassic marine reptiles from China

J.-L. Li<sup>1,2</sup>, J. Liu, C. Li

*Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences,*

<sup>2</sup>Corresp. author: li.jinling@pa.ivpp.ac.cn

Marine deposits of the Triassic are extensive in southern and southwestern China. The first Triassic marine reptile, a small pachypleurosaurid, *Keichousaurus hui*, was described from Xingyi, Guizhou Province by C. C. Young in 1958. By the mid-1990s, some sauropterygians, ichthyosaurs and hupehsuchids had been reported from the Lower-Middle Triassic of Anhui, Guizhou, Yunnan and Hubei Provinces, Guangxi Zhuang Autonomous Region, and Tibet Zang Autonomous Region. Although all these marine reptile fossils, except those of *Keichousaurus hui* and *Hupehsuchus nanchangensis*, are incompletely preserved, they are significant in understanding the phylogenetic relationship of relative groups and their paleobiogeography. In 1998 numerous complete skeletons of Triassic marine reptiles, including the first placodont and thalattosaur known in China, were recovered from a new horizon and region, the Wayao Member, Falang Formation of Guanling, Guizhou. As a result, all the major groups of Triassic marine reptiles found in west Tethys, including Eosauropterygia, Placodontia, Ichthyosauria, and Thalattosauria, are present in the Triassic of southern and southwestern China.

1) Remains of marine reptiles from the Chinese Lower Triassic are relatively scattered, and six genera and species have so far been described in Guangxi, Anhui, and Hubei provinces. *Chaohusaurus geishanensis* Young et Dong, 1972 from Qinglong limestone (now the Nanlinghu Formation), Chaoxian County, Anhui Province, is a small ichthyosaur about 0.5 m long with a short and narrow snout, labio-lingually widened posterior teeth, and unusually large forefins. Motani and You (1998) considered the other two species, *Chensaurus chaoxianensis* (Chen, 1985) and *Chensaurus faciles* (Chen, 1985) from the same locality as the synonym of *Chaohusaurus geishanensis*.

*Kwangsisaurus orientalis* Young, 1959 was established on a partial postcranial skeleton collected from the Lower Triassic, Wuming, Guangxi. It was once assigned to Nothosauridae (Handbook of

Chinese Vertebrate fossils, 1979; Sun et al., 1992), but was referred to Pistosauroidea (Sauropterygia) by Rieppel (1999) due to its plate-like coracoid, long and distally expanded transverse process on dorsal vertebrae, three carpal ossifications, and curved humerus without an entepicondylar foramen.

*Nanchangosaurus suni* Wang, 1959 known from Daye Formation in Nanzhang, Hubei, is similar to *Hupehsuchus nanchangensis* (see below) in having a long, flattened rostrum, long neck with short cervical ribs, and small dermal plates over the posterior cervical and trunk vertebrae. These characters give strong evidence of the sister-group affinity for the two genera (Carroll and Dong, 1991). Because the two genera are endemic to eastern Asia, an order and a family (Hupehsuchia Nanchangosauridae) were erected.

*Keichousaurus yuananensis* (Young, 1965a), *Hanosaurus hupehensis* (Young, 1972a) and *Hupehsuchus nanchangensis* (Young, 1972b) are found in the Jialingjiang Formation in Hubei. The age of the Jialingjiang Formation has been much debated. It was once considered as Middle Triassic, but ages of late Early Triassic-early Middle Triassic (Chen and Jin, 1996) or late Early Triassic (Yang and Zhang, 2000) were then suggested. Since all the three taxa, *Keichousaurus yuananensis*, *Hanosaurus hupehensis*, and *Hupehsuchus nanchangensis*, are found in the beds below the uppermost part of the Jialingjiang Formation (attributed to Middle Triassic in the concept of the Jialingjiang Formation, Chen and Jin, 1996), they are most likely to be Olenekian. *Keichousaurus yuananensis* is larger than the type species, *K. hui*, and was established based on an incomplete skeleton. Similar to the latter, its humerus is longer than its femur. The outer margin of the ischium in *K. yuananensis* is crooked.

*Hanosaurus hupehensis*, the holotype of which is incomplete, was originally referred to as a thalattosaur. Rieppel (1998b) assigned it to Pachypleurosauridae based on the following characteristics: nasals meet each other in the midline of the skull; the upper temporal fossa is elongate and curved; the transverse process is developed in caudal vertebra; there is a lack of obvious evidence for the presence of the supratemporal bone. *Keichousaurus* and *Hanosaurus* form two consecutive sister groups to the European pachypleurosaur.

*Hupehsuchus nanchangensis* is different from *Nanchangosaurus* in having three layers of dermal plates above the neural spines in the trunk

region and in the division of the neural spines into proximal and distal elements. The most conspicuous features of the taxon are its spindle-shaped and laterally compressed body and paddle-like limbs that reflect adaptation for high-speed swimming (Carroll and Dong, 1991). Because of the poor record of diapsid reptiles from the Late Permian to Early Triassic, the relationships of Hupehsuchia to other taxa are uncertain.

Based on an incomplete skeleton from Luxi, Yunnan, Young (1978) identified a new species of *Kwangsisaurus*—*K. lusiensis*. Rieppel (1999) thought that *K. lusiensis* might represent a taxon closely related to *Lariosaurus*.

2) Marine reptiles from the Middle Triassic in China, including six genera and species of sauropterygians and ichthyosaurs, are comparatively abundant and diverse. Among them, *Mixosaurus maoteiensis* Young, 1965 is the first recorded ichthyosaur in China, and is represented by an incomplete postcranial skeleton discovered in Renhuai, Guizhou (Young, 1965b). This taxon, with deep amphicoelous vertebrae, expanded rectangular coracoid, and short humerus, resembles the European *Mixosaurus cornalianus* of the same age.

A damaged lower jaw and some disarticulated postcranial bones found in the Middle Triassic (now assigned to the Member I of Guanling Formation) Qingzhen, Guizhou, were recognized as a nothosaur and named as *Chinchenia sungi* (Young, 1965a). This taxon was referred to Pistosauridae on the basis of the amphicoelous vertebrae and the distal expansion of the transverse process. Cladistic analysis shows that *Chinchenia*, *Kwangsisaurus* and *Corosaurus* are trichotomous in the base of Pistosauroidae; these pistosaurs are closely related to members in North America (Rieppel, 1999).

The second species of sauropterygian, *Sanchiaosaurus dengi* (Young, 1965a), is represented by a tooth, several pieces of skull bone, and most of a postcranial skeleton from the Member I, Guanling Formation in the suburb of Guiyang City, Guizhou. As indicated by Rieppel (1999), *S. dengi* is characterized by an ilium without a trace of a preacetabular process, and by a dorsal blade of the ilium that is broad, bulbous, and ornamented with densely set tubercles. Although the material of *Sanchiaosaurus* is not complete, it can be deduced from cladistic analysis that it is the sister-group of all other nothosaurs.

*Keichousaurus hui* Young, 1958 and *Nothosaurus*

sp. were collected from the Zhuganpo Member of Falang Formation in Xingyi, Guizhou. Since the establishment of *K. hui*, countless complete and incomplete skeletons of the taxon have been excavated from Xingyi and adjacent areas. It has a short rostrum, long neck, and broad ulna. Furthermore, its temporal region is relatively elongated. The growth of the humerus is positively allometric during its lifetime, indicating an important role of the forelimbs in locomotion. *K. hui* is deduced to be an ovoviviparous reptile (Lin and Rieppel, 1998).

*Nothosaurus* sp., represented by an incomplete skull, was originally named as *Shingyisaurus unexpectus* (Young, 1965a) and was re-described as a nothosaur based on cranial traits such as pointed conical teeth with striated enamel surfaces, a retracted pineal opening, and contact of the frontal and parietal bones behind the fore-margin of the temporal fossa (Rieppel, 1998a).

3) The fossil localities of the Upper Triassic marine reptiles are distributed in Tibet and Guizhou only. *Himalayasaurus tibetensis* Dong 1972 found in Nyalam, Tunlong, Tibet, is a large ichthyosaur more than 10m long. It has a long rostrum, bioconcave centrum, and a long and narrow scapular bone. Teeth are subpleurodont in the upper jaw and pleurodont in the lower jaw. Another ichthyosaur from Nyalam and Tingji, Tibet, is *Tibetosaurus tingjiensis* (Young, Liu et Zhang, 1982), represented by poorly preserved specimens.

The most important marine reptiles of the Upper Triassic were found in the Wayao Member of the Falang Formation of Guanling, Guizhou. About ten genera, including ichthyosaurs, thalattosaurs and placodonts (the latter two new in China), have been reported. Ichthyosaurs are dominant in the collections. A large number of skeletons of these animals, ranging from several dozen centimeters to 5-6 m in length, were recovered, and probably represent three to four new taxa. Among these, *Qianichthyosaurus zhoui* Li, 1999 has three special features, earlier unreported in other ichthyosaurs. First, the vertebral column has a great arch upwards. Second, the three main bones of the hind limb are slightly stronger than those of the forelimb. Last, in ratio to the cranium, *Qianichthyosaurus* has proportionally the largest orbit in the Order Ichthyosauria. It is similar to shastasaurids from the Triassic in size and shape of limbs, and resembles *Ophthalmosaurus* and *Baptanodon* of the Jurassic in skull shape.

*Sinocyamodus xinpuensis* Li 2000 is the first record of placodonts in China and in regions outside Europe, North Africa and the Middle East. The spe-

cies is characterized by its relatively small size, distinctly elongated orbit, pectoral and pelvic girdles located outside carapace, and limbs and tail covered with osteoderms.

*Anshunsaurus huangguoshuensis* Liu, 1999 was originally referred to Eusauroptrygia, but was transferred to Thalattosauria based on characteristics such as the elongated and tapering premaxillary rostrum, retracted nares, contact of premaxilla and frontal bones, and reduced upper temporal fossa. Thalattosaurs are a monophyletic clade of marine reptiles known from the Middle and Late Triassic of Europe and North America. Phylogenetic analysis indicates that *Anshunsaurus* is the sister taxon of *Askeptosaurus italicus* from the Middle Triassic southern Alpine region, but it differs from the latter in the fusion of postorbital and postfrontal bones, the closed upper temporal fenestra, and the elongated posterior process of the jugal etc. (Rieppel and Liu, 2000). *Xinpusaurus* cf. *X. suni* is the second genus of thalattosaurs found in Guanling, Guizhou. It is distinguished from *Askeptosaurus* and *Anshunsaurus* by its relatively small size and its short, slightly ventrally bent snout. A preliminary phylogenetic analysis indicates potential trans-Pacific relationships: *Xinpusaurus* is more closely related to the North American genus *Nectosaurus* (Liu and Rieppel, 2001).

Yin et al. (2000) described nine genera and ten species (among them nine new genera and species) of sauropterygians and ichthyosaurs based on materials discovered in the Wayao Member, Falang Formation. These are part of a private collection, and are not freely accessible to other paleontologists; therefore, until the material can be reviewed and restudied in detail, we do not include these genera and species in the list of marine reptiles.

It is worth mentioning that there have been varying opinions on some genera and fauna. For example, there is a strong argument about the Zhuganpo Member, Falang Formation yielding *Keichousaurus* fauna. It was considered as Middle Triassic (Young, 1958; Chen, 1985), Middle-Late Triassic (Dong et al, 1997), or Late Triassic (Wang, 1996; Wang, 1998). Comparison of Chinese marine reptiles with their counterparts in Europe lead us to agree with Young (1958) and Chen (1985), so we assign the Zuganpo Member to the Middle Triassic. That is, we assign a Ladinic age.

### Acknowledgements

We gratefully acknowledge Dr. Olivier Rieppel for his helpful discussion on Triassic marine reptiles of China. Our cordial thanks are due to Dr. Michael

J Orchard (Geological Survey of Canada) and Dr. Ruth Hubert (Louisiana State University) for assistance with the manuscript.

This research was supported by National Natural Science Foundation of China (40072010) and Innovation Found of Chinese Academy of Sciences (KZCX3-J-02).

### References

- Carroll R, Dong Z-M, 1991: Hupehsuchus, an enigmatic aquatic reptile from the Triassic of China, and the problem of establishing relationship. *Phil. Trans. R. Soc. Lond. B* 331:131-153.
- Chen G-X, Jin J-W, 1996: Stratigraphy (lithostratic) of Hubei Province. Multiple classification and correlation of the stratigraphy of China. 42, China University of Geosciences Press. 1-284.
- Chen Z-F, 1985: Stratigraphical position of *Keichousaurus hui* Young of Middle Triassic and its significance in southwestern Guizhou. *Geol. Guizhou*, 5: 289-290.
- Dong W-P et al, 1997: Stratigraphy (lithostratic) of Guizhou Province. Multiple classification and correlation of the stratigraphy of China. 52, China University of Geosciences Press. 1-306.
- Dong Z-M, 1972: An ichthyosaur from the region of Qomolangma Feng, Tibet, China. *Mem. IVPP. Acad. Sin.*, A 9: 7-10.
- Li C, 1999: Ichthyosaur from Guizhou China. *Chinese Science Bulletin* 44(14):1429-1333.
- Li C, 2000: Placodont (Reptilia, Placodontia) from Upper Triassic of Guizhou, southwest China. *Vert. PalAsiat.* 38(4): 314-317.
- Lin K-B, Rieppel O, 1998: Functional morphology and ontogeny of *Keichousaurus hui* (Reptilia, Sauropterygis). *Fieldiana Geol.* n s. 39: 1-35.
- Liu J, 1999: Sauropterygian from Triassic of Guizhou, China. *Chinese Science Bulletin.* 44(4):1312-1315.
- Liu J, Rieppel O, 2001: The second thalattosaur from the Triassic of Guizhou, China. *Vert. PalAsiat.* 39(2): 77-87.
- Motani R, You H-L, 1998: Taxonomy and limb ontogeny of *Chaohusaurus geishanensis* (Ichthyosauria), with a note on the allometric equation. *J. Vert. Paleont.* 18(3):533-540.

- Rieppel O, 1998a: The status of *Shingyisaurus unexpectus* from the Middle Triassic of Kweichou, China. *J. Vert. Paleont.* 18(3): 541-544.
- Rieppel O, 1998b: The systematic status of *Hanosaurus hupehensis* (Reptilia, Sauropterygia) from the Triassic of China. *J. Vert. Paleont.* 18(3):545-557.
- Rieppel O, 1999: The sauropterygian genera *Chinchenia*, *Kwangsisaurus*, and *Sanchaosaurus* from the Lower and Middle Triassic of China. *J. Vert. Paleont.* 19(2): 321-337.
- Rieppel O, Liu Jun, 2000: The first record of a thalattosaur reptile from the Late Triassic of Southern China (Guizhou Province, PR China). *J. Vert. Paleont.* 20(3): 507-514.
- Sun A-L, Li J-L, Ye X-K, et al, 1992: The Chinese Fossil Reptiles and Their Kins. Science Press, Beijing and New York, 1-260.
- Wang C-Y, Kang P-Q, Wang Z-H, 1998: Conodont-based age of the *Keichousaurus hui* Young, 1958. *Acta Micropalaeont. Sinica*, 15(2): 196-198.
- Wang K-M, 1959: Discovery of a new reptile fossil from Hubei, China. *Acta Palaeont. Sinica*. 7(5):367-378.
- Wang L-T, 1996: A discussion on horizon and age of *Keichousaurus hui* occurrence. *Guizhou Geology*, 13(3): 209-212.
- Yang Z-Y, Zhang S-X, 2000: Stratigraphical Lexicon of China, Triassic. Geology Publishing House. Beijing, 1-139.
- Yin Gongzeng, Zhou Xiugao, Cao Zetian, Yu Youyi, Luo Yongming, 2000: A preliminary study on the Early Late Triassic marine reptiles from Guanling, Guizhou, China. *Geology-Geochemistry*. 28(3): 1-23.
- Young C C, 1958: On the new Pachypleurosauroidea from Keichow, southwest China. *Vert. PalAsiat.* 2(2-3): 69-81.
- Young C C, 1965a: On the new nothosaurs from Hupeh and Keichow, China. *Vert. PalAsiat.* 9(4):315-356.
- Young C C, 1965b: On a revised determination of a fossil reptile from Jenhui Kweichow with note on a new ichthyosaur probably from China. *Vert. Pal. Asiat.* 9(4): 368-375.
- Young C C, 1972a: A thalattosaur from Nanchang, Hubei, China. *Mem. IVPP. Acad. Sin.*, A9: 17-27.
- Young C C, 1972b: *Hupehsuchus nanchangensis*. *Mem. IVPP. Acad. Sin.*, A9:28-34.
- Young C C, 1978: *Kwangsisaurus lusiensis* from Yunnan. *Vert. PalAsiat.* 16(4): 222-224.
- Young C C, Dong Z-M, 1972: An ichthyosaur (*Chaohusaurus geishanensis*) from Anhui, China. *Mem. IVPP. Acad. Sin.*, A 9: 11-13.
- Young C C, Liu D-S, Zhang M-L, 1978: *Ichthyosaur from Tibet, China. Monograph on Mount Xixia Bangma, Scientific Expedition, 1964. Science Press.* 350-355.



Triassic Marine Reptiles

	Ichthyosaurs	sauropterygian		thalatosaurs	hupehsuchian
		eosauropterygian	placodonts		
3	<i>Himalayasaurus tibetensis</i> <i>Tibetosaurus tingiensis</i> <i>Qianichthyosaurus zhoui</i>		<i>Sinozyamodus xinpuensis</i>	<i>Anshunsaurus huangguosihuensis</i> <i>Xinpusaurus cf. X.suni</i>	
2	<i>Mixosaurus maotiensis</i>	<i>Keichousaurus hui</i> <i>Nothosaurus</i> sp. <i>Chinchenia sungi</i> <i>Sanchiaosaurus dengi</i> <i>Kwangsisaurus lusiensis</i>			
1	<i>Chaohusaurus geishanensis</i>	<i>Kwangsisaurus orientalis</i> <i>Keichousaurus yunanensis</i> <i>Hanosauruw hupehensis</i>			<i>Hupehsuchus nanchangensis</i> <i>Nanchangosaurus suni</i>

## A proposed area for the study of the Accessory Section and Point of the Terrestrial Permian-Triassic Boundary

PENG Yuanqiao<sup>1,3</sup>, YIN Hongfu<sup>1</sup>, YANG Fengqing<sup>1</sup> and WANG Shangyan<sup>2</sup>

<sup>1</sup> Faculty of Earth Science, China University of Geosciences, Wuhan 430074, Hubei, China

<sup>2</sup> Guizhou Geological Survey, Guiyang 550004, Guizhou, China

<sup>3</sup>Corresp. author: [digushi@cug.edu.cn](mailto:digushi@cug.edu.cn)

After the establishment of the Global Stratotype Section and Point (GSSP) of the Permian-Triassic boundary (PTB), definition of the Accessory Section and Point (ASP) of the Terrestrial Permian-Triassic Boundary (TPTB) is now on the agenda. However, all good TPTB sections so far known have the following shortcomings: (1) the exact TPTB horizon is difficult to define with high-resolution fossils; (2) correlation between marine and terrestrial PTBs is hard to attain with accuracy. These, of course, enhance the understanding of the global life crisis from marine to land across the Paleozoic-Mesozoic transition. In western Guizhou and eastern Yunnan of southwestern China, some fossiliferous PTB sections from marine via paralic to land facies are well developed, allowing bed-to-bed correlation of the PTB sequences among themselves. Fortunately, the marine PTB sequence in this area is almost the same as the Meishan section, where the GSSP of the PTB locates, which may provide a reliable auxiliary marker for high-resolution demarcation of the TPTB. These merits in western Guizhou and eastern Yunnan made it a good place to study the ASP of the TPTB, so we propose to study the ASP of the TPTB in this area.

Keywords: Terrestrial Permian-Triassic boundary (TPTB), Accessory Section and Point (ASP), Western Guizhou and eastern Yunnan

The Global Stratotype Section and Point (GSSP) of the Permian-Triassic boundary (PTB) was ratified by the IUGS Executive Committee in March, 2001. Thus, the GSSP of the PTB is defined at the base of Bed 27c, the Meishan Section D, Changxing County, Zhejiang Province, China, at the horizon where the conodont *Hindeodus parvus* first appeared (Yin et al., 2001).

With the establishment of the GSSP of the PTB, the Accessory Section and Point (ASP) of the Terrestrial Permian-Triassic Boundary (TPTB) should be defined as well. Cowie et al. (1986) pointed out that the ASP is especially useful in two clearly different kinds of strata such as the correlation of the New Red Sandstone and the marine Triassic. The marine bio-events across the Paleozoic-Mesozoic transition have long been researched along with the study of the GSSP of the PTB, but the Paleozoic-Mesozoic life crisis on land is hard to estimate due to the uncertainty of the TPTB. Therefore, high-resolution definition and correlation of the TPTB or the terrestrial Paleozoic-Mesozoic boundary is now on the agenda.

### Status quo of the TPTB research

The terrestrial deposits of the Permian-Triassic are widely developed all over the world. The global distribution of the TPTB strata is no less than that of the marine PTB. However, the study of the TPTB is not so successful as the marine PTB, and the definition and correlation of the TPTB is more difficult than the marine PTB. Some investigators have put forward proposals for the construction of the special time scale (parallel to the existing marine one) for continental deposits (Lozovsky, 1991; Lucas et al., 1992). Accordingly, the search and the establishment of the ASP of the TPTB should be proposed as well. However, researches on the subdivision and correlation of the TPTB sequence and the fauna feature on land are relatively weak. Until now, few type sections of the TPTB have been proposed.

At present, some scholars are inclined to search the ASP of the TPTB in South Africa (Lucas et al., 1996). The Permian-Triassic Beaufort Group of Karoo basin in South Africa belongs to the sediments of a large intracratonic retro-arc foreland basin in southwestern Gondwana (Smith, 1995). Many vertebrate fossils were found and well studied in this area (Kitching, 1977). The *Cistecephalus* Zone and *Lystrosaurus* Zone in this area correspond to the *Dicynodon* fauna and *Lystrosaurus* fauna respectively in Xinjiang, northwestern China (Zhao, 1980; Cheng, 1986). It has become one of the most important areas for the study of the TPTB all over the world, but the study of other kinds of fossils was not as thorough as that of the vertebrates (D'Engelbronner, 1996). Many difficulties are met when attempts are made to establish correlation between the various regional scales, or to locate the continental tetrapod biozones on the universal stratigraphic scale (Battail, 1995; Shishkin, 1994). Some of the problems which are encountered in global correlation are illustrated with the

well known example of the biozones of the Beaufort Group of Karoo basin in South Africa by Battail (1995).

The Chinese mainland lay in the east part of Tethys during Pangea time and represents one of the most developed areas of terrestrial Permian and Triassic in the world. The terrestrial Permian and Triassic in China are mainly distributed to the north of Kunlun-Qinling Mts. Among them, the Dalongkou section, Jimsar, Xinjiang is the most exhaustively studied TPTB in China, and its biostratigraphy is well studied. The TPTB defined by integrative subdivision of vertebrates, bivalves, ostracods, plants and palynomorphs can be well correlated among different continents (Yang et al., 1988; Zhou et al., 1997). The eventostratigraphic studies, such as magnetostratigraphy (Li et al., 1997), sedimentary events and climate change events (Yang et al., 1988, 1992), have been carried out as well. Clearly, it represents one of the world's best records of the terrestrial Permian-Triassic transition, and further study of this section could provide more precise correlation to the standard global chronostratigraphic scale and a more detailed understanding of terrestrial biotic changes across the PTB (Lucas et al., 1996). Cheng (1993) and Cheng and Lucas (1993) proposed that the Dalongkou section be considered as a potential auxiliary (non-marine) GSSP for the PTB.

Some other areas, such as the Moscow syncline in Russia and the Cis-Ural region in Kazakhstan, the Bowen basin and Sydney basin in eastern Australia, and the Zechstein basin in Germany and its surrounding countries, developed typical TPTB sections and have been well studied as well.

At present, however, all the preferable TPTB sections in the world have the following shortcomings. (1) The exact TPTB position is hard to define with high-resolution due to sporadic occurrence of vertebrates in few beds. As a result, between the assured Permian and the assured Triassic, there are always intervals of several tens of meters of uncertain age. Sometimes, mixed fauna, which cannot be correlated with marine GSSP, can be found in it. Among them, different kinds of fossils have different time ranges, so it is hard to demarcate the assured TPTB at those sections. Therefore, the life crisis on land across the TPTB and its sequence are hard to estimate as well. (2) The correlation between marine and terrestrial PTBs is hard to be attained with accuracy. These, of course, hamper the understanding of the global life crisis across the Paleozoic-Mesozoic transition.

If we try to establish the ASP of the TPTB in a wholly

continental area, correlation between marine and continental PTBs will be difficult. However, if an area contains both marine and terrestrial PTBs via paralic facies, there is an opportunity to correlate all those different facies directly, and the problems mentioned above might be solved. At last, we found an area (western Guizhou and eastern Yunnan, China) coincident to this condition. In western Guizhou and eastern Yunnan, apparently continuous depositions of PTB strata from marine via paralic facies to land are very well recorded. These strata are fossiliferous. The most advantageous phenomenon is that the marine PTB sequence in this area is almost the same as the Meishan section, the locality of the GSSP of the PTB, allowing bed-to-bed correlation of the type PTB strata with the marine PTB strata of this area, and thus also with the TPTB strata there. The merits of sections in western Guizhou and eastern Yunnan made this area a good place to study the ASP of the TPTB.

### General geology in western Guizhou and eastern Yunnan

The terrestrial strata across the PTB in western Guizhou and eastern Yunnan are composed, in ascending order, of Late Permian Xuanwei Formation, Late Permian and Early Triassic Kayitou Formation, and Early Triassic Dongchuan Formation. Based on detailed study of biostratigraphy, Nanjing Institute of Geology and Palaeontology, Academia Sinica (1980) gave a systematic conclusion of the strata in western Guizhou and eastern Yunnan. They attributed the terrestrial coal-bearing strata, with occasional marine interlayers, into Xuanwei Formation. It is overlain by Kayitou Formation (or Kayitou beds) with conformity. The lithologies of Kayitou Formation are almost the same as that of Xuanwei Formation except no coal beds are interbedded. According to fossils such as bivalves, ostracods and palynomorphs in the Kayitou Formation, it was attributed either to the earliest stage of Early Triassic by the research group of Nanjing Institute of Geology and Palaeontology, Academia Sinica (1980), or to late Late Permian and early Early Triassic (Wang and Yin, 2001). The Dongchuan Formation also overlies Kayitou Formation with conformity. There are clear differences of lithologic colors between Xuanwei Formation, Kayitou Formation and Dongchuan Formation. At the outcrop sections, the gray to gray-green strata belong to the Xuanwei Formation, the variegated strata belong to the Kayitou Formation, while the amaranth(?) strata belong to Dongchuan Formation. The nonmarine PTB strata in western Guizhou and eastern Yunnan all belong to these types and can be well corre-




Figure 1. Sketch map of the lithofacies and paleogeography from Late Permian (Changhsingian) to Early Triassic in western Guizhou and eastern Yunnan (after Yao et al., 1980).

lated regionally.

There are many excellent PTB sections from ma-

rine via paralic to land in western Guizhou and eastern Yunnan. This area is one of the ideal regions to study the subdivision and correlation of the PTB strata from marine to land. Based on strati-

Common sequence across marine PTB in Yangtze platform			Sequence across the PTB at the Meishan section			Sequence across the TPTB at the Chahe section		
Overlying strata	marl (or calcareous mudstone)		29		marl	57	siltstone and mudstone	
PTBST	Top Clay	clay (or/and mudstone) (0-10 cm)	28		clay (3-5 cm)	56b	clay (10 cm)	
	Boundary Rock	micritic limestone (or marl) (10-30 cm)	T	27d	micritic limestone (15-17 cm)	56a	clay (25 cm)	
				27c				
			P	27b		55	mudy siltstone (25 cm)	
				27a				
		26		'Black Clay' (shale) (5-12 cm)				
Bottom Clay	clay (or/and mudstone) (1-20 cm)	25		'White Clay' (0.5-5 cm)	54	clay (20 cm)		
Underlying strata	siliceous micritic limestone (or siliceous mudstone, silicalite)		24e		siliceous micritic limestone	53	silty sandstone and fine sandstone	

Figure 2. Common sequence across the PTB from marine to land in Yangtze region and its correlation.

graphic study of many sections in western Guizhou and eastern Yunnan from the Late Permian to the Early Triassic, the research group of Nanjing Institute of Geology and Palaeontology, Academia Sinica (1980) drew the sketch evolutionary map of lithofacies and paleogeography of western Guizhou and eastern Yunnan during the Permian-Triassic transition (Fig.1). From this figure it is evident that not only marine PTB (clastic facies and carbonate facies) but also non-marine PTB (paralic facies and terrestrial facies) can be found in this area. The shoreline of Changhsingian rarely regressed to the east of Nayong-Liuzhi line, and the shoreline lay approximately to the east of Shuicheng-Qinglong line at the end of the Permian. At the beginning of the Triassic, the seawater transgressed a little to the west, and the shoreline moved accordingly westward to Xuanwei-Fuyuan line. Thus the genuine TPTB sections should be searched to the west of Xuanwei-Fuyuan line. All three reported TPTB sections, Laibin section, Xuanwei, Yunnan (Nanjing Institute of Geology and Palaeontology, Academia Sinica, 1980), Zhejue section and Chahe section, Weining, Guizhou (Wang and Yin, 2001), lie in the western area.

### Similar bed sequence across the PTB from marine to terrestrial facies in western Guizhou and eastern Yunnan

Latest study in western Guizhou and eastern Yunnan reveals a set of TPTB sequences which possess almost the same lithology and event associations as the marine Permian-Triassic Boundary Stratigraphic Set (PTBST) in South China. There are also one to two clayrock beds across

the TPTB in western Guizhou and eastern Yunnan. The results of X-ray diffraction indicate that the main mineral compositions of those clayrocks are illite-montmorillonite and kaolinite, which are almost the same as the compositions of the marine PTB clayrocks. Zircon, apatite, hexagonal dipyrmaid quartz and some other accessory minerals of acidic lava have been found in the TPTB clayrocks at Chahe section, Weining of Guizhou Province and Mide section, Xuanwei of Yunnan Province. Shape parameters (length and width) of the zircons and their total frequencies are recorded. These data show that they are the same as the dirt bands in the coal beds of the Xuanwei Formation, which were considered to be formed by weathering of volcanic sediments (Zhou and Ren, 1983). In addition, transparent hyaline micro-spherules are occasionally found in the boundary clayrocks of the two sections. A lot of transparent hyaline micro-spherules and black metal micro-spherules are also found in the TPTB clayrocks at Zhejue section, Weining of Guizhou Province. Transparent hyaline micro-spherules are usually round, clear and transparent. Black metal micro-spherules are also round, some appear teardrop shaped and some have irregular cracks on their surfaces. Both types of the micro-spherules appear with complete extinction under the cross polarizer. Those characters of the micro-spherules indicate that they are formed through quick condensation of melted materials. Consequently, It is proposed that the TPTB clayrocks are the result of volcanism, which coincides to the formation of the marine PTB clayrocks of South China (Yin et al., 1989).

A typical three-layer structure consisting, in ascend-

ing order, of claystone (bed 54), muddy siltstone (bed 55) and claystone (bed 56) appears across the TPTB at the Zhejue section. This kind of TPTB association represents the same as that of the Meishan section, type section of the PTBST (Fig.2) (Peng et al., 2001). The following characteristics are found in the underlying and overlying strata of the terrestrial PTBST. (1) Biostratigraphic aspect: the palynomorph assemblages show clear differences between the underlying and overlying strata of bed 54. Typical members of the Triassic are only abundant in bed 54 and its overlying strata. Plant fossils are abundant in the underlying strata of bed 54 and are few in the overlying strata (Wang and Yin, 2001). (2) The susceptibility appears distinctly different between the underlying and the overlying strata of bed 54 and becomes increasingly higher in ascending order (Wang and Yin, 2001). This kind of changing trend is almost the same as that across the marine PTB in South China and can be correlated with high-resolution (Peng et al., 2000). (3) The data of  $\delta^{13}\text{C}_{\text{org}}$  change abruptly across bed 54 and then recover.  $\delta^{13}\text{C}_{\text{org}}$  data are relatively lower and stable in the underlying strata of bed 54 (-24.51~ -26.98‰) and higher and changeable in the overlying strata (-17.63~ -27.20‰) (Wang and Yin, 2001). These characteristics indicate that the terrestrial PTBST might also represent the high-resolution chronostratigraphic PTB.

The continuous clayrocks from marine via paralic facies to land in western Guizhou and eastern Yunnan provide a reliable auxiliary marker for high-resolution demarcation of the TPTB. Thus, in this area exists potential for high-resolution subdivision and correlation of the PTB from marine to land. This can then help us understand the terrestrial and global life crisis across the Paleozoic-Mesozoic transition.

### Conclusions

(1) There are many excellent PTB sections from marine via paralic facies to land in western Guizhou and eastern Yunnan. This area is one of the ideal regions to study the subdivision and correlation of the PTB from marine to land. Thus, we propose to study the ASP of the TPTB in this area.

(2) Marked by the PTB clayrock beds, with the study of the biostratigraphy and some other eventostratigraphic characters such as susceptibility, carbon isotope and radiometric dating across the PTBST, the high-resolution correlation framework of the PTB from marine via paralic to land can be set up in western Guizhou and eastern Yunnan. This will help to establish the high-resolution correlation framework of the PTB in South

China and over the world.

(3) Establishment of high-resolution chronostratigraphic TPTB in western Guizhou and eastern Yunnan will help us to understand the global life crisis from marine to land across the Paleozoic-Mesozoic transition.

However, the work in this area is preliminary and further study is necessary. We are appreciative of all cooperation with research in this area.

### Acknowledgements

This work is a part of the research programs supported by the National Science Foundation of China and the China Geological Survey.

### References

- Battail B., 1995, Continental Permo-Triassic biostratigraphic scales based on tetrapod vertebrates: a few problems. *Bull. Soc. Geol. France*, 166: 527-535.
- Cheng Zhengwu, 1986, Vertebrates. In: *Institute of Geology, Chinese Academy of Geological Sciences and Institute of Geology, Xinjiang Bureau of Geology and Mineral Resources (eds.), Permian and Triassic strata and fossil assemblages in the Dalongkou area of Jimsar, Xinjiang*. Beijing: Geological Publishing House, 207-218.
- Cheng Z., 1993, On the discovery and significance of the nonmarine Permo-Triassic transition zone at Dalongkou in Jimsar, Xinjiang, China. In: Lucas S.G. and Morales M. (eds.), *The nonmarine Triassic*. New Mexico Museum of Natural History and Science Bulletin, 3: 65-67.
- Cheng Z. and Lucas S.G., 1993, A possible non-marine GSSP for the Permian-Triassic boundary. *Albertiana*, 12: 39-44.
- Cowie, J.W., Ziegler, W., Boucot, A.J., Bassett, M.G. and Remane, J., 1986, Guidelines and statutes of the International Commission on Stratigraphy. *Cour. Forsch.—Inst. Senckenberg*, 83: 1-14.
- D'Engelbronner E.R., 1996, New palynological data from Karoo sediments, Mana Pools basin, Northern Zimbabwe. *J. Afr. Earth Sci.*, 23: 17-30.
- Kitching J.W., 1977, The distribution of the Karoo vertebrate fauna. *Bernard Price., Inst. Palaeont. Res. Mem.*, 1: 1-131.

- Li Yongan, Wu Shaozu and Sun Dongjiang, 1997, The magnetostratigraphy across the Permian-Triassic boundary in Tianshan area, China. *Xinjiang Geology*, 15(3): 227-235 (in Chinese with English abstract).
- Lozovsky V.R., 1991, Proposal for a new working group on continental beds at the Permian/Triassic boundary in the continental series. *Permophyles*, 19: 7-11.
- Lucas S.G., Lozovsky V.R., Cheng Zhengwu, 1992, Proposal to STS for formation of a working group on the nonmarine Triassic timescale. *Albertiana*, 10: 41.
- Lucas S.G., H. Kozur, R. Molina-Garza and J. Geissman, 1996, Reconsideration of Dalongkou as an auxiliary GSSP for the Permian-Triassic boundary. *Albertiana*, 18: 10-11.
- Nanjing Institute of Geology and Palaeontology, Academia Sinica, 1980, Late Permian coal-bearing strata and palaeontological fauna in western Guizhou and eastern Yunnan. Beijing: Science Press, 277pp (in Chinese).
- Peng Yuanqiao, Tong Jinnan, Hans J. Hansen and Guang R. Shi, 2000, Susceptibility and its significance across the Permian-Triassic boundary in South China. *Geology and Mineral Resources of South China*, 2: 22-28 (in Chinese with English abstract).
- Peng Yuanqiao, Tong Jinnan, G.R. Shi and Hans J. Hansen, 2001, The Permian-Triassic Boundary Stratigraphic Set: characteristics and correlation. *Newsletters on Stratigraphy*, 39(1): 55-71.
- Shishkin M.A., 1994, Problems of global correlation of the continental Triassic on the basis of tetrapods. In: Guex J. and Baud A. (eds.), *Recent developments on Triassic stratigraphy. Mém. de Géol. (Lausanne)*, 22: 121-126.
- Smith R.M.H., 1995, Changing fluvial environments across the Permian-Triassic boundary in the Karoo Basin, South Africa and possible causes of tetrapod extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 117: 81-104.
- Wang Shangyan and Yin Hongfu, 2001, Study on terrestrial Permian-Triassic boundary in eastern Yunnan and western Guizhou. Wuhan: China University of Geosciences Press, 88pp (in Chinese with English abstract).
- Yang Jiduan, Qu Lifan, Zhou Huiqin, Cheng Zhengwu, Zhou Tongshun, Hou Jingpeng, Li Peixian and Sun Suying, 1988, Continental Permian-Triassic boundary and event. *Geoscience*, 2(3): 366-374 (in Chinese with English abstract).
- Yang Jiduan, Qu Lifan, Zhou Huiqin, Cheng Zhengwu, Zhou Tongshun, Hou Jingpeng, Li Peixian, Sun Shuying, Wu Shaozu, Li Daiyun and Long Jiarong, 1992, Classification and correlation of nonmarine Permo-Triassic boundary in China. In: Sweet W.C., Yang Zunyi, J.M. Dickins & Yin Hongfu (eds.), *Permo-Triassic events in the eastern Tethys*. London: Cambridge University Press, 56-59.
- Yao Zhaoqi, Xu Juntao, Zheng Zhuoguan, Zhao Xiugu and Mo Zhuanguan, 1980, Late Permian biostratigraphy in western Guizhou and eastern Yunnan and the Permian-Triassic boundary. In: Nanjing Institute of Geology and Palaeontology, Academia Sinica (ed.), *Late Permian coal-bearing strata and palaeontological fauna in western Guizhou and eastern Yunnan*. Beijing: Science Press, 1-69 (in Chinese).
- Yin Hongfu, Huang Siji, Zhang Kexin, Yang Fengqing, Ding Meihua, Bi Xianmei and Zhang Shuxing, 1989, Volcanism at the Permian-Triassic boundary in South China and its effects on mass extinction. *Acta Geologica Sinica*, 2: 169-181 (in Chinese with English abstract).
- Yin Hongfu, Zhang Kexin, Tong Jinnan, Yang Zunyi and Wu Shunbao, 2001, The Global Stratotype Section and Point (GSSP) of the Permian-Triassic Boundary. *Episodes*, 24(2): 102-114.
- Zhao Xijin, 1980, Mesozoic vertebrate biostratigraphy in North Xinjiang. *Monograph of Institute of Vertebrate Paleontology and Paleoanthropology, Academia Sinica, Series A, Number 15*, Beijing: Science Press (in Chinese).
- Zhou Yiping and Ren Youliang, 1983, Formation of some dirt bands (Tonteins) in the coal beds of the Upper Permian Xuanwei Formation and their significance. *Yunnan Geology*, 2(1): 38-44 (in Chinese with English abstract).
- Zhou Tongshun, Li Peixian, Yang Jiduan, Hou Jiping, Liu Shuwen, Cheng Zhengwu, Wu Shaozu and Li Yongan, 1997, Stratotype section of non-marine Permian-Triassic boundary in China. *Xinjiang Geology*, 15(3): 211-226 (in Chinese with English abstract).

## Meeting Reports

### Oman Pangea Symposium and Fieldmeeting, Muscat, Sultanate of Oman, 7-20 January, 2001

Aymon Baud<sup>1</sup>, Jean Marcoux<sup>2</sup>,  
Benoit Beauchamp<sup>3</sup>

<sup>1</sup> BFSH2-UNIL, CH-1015 Lausanne, Switzerland, [aymon.baud@sst.unil.ch](mailto:aymon.baud@sst.unil.ch), <sup>2</sup> Paris VII University, Sciences physiques de la terre, Pl. Jussieu 2, F-75251 Paris Cedex 05 France, <sup>3</sup> Geological Survey of Canada, 3303, 33 street NW, Calgary, Alberta T2L 2A7, Canada

The preparation of the International Conférence on the Geology of Oman organised by the Oman Ministry of Commerce and Industry in cooperation with Sultan Qaboos University and the Mineralogical Institute of Bern University was a unique opportunity for a Pangea Symposium proposal in this part of the world. This was done in 1999 by A. Baud and J. Marcoux and accepted the same year by the Organising Committee.

Sponsored by the Subcommittee on Triassic Stratigraphy among other institutions, the Symposium and fieldmeeting was to provide a forum to geologists who are interested in the Pangean time interval for discussing global changes related to Pangea integration and North Gondwana and Central Tethys evolution. Members of the International Subcommittees on Permian and on Triassic Stratigraphy were invited. Sedimentologists, stratigraphers and paleontologists working within the Permian and Triassic time interval and interested by topics as diverse as biotic crisis, extinction, recovery and evolution across the Palaeozoic-Mesozoic transition used the opportunity to discuss, observe and sample the spectacular Permian and Triassic outcrops of Oman's former continental margin, from shallow shelf to deep marine sediments and sea mounds and to participate in the Pangea Symposium.

A comprehensive report has been published in *Episodes*, vol. 21/2, p. 126-127 and here we are giving the main results concerning the Triassic stratigraphy.

The thematic of the **Pre-Conference Excursion**

No. A01, from January 8 - 11, 2001, was «Permo-Triassic Deposits, from shallow platform to Basin and Seamounts». It was led by A. Baud, F. Bechennec, L. Krystyn, J. Marcoux R. Maury and S. Richoz. Sixteen participants attended this fieldtrip with great interest and took part in the lively discussions on the outcrops.

Very interesting new data concerned the late Triassic and Triassic-Jurassic transition and were presented by Leopold Krystyn. The first occurs in the Baid exotic block which is a worldwide unique exposure because it witnesses an Early Mesozoic stable pelagic environment. It persisted with just minor changes for more than 100 million years. It represents a tiny piece of a paleogeographic realm where more stable conditions should have provided better survival chances across the T-J crisis interval than the environmentally stressed shallow shelf regions of the oceans. This realm must have been widespread in the tropical belt of the Neotethys ocean but has been lost nearly completely by later subduction and collision. An outcrop of about 20m thick, predominantly nodular thin bedded red ammonoid bearing limestones comparable to the ammonitico rosso facies of the Southern Alps and the Adnet limestone of the Austrian Calcareous Alps have their base in the Norian and their top in the Sinemurian!

The second Triassic outcrop concerns the Al Aqil Block with the Aqil breccia. The reconstructed stratigraphic sequence consists of basal pillow basalts followed mainly by pelagic sediments on which the Aqil breccia is resting as debris flow with erosional contact on the lower fan sediments underlain by dark brown radiolarites. Among the described blocks, there is a basal forereef breccia (Lower Carnian) a crinoidal-brachiopod packstone (Upper Carnian) and finally a cephalopod Hallstatt-type limestone unit (Norian-Sinemurian).

The third Triassic outcrop concerns the Upper Triassic reefal platform margin of the Kawr Group. A careful survey of the base of the Misfah formation has led to the vertical discrimination of 1) a thin basal interval (about 5m, dated of Tuvalian 3) overlain by 2) allodapic lower slope deposits (50m) changing into 3) reefal limestones (appr. 100m) which are finally topped by 4) cyclic bedded lagoonal megalodont-bearing limestones of several 100m in thickness (Norian).

Between Jan. 12-16, 2001, the Conference on the Geology of Oman has attracted about 400 scientists from all over the world. As part of the Conference, the 2-day **Pangea Symposium** started on Jan. 14 in which 18 oral communications and 5



posters where presented. Between 50 and 100 scientists followed the Symposium. In their introduction, A. Baud and B. Beauchamp stated the general purposes of the Pangea project and emphasised the international cooperation. Concerning the Triassic sediments, the main items of scientific progresses are summarised as follow: Bernecker, M. (Second-order cycle development of the Arabian platform and Hawasina seamounts: Permian and Triassic outcrop data from central Oman), Cordey, F., Baud, A., Béchenec, F., Gorican, S. Krystyn, L. and Robin C. (Permian-Triassic deep water sediments of the Wadi Wasit revisited).

Recent isotope studies were presented by Richoz, S., Atudorei, V., Baud, A. and Marcoux, J. (Upper Permian to lower Triassic carbon isotope record: review and new data in the Oman Mountains, from the shallow platform to the basin) and Richoz, S., Baud, A., Marcoux, J. and Cordey, F. (Lower Triassic carbon isotope stratigraphy of the Sumeini slope deposits (Maqam C, NW Oman).

New data on Permian-Triassic boundary were the subject of a talk by Baud, A., Cordey, F., Krystyn, L. Marcoux, J. and Richoz, S. (The Permian-Triassic boundary in Oman, a review) and of a poster by Krystyn, L., Richoz, S. and Baud, A. (A Unique Permian-Triassic Boundary section from Oman): this is the first discovery of a complete dated Griesbachian coquinite limestone succession in Oman. This facies is unknown in other part of the Tethys.

About Triassic paleoclimates, stratigraphy, magnetic insight and palynofacies, the first presentation has been given by MacDonald, W. and Ellwood, B. on «Magnetic Insights into Permo-Triassic Pangea».

Stratigraphy of Triassic sediments was presented by Bachmann, G.H., Brueckner-Roehling, S., Exner, M., Kedzierski, J. and Szurlies, M. (Sequence Stratigraphy of the Scythian-Anisian Transgression, Triassic Type Region, Germany), by Mandl, G. W. (From Triassic Sea to Cretaceous Orogen - The Austroalpine Sector of the Tethyan Shelf (Eastern Alps, Austria), and by Kozur, H. W. (Ladinian and Carnian palaeogeography of southern Turkey and its importance for the development of the Triassic Tethyan faunal provinces).

Sedimentology, correlations, paleoecology and palynology of the Permian-Triassic boundary interval were the subject of 4 presentations: - Brookfield, M. (Sedimentology of the Permo-Triassic boundary sections in Kashmir, India), - Twitchett, R. J. (High resolution, global correlation

of the Permian-Triassic interval), - Twitchett, R. J. and Looy, C. V. (Rapid and synchronous collapse of end-Permian marine and terrestrial ecosystems), - Spina, A., Cirilli, S and Baud, A. (Palynology of the uppermost Permian - basal Triassic successions in the High Arctic (Canada) and comparison with some PTB Gondwanian localities).

All the abstracts of the Pangea Symposium are available on the Web at <http://www.geoconfoman.unibe.ch/> and have been published in the last issue of GeoArabia.

The Symposium was followed on Jan. 17 by a 4-day **post-conference Excursion (B01)** on the Permo-Triassic Deposits from Shallow Water to Base of Slope and basin leaded by A. Baud, F. Bechenec, F. Cordey, J. Marcoux, R. Maury, J. le Metour and S. Richoz.

New data on Lower Triassic succession (Member C, Maqam Formation) were presented by S. Richoz and A. Baud in the Wadi Maqam and Wadi Shuayab. It consists of a very thick unit (900m) essentially of platy limestones, calcarenites and calcirudites. It comprises mainly grey-beige calcilutite, laminated and flaggy, interbedded with sparse beds of fine-grained calcarenite in cm beds. Channelizing beds of intraformational calcirudite are also part in this succession which constitutes the great part of the outcrops of the Sumeini Group. The low negative values of the Carbon isotope within this thick Lower Triassic unit contrast with the high positive values found in the underlying Upper Permian Member B (shift of 6 ‰ at the Permian-Triassic transition).

## **The Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events, Changxing, China, August 10-13, 2001**

**Yin Hongfu<sup>1</sup>, Tong Jinnan<sup>2</sup>**

<sup>1</sup> *China University of Geosciences, Wuhan 430074, China, hfyin@cug.edu.cn*, <sup>2</sup> *China University of Geosciences, Wuhan 430074, China, jntong@cug.edu.cn*

The Paleozoic-Mesozoic transition was the greatest geological turning point during Phanerozoic history, yielding the biggest biotic mass extinction and other extraordinary fatal events. So it has long been a hot topic for geologists and the Permian-Triassic boundary has received considerable attention. After the untiring and arduous effort of Permian-Triassic boundary geologists for decades and of others for over twenty years, the Global Stratotype Section and Point (GSSP) of the Permian-Triassic Boundary (PTB) was finally ratified at the base of Bed 27c of Meishan Section D, Changxing County, Zhejiang Province, South China by the International Union of Geological Sciences

(IUGS) in early March 2001. This marks a great progress in the study of Permian-Triassic boundary stratigraphy and geological events. As a result of the studies on the Permian-Triassic boundary and transitional events and geological processes over the world there is now a unitary chronostratigraphic framework.

The international Symposium on the GSSP of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events was held in Changxing County, Zhejiang Province, China on August 10-13, 2001, chaired by Professor Yin Hongfu, the leader of the Permian-Triassic Boundary Working Group (PTBWG), co-organized by China University of Geosciences, Nanjing Institute of Geology and Paleontology, China Geological Survey and the Government of Changxing County, and co-sponsored by China University of Geosciences, Nanjing Institute of Geology and Paleontology, National Natural Science Foundation of China, China Geological Survey, China National Committee of Stratigraphy, Paleontological Society of China, Land and Resources Bureau of Zhejiang Province, Subcommittee of Triassic Stratigraphy (STS), Subcommittee of Permian Stratigraphy (SPS) and the Global Sedimentary Geological Program (GSGP). A formal ceremony unveiling the monument marking the PTB GSSP took place on site at Meishan Section D during the symposium. About



*Picture 1: The opening of the International Symposium on the GSSP of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events*

60 geologists from 13 countries attended the scientific activities and over 100 other members from the national and local governmental offices and news offices presented the opening and monument ceremonies. The vice-chairman of the International Commission of Stratigraphy (ICS) (Richard Lane), the vice-minister of Ministry of Land and Resources and the director of China National Committee of Stratigraphy (Shou Jiahua), the vice-director of the Head Office of the National Environment Protection (Song Ruixiang), the vice-governor of the Zhejiang Province (Zhang Mengjin) and others were specially invited to present at the symposium and/or ceremonies.

The opening of the symposium (Picture 1) was first held in the downtown of Changxing County during the morning of August 10. Professor Yang Zunyi, 94 years old, chaired the opening and briefly introduced the history of the decision of the GSSP at Meishan. He expressed his excitement and appreciation for the final decision of the PTB GSSP after an extensive, painstaking, and thorough comparison and selection process. He said, "As you know, in the early years the ammonoid *Otoceras* was considered as the index fossil of the Permian-Triassic boundary. In 1986, the conodont *Hindeodus parvus* was proposed to substitute *Otoceras* as the boundary marker, which later obtained the majority approval of the Permian-Triassic Boundary Working Group. In 1993, the PTBWG selected four candidates for the stratotype of the boundary, i.e., Meishan of Zhejiang, Guryul Ravine of Kashmir, Shangsi of Sichuan, and Selong of Tibet. In 1996 nine members of the PTBWG published a formal recommendation to set the Permian-Triassic boundary at the first appearance of *Hindeodus parvus*, Bed 27c of Meishan. From 1999 to 2000, the proposal for Meishan as the GSSP of the Permian-Triassic boundary passed three separate ballots. In March of this year, the IUGS Executive Committee finally ratified that the GSSP of the Permian-Triassic Boundary be defined at the base of Bed 27c, Meishan Section D, Changxing County, Zhejiang Province, China, at the horizon where the conodont *Hindeodus parvus* first appeared." Subsequently, some leaders from the international scientific organizations and from the national and local governmental offices addressed their congratulations to the opening of the symposium.

After the opening of the symposium, all conventioners immediately took buses to Meishan Section D to present the ceremony of unveiling the monument marking the PTB GSSP on site. The monument ceremony was chaired by the leader of the PTBWG, Professor Yin Hongfu. He said, "According to the regulation of the International Com-



Picture 2: The GSSP monument of the Permian-Triassic boundary at Meishan Section D

mission of Stratigraphy, the type-locality of the GSSP should be open to all scientists for investigation and sampling regardless of their nationality, political attitude, race or religion, and a preservation stele or monument should be established on the site in order to protect the stratotype. The first demand has been satisfied by the document of State Council of China, September 17, 1999, which declared the openness of the whole Changxing County, including Meishan area, to foreigners. Now we will fulfill the second demand by establishing and unveiling the protection monument."

The chairman of the STS, Dr. Michael Orchard, and the head of the Changxing County declared the importance of the Meishan section and its protection, separately. Richard Lane, Shou Jiahua, Song Ruixiang and Zhang Mengjin together unveiled the monument.

The monument of the PTB GSSP is 9.09 m high and composed of three parts: the base, the body and the top (Picture 2). The body is bipartite. The lower part has two blocks, meaning Dyas in Ger-

man, now called Permian. The upper part has three long columns, implying Trias in German, now called Triassic. The top consists of a “golden spike” wedg-

column are the Chinese translation of the PTB GSSP and the three organizations in charge of it. The carved characters on the lower body records

Picture 3 Group photo in front of the GSSP of the Permian-Triassic boundary



ing into the rocks and the index fossil of the spike—*Hindeodus parvus*. The carved characters on the

the main research units, the supporting organizations of the research and the monument. At the



Picture 4 The protection stele of the Changxing Limestone

back is a diagram showing the multidisciplinary correlation of the PTB strata. In front of the monument is a marble stone, where carved in Chinese, is a record of the course of the choice of this GSSP. A group photo was taken thereafter in front of the monument for all symposium participants and the official leaders as a souvenir (Picture 3).

To celebrate the completion of the monument and the opening of the symposium, a Commemoration Envelope of the GSSP of the Permian and Triassic Boundary was co-issued by the Philately Company of Zhejiang Province, the Government of Changxing County and the Post-office of Changxing County. A brief ceremony for the first-day cover and the signature followed the opening.

The scientific sessions started from the afternoon of August 10 on. This symposium received and printed 47 abstracts and 33 papers were orally reported during the symposium, including 9 keynotes. Yin Hongfu first summarized the basic geology and achievements of the Meishan section and the GSSP of PTB, and then the scientific sessions went on. The reports and discussions focussed on Permian-Triassic boundary geology, dealing with all aspects related to the boundary and the transition. The contents and achievements of the reports and discussions are roughly summarized below:

1. Boundary chronology: Dr. Roland Mundil re-

ported new ages for the clay beds around the Permian-Triassic boundary of the Meishan section based on the U/Pb dating of zircons. The age of the Permian-Triassic boundary at the GSSP is suggested to be 253 Ma. The method is believed to be more reliable and the result is proven by their research at the Shangsi section. This result, if proved by re-investigations, will cast significant influence on the duration of the Upper Permian, Lower Triassic and the Permian-Triassic crisis.

2. Boundary events: Research on the Permian-Triassic boundary has yielded new evidence. The quantitative statistics on a database of fossils, the discovery of fullerenes ( $C_{60}$  and  $C_{70}$ ), and anomalous sulfur and strontium isotope excursions, and new curves of carbon isotopes, as well as new dating on the boundary strata, compel us to further consider the causes of the great transitional events. A few papers especially suggested an extraterrestrial origin for the mass extinction.

3. Boundary correlation: Many reports dealt with new data at the Permian-Triassic boundary from various regions around the world, including Arctic Canada, Norway, Russia, Japan, Thailand, Iran, Saudi Arabia, Oman, and Vietnam as well as several provinces and Tibet of South China. The new data are correlated to the standard sequence at the Meishan section and support the boundary data as well as demonstrate a unitary boundary strati-



*Picture 5 Steps to the base of the Changxing Limestone at Section D*

graphic pattern. However, the magnetostratigraphic sequence of the terrestrial Permian-Triassic boundary strata in Xinjiang reported by Dr. John Lyons is clearly different from the marine sequence.

4. Transitional biota: Besides the mass extinction pattern at the Permian-Triassic boundary, the evolutionary processes and lineages of several characteristic taxa were examined, for example, conodont lineages and paleogeographic provincialism, radiolarian extinction at the end of the Permian and recovery in the early Triassic as well as a turnover between the Early and Middle Triassic, the discovery of Triassic vertebrates in Guizhou and their radiation during the Middle Triassic. It is notable that the restudy of *Reduviasporonites* redefines and opposes a disastrous fungal event.

5. Stratigraphy below and above the boundary: Some reports addressed advances in the study of the Lopingian and Changhsingian below the Permian-Triassic boundary and the Lower Triassic above the boundary. The bases of the Wuchiapingian and Changhsingian stages of the Lopingian and the Induan-Olenekian boundary were discussed. Many reports presented important advances in Lopingian and Lower Triassic conodont and ammonoid biostratigraphy, the biostratigraphic correlation between the Boreal and

Tethyan sequences, and of new knowledge on the continuation of some stratigraphic sequences in the new biostratigraphic framework.

Meanwhile, studies indicate that the reconstruction of the Permian and Triassic eastern Tethys shows an archipelagic ocean full of islands, remarkably different from previous reconstructions. The paleoclimatic and paleoceanographic interpretations thus have to be quite different from those previously published. A lecture was also presented on Carboniferous conodont events prior to the formation of Pangea and the tectono-stratigraphy surrounding the North American transcontinental arch.

During the symposium, STS and SPS business meetings were conducted and a conodont workshop was organized. The SPS meeting was chaired by the vice-chairman of the SPS, Dr. Clinton Bruce Foster. At the meeting Dr. Charles Henderson, the secretary-general of the SPS, reported on the work of the SPS since the 31st IGC, its plan for the future and on preparations for attending the 32nd IGC. Recent advances and results in the study of the bases of the Lopingian and the Changhsingian were addressed and discussed. It was approved that unnecessary controversies could be avoided and that the Guadalupian-Lopingian boundary should be fixed against time soon.



Picture 6 The bed numbers carved on rocks at the boundary of Section D

The STS meeting was chaired by Dr. Michael Orchard, chairman of the STS. The meeting emphasized the boundaries between the stages within the Triassic. The base of the Triassic (Induan Stage) has been fixed, which provides an excellent example for the other Triassic GSSPs. The base of the Olenekian faces problems in that the stage is from the Boreal biogeographic realm and the proposed candidates of the boundary from the Vladivostok area may not meet the GSSP requirements. Thus it was suggested that the stage be re-selected. South China qualifies for this GSSP. The base of the Anisian is expected to be the second Triassic GSSP in the near future, but the work in Romania should be quickened. The tuff beds around the Lower and Middle Triassic boundary in South China would provide age data for this boundary. The base of the Ladinian has been extensively argued in terms of three different markers. As lots of work has been done at this boundary, it was advised that a vote could be done sooner. However, it is feared that the ballot may not be successful if no predominant selection is achieved before the vote. The base of the Carnian has been studied in Italy and Spiti and some good results have been achieved recently, but the pace of this work should also be quickened. The base of the Norian has seen little work so far, but a new section in North America is quite good for biostratigra-

phy although the magnetostratigraphy is not reliable due to an overprint. The base of the Rhaetian is to be done in the future.

The conodont workshop was chaired by Drs. Mei Shilong and Charles Henderson. Conodonts become the key fossils in the definition of the boundaries between and within the Permian and Triassic, but the taxonomic position of some conodonts is as yet indeterminate and the classification of conodonts is under dispute. A classification scheme based upon population features was proposed and discussed and it was confirmed to be valuable for Permian and Triassic conodont studies. The conodonts marking the Permian-Triassic boundary and its lineage were reviewed as well.

The mid-symposium field excursion provided all participants an opportunity to closely examine all of the Meishan sections. Section D, where the PTB GSSP is located, is also the location where the entire Changxing Limestone outcrops; it was protected with a stele by the governments early in the 80's (Picture 4). Here, all but the base and top of the Changxing Limestone are in the middle part of the hillside, which causes trouble for observation and investigation. With the cooperation of the local government, the Meishan section has been entirely repaired and protected. The boundaries

were especially revealed by the preparation and a series of firm cemented steps extends to the boundaries (Picture 5) and the standard lithostratigraphic bed numbers were carved on the rocks (Picture 6), which is very convenient for observation and for research at the section. Now the boundary strata of Section D (the PTB GSSP) are strictly protected. In the west part, about 80 m from the GSSP, a section is exposed for sampling to meet the needs of researchers. As the case stands, the Permian-Triassic boundary in the Meishan area extends along the hillside and it is exposed within a series of abandoned quarries. For this excursion the local government built a road along the foot of the hill to run through all the sections. Therefore, we could trace the Permian-Triassic boundary westward from Section D to Section A and achieve a complete knowledge on the standard boundary sequence. This would be of great importance for the extension of the PTB GSSP around the world.

For better understanding the PTB GSSP at the Meishan section, the China University of Geosciences professionally designed 24 poster pages showing the major achievements of various studies at the Meishan section and posted them during the symposium. Meanwhile, the China University of Geosciences and the Nanjing Institute of Geology and Paleontology jointly collected and exhibited hundreds of publications on the Permian and Triassic boundary and related studies that were accomplished by both institutions during various stages, as well as some of the most important fossil specimens from the Meishan section. So it provided the best opportunity for extensively understanding the Meishan section and the Permian-Triassic boundary sequence. This formed crucial parts of the symposium together with the field excursions.

The pre-symposium field excursion was executed on August 8-9 and nine participants including the guiders attended this tour to view the regional stratigraphy in the Meishan area. Four sites were scheduled to examine Carboniferous to Triassic sequences and some key boundaries within them, the variation of Changhsingian facies, and the spatial extension of Permian-Triassic boundary strata. From the excursion, the specialists were convinced of the superiority of the Permian-Triassic boundary sequence at Meishan for the regional geologic setting.

The post-symposium field excursion to Chaohu, Anhui Province had 15 attendee, including the chairman of the STS, Dr. Michael Orchard, the vice-chairman of the STS, Prof. Yin Hongfu (vice-chairman of the China National Committee of Stratigra-

phy), the vice-chairman of the STS, Dr. Yuri Zakharov (leader of the Induan-Olenekian Boundary Working Group), and the secretary-general of the SPS, Dr. Charles Henderson. The excursion had five stops. The first stop dealt with the Carboniferous and Permian boundary with emphasis on the difference between the global and the traditional Chinese Carboniferous-Permian boundaries. The Permian stratigraphic sequence was reviewed as well. The other four stops focussed on the Upper Permian and Lower Triassic. They were especially designed to closely examine the whole Lower Triassic sequence, fossil assemblages, sedimentary facies and key boundaries such as the Permian-Triassic, Induan-Olenekian and Lower and Middle Triassic boundaries. The Induan-Olenekian boundary received considerable attention and discussion. It is believed that the Lower Triassic of Chaohu is quite competent as a potential GSSP for the Induan-Olenekian boundary, because: (1) the Lower Triassic sections have a good geographic condition with excellent traffic and good stratigraphic exposure on the roadsides. The sections are in the suburb of the Chaohu City, only 3 km from the downtown. Chaohu is a medium-sized city of the Anhui Province, 60 km from Hefei, the capital of the Anhui Province, communicated by railway and freeway. Moreover, it is only 90 km from Chaohu to Nanjing, the capital of the Jiangsu Province; (2) the Lower Triassic of Chaohu has abundant fossils. Most of the crucial ammonoids and conodonts have been collected so that the boundary could be clearly defined and correlated. In the meantime, the sections yield rich bivalves and some vertebrates and they are important sites producing rare reptile fossils; and (3) the Lower Triassic sequence at Chaohu is complete. The sedimentary sequences almost could be correlated with those of North America in terms of sequence stratigraphy. The cyclic sediments are ideal for high-resolution cyclostratigraphic study.

Two groups of 14 scientists went to Guizhou for the Permian-Triassic boundary sequences from the marine to continental facies after the symposium in Changxing. One group of five men went alone, as they needed more time to do field work there. The other group of nine attendees went on the scheduled post-symposium field excursion to Guizhou. The viewed sections were far from each other and in different facies so that the excursion was quite long. As it is in a mountainous area in the western Guizhou, the traffic was also slow. At the same time, many parts of the highway are currently under construction due to the national program of developing the west in China, which makes the traffic even slower, so that we had to cancel one previously scheduled stop. The marine Per-



mian-Triassic boundary section in Zhongzhai of the Liuzhi County was examined. The lithology at the boundary is entirely different from that at the Meishan section and mainly composed of clastic rocks. But it contains rich marine fossils similar to those in Meishan. The boundary stratigraphic sequence of clay beds and boundary limestone at the Meishan section was also observed here except that the boundary limestone here contains more arenaceous elements. It is a marine sequence with a different facies (littoral) from the Meishan section, that represent transitional facies from marine to continental, so it was quite attractive and interesting to the visitors.

The observed continental Permian-Triassic boundary sequence is the Zhejue section in the Weining County at the border between the Guizhou and Yunnan provinces. The boundary strata consist entirely of continental sediments yielding mainly plant fossils except for the boundary clay beds, which is intercalated by a medium bed of sandstone to form a three-bed sequence similar to the boundary sequence at the Meishan section. Although the boundary correlation based only upon the three-bed boundary sequence might be quite arbitrary and adventurous, it is convincing that this is an instructive effort.

### Acknowledgements

The symposium was financially supported by China University of Geosciences (Wuhan), Nanjing Institute of Geology and Paleontology, China Geological Survey, China National Committee of Stratigraphy, National Natural Science Foundation of China and the Government of Changxing County. The GSSP monument was financed by the National Natural Science Foundation of China, the Ministry of Science and Technology, China Geological Survey, China University of Geosciences and the Land and Resources Office of Zhejiang Province. We extend appreciation to Drs. Michael Orchard and Charles Henderson for their kind comments and revisions to an earlier draft manuscript of this report.

## Report on the field excursion in western Guizhou and eastern Yunnan

Peng Yuanqiao<sup>1,3</sup>, Wang Shangyan<sup>2</sup> and Gu Songzhu<sup>1</sup>

<sup>1</sup> Faculty of Earth Science, China University of Geosciences, Wuhan 430074, Hubei, China,

<sup>2</sup> Guizhou Geological Survey, Guiyang 550004, Guizhou, China

<sup>3</sup>corrsp. author: digushi@cug.edu.cn

After the International Symposium on the Global Stratotype Section and Point (GSSP) of the Permian-Triassic boundary (PTB) and the Paleozoic-Mesozoic Events, nine scholars from home and abroad took part in the field excursion on the PTB strata from marine to land in western Guizhou and

eastern Yunnan on 14-18 August, 2001. The field excursion was lead by Dr. Peng Yuanqiao (CUG), Dr. Wang Shangyan (GGS) and Dr. Gu Songzhu (CUG). Participants included Dr. Aymon Baud (Geological Museum of Switzerland) and his wife Monique Baud, Dr. Tadeusz Peryt (Polish Geological Institute), Dr. Takeshi Ishibashi (Kyushu University of Japan), Dr. Kunio Kaiho (Tohoko University of Japan) and Dr. Cao Changqun (Nanjing Institute of Geology and Palaeontology, Academia Sinica). This field party examined two kinds of facies of the PTB strata: the marine Permian and Triassic of Liuzhi area, Guizhou and the terrestrial Permian and Triassic of Weining area, Guizhou.

Western Guizhou and eastern Yunnan are in the southwestern part of China, to the south of Sichuan and to the north of Guangxi (Fig.1). The area lies in the east part of Yunnan and Guizhou plateau with the average elevation of 1000 meters. Typical Karst landforms are widely developed in the whole area. Road access is improving through a large-




Figure 1 Geographical setting of western Guizhou and eastern Yunnan  
1-Laibin section; 2-Zhejue section; 3-Chahe section; 4-Zhongzai section; 5-Jiaozishan section, 1, 2 and 3 are terrestrial PTB sections, 4 is neritic clastic PTB section, 5 is neritic carbonate PTB section;



Picture 1 The Permian-Triassic boundary at Stop 2 (North Pingdingshan Section)

scale Road Refit Project currently in progress.

Two sections were arranged to be surveyed for this five days' field excursion: the marine Zhongzai section, Liuzhi, Guizhou Province and the terrestrial Zhejue section, Weining, Guizhou Province.

Stop 1: Zhongzai section, Liuzhi, Guizhou Province—neritic clastic PTB section

Zhongzai section is located in a road-cut at the north part of Zhongzai village (or Heilaga village), Langdai, Liuzhi County, Guizhou Province. The well exposed Upper Permian and the Lower Triassic succession differ lithologically from the neritic carbonate PTB Meishan section — the Global Stratotype Section and Point (GSSP) of the PTB. The lithologies near the PTB at the Zhonghai section are composed of clastics containing abundant fossils, but the PTB sequence of clay beds and the corresponding boundary limestone (sandstone) is clear. This section possesses shore facies characteristics and should be the key for the high-resolution correlation from marine to land.

Stop 2: Zhejue section, Weining, Guizhou Province—terrestrial PTB section

Zhejue section is located at Gongping village, Zhejue town, Weining County, Guizhou Province. It is cut by national highway 326 from Weining to Xuanwei. The Upper Permian and the Lowermost Triassic are well exposed and consist of terrestrial clastics containing abundant plant fossils. At this section, the PTB is marked by two clay beds with the same structure and event sequence as that of the Meishan section. The main difference between the two sections is that the boundary limestone at the Meishan section is replaced by a sandstone bed at the Zhejue section. Although the high-resolution correlation of the boundary sequence at the two sections needs to be strengthened by further study in biostratigraphy and magnetostratigraphy, it is believed that the Zhejue section could serve as an Accessory Section and Point (ASP) of the PTB. In its favour are: (1) good stratigraphic exposure and accessibility, (2) abundant fossils including plants and palynomorphs, (3) similarity of the stratigraphic set compared with the marine PTB sequence at Meishan section and (4) the boundary clay beds formed by volcanism at Zhejue section that might provide us opportunity to test the direct age for the terrestrial PTB.



*Picture 2 The Stratigraphic sequence around the Induan-Olenekian boundary (and sequence surface) at Stop 2 (North Pingdingshan Section)*



*Picture 3 The Stratigraphic sequence around the Induan-Olenekian boundary at Stop 3 (West Pingdingshan Section)*



## New Triassic Literature

### Triassic Bibliography

**G. Warrington, Kerp H., Kuerschner, W.M., Lucas, S.**

- Abdala, F., 2000. Catalogue of non-mammalian cynodonts in the Vertebrate Paleontology Collection of the Instituto Miguel Lillo, Universidad Nacional de Tucuman, with comments on species. *Ameghiniana*, 37: 463-476.
- Acheche, M. H., M'Rabet, A., Ghariani, H., Ouahchi, A. & Montgomery, S. L. 2001. Ghadames, basin, southern Tunisia: a reappraisal of Triassic reservoirs and future prospectivity. *AAPG Bulletin*, 85 (5): 765-780.
- Ahlberg, A. & Olsson, I. 2001. Petroleum assessment of the Mesozoic succession in the Höllviken Graben and on the Skurup Platform, southern Sweden. *GFF*, 123: 85-95.
- Anon. 2001. MICRO-ORGANISM uncertainty casts doubt on eXtinction theory. *AusGEO News* 61, 8-9.
- Árkai, P. Alpine regional metamorphism in the main tectonic units of Hungary: a review. *Acta Geologica Hungarica*, 44 (2-3): 329-344.
- Balini, M., Krystyn, L., Nicora, A. & Torti, V. 2001. The Ladinian-Carnian boundary succession in Spiti (Tethys Himalaya) and its bearing on the definition of the GSSP for the Carnian stage (Upper Triassic). *Journal of Asian Earth Sciences*, 19 (3A): 3-4.
- Balogh, K. & Pécskay, Z. 2001. K/Ar and Ar/Ar geochronological studies in the Pannonian-Carpathians-Dinarides (PANCARDI) region. *Acta Geologica Hungarica*, 44 (2-3): 281-299.
- Battail, B. & Surkov, M. V. 2000. Mammal-like reptiles from Russia. Pp.86-119 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Bédir, M., Boukadi, N., Tlig, S., Ben Timzal, F., Zitouni, L., Alouani, R., Slimane, F., Bobier, C. & Zargouni, F. 2001. Subsurface Mesozoic basins in the central Atlas of Tunisia: tectonics, sequence deposit distribution, and hydrocarbon potential. *AAPG Bulletin*, 85 (5): 885-907.
- Benatov, S. 1999. Fasnian ammonoids and Anisian-Ladinian boundary in western Bulgaria. *Review of the Bulgarian Geological Society*, 60 (1-3): 29-37.
- Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds). 2000. *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press, xxxix+696pp.
- Beyer, K. & Lundschein, B. A. 2000. Magnetostratigraphy as a potential tool for correlation in the Murzuq Basin illustrated by an example from the Triassic Snorre Reservoir in the Northern North Sea. Pp.17-29 in Sola, M. A. & Worsley, D. (eds) *Geological Exploration in Murzuq Basin*. Proceedings of the geological conference on exploration in the Murzuq Basin, Sabah, 20-22 September, 1998. Elsevier
- Blein, O., Lapierre, H., Schweikert, R. A., Pécher, A. & Reynaud, C. 2001. Volcanisme triasique calco-alcalin à shoshonitique du Nevada occidental. *Bulletin de la Société géologique de France*, 172 (2): 189-200.
- Blendinger, W. 2001. Triassic carbonate buildup flanks in the Dolomites, northern Italy: breccias, boulder fabric and the importance of early diagenesis. *Sedimentology*, 48: 919-933.
- Bohacs, K. M., Carroll, A. R., Neal, J. E. & Mankiewicz, P. J. 2000. Lake-basin type, source potential, and hydrocarbon character: an integrated sequence-stratigraphic-geochemical framework. *AAPG Studies in Geology*, 46: 3-34.
- Bonaparte, J. F., Ferigolo, J. & Ribeiro, A. M. 2001. A primitive Late Triassic 'ictidosaur' from Rio Grande do Sul, Brazil. *Palaeontology*, 44 (4): 623-635.
- Boncheva, I. 1999. Conodont data on the age of a part of the sediments in the scope of the Stakevci Formation, (northwest Bulgaria). *Review of the Bulgarian Geological Society*, 60 (1-3): 25-28.
- Boulvain, F., Belanger, I., Delsate, D., Dosquet, D., Ghysel, P., Godefroit, P., Laloux, M., Roche, M., Teerlynck, H. & Thorez, J. 2000. New lithostratigraphic, sedimentological, mineralogical and palaeontological data on the Mesozoic of Belgian Lorraine: a progress report. *Geologica Belgica*, 3 (1-2): 3-33.

- Brew, G., Barazangi, M., Al-Maleh, A. K. & Sawaf, T. 2001. Tectonic and geologic evolution of Syria. *GeoArabia*, 6 (4): 573-616.
- Buckovic, D., Jelaska, V., Tesovic, B.C., 2001. Facies variability in Lower Liassic carbonate successions of the Western Dinarides (Croatia). *Facies*, 44: 151-162.
- Budai, T., Csillag, G., Vörös, A. & Dosztály, L. 2001. Middle to Late Triassic platform and basin facies of the Veszprém Plateau (Transdanubian Range, Hungary). *Földtani Közlöny*, 131 (1-2): 37-70.
- Budai, T., Csillag, G., Vörös, A. & Lelkes, G. 2001. Middle to Late Triassic platform and basin facies of the Eastern Bakony Mts. (Transdanubian Range, Hungary). *Földtani Közlöny*, 131 (1-2): 71-95.
- Carrano, M. T. & Wilson, J. A. 2001. Taxon distribution and the tetrapod track record. *Paleobiology*, 27 (3): 564-582.
- Carrillat, A. 2001. Palaeoenvironmental reconstruction of the Mufara Formation (Upper Triassic, Sicily): biostratigraphy, organic facies, sedimentological and geochemical approach. *Terre & Environnement*, 27: 1-269. Section des Sciences de la Terre, Université de Genève.
- Carter, A., Roques, D., Bristow, C. & Kinny, P. 2001. Understanding Mesozoic accretion in southeast Asia: significance of Triassic thermotectonism (Indosinian orogeny) in Vietnam. *Geology*, 29 (3): 211-214.
- Cartwright, J., Stewart, S. & Clark, J. 2001. Salt dissolution and salt-related deformation of the Forth Approaches Basin, UK North Sea. *Marine and Petroleum Geology*, 18: 757-778.
- Catuneanu, O. & Elango, H. N. 2001. Tectonic control on fluvial styles: the Balfour Formation of the Karoo Basin, South Africa. *Sedimentary Geology*, 140: 291-313.
- Çerikcioglu, B. 2001. An approach to the origin of Kocaeli Triassic dolomites based on their isotopic and petrographic characteristics. *Geological Bulletin of Turkey*, 44 (1): 37-44.
- Chalbaoui, M., Ben Dhia, H. & Turki, M. 2000. Nouvelles données sur la tectonique de la région des Chotts de Tunisie du Trias au Crétacé inférieur. *Africa Geoscience Review*, 7 (4): 355-363.
- Charlton, T. R. 2001. Permo-Triassic evolution of Gondwanan eastern Indonesia, and the final Mesozoic separation of SE Asia from Australia. *Journal of Asian Earth Sciences*, 19: 595-617.
- Chatalov, A. 2000. Marine phreatic cements in the Triassic limestones from the western Balkanides. *Geologica Balcanica*, 30 (1-2): 33-48.
- Chatalov, A. 2001. Deformational structures in the Iskur Carbonate Group (Lower – Upper Triassic) from the Western Balkanides. *Geologica Balcanica*, 30 (3-4): 43-57.
- Chatalov, A. & Stanimirova, T. 2001. Diagenesis of Spathian-Anisian dolomites from the western Balkanides, Bulgaria. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, 2001 (5): 301-320.
- Chikhaoui, M., Braham, A., Turki, M.M., 2001. Evidence of the non-interbedding of the Triassic evaporites within the Turonian sediments in the Koudiat Sidii area (north-western Tunisia). *Comptes Rendus de l'Académie des Sciences Sér., II Fasc. A - Sciences de la Terre et des Planètes*, 332: 689-694.
- Clark, J.M., Sues, H.D., Berman, D.S., 2000. A new specimen of *Hesperosuchus agilis* from the Upper Triassic of New Mexico and the interrelationships of basal crocodylomorph archosaurs. *Journal of Vertebrate Paleontology*, 20: 683-704.
- Colbert, E. H. & Olsen, P. 2001. A new and unusual aquatic reptile from the Lockatong Formation of New Jersey (Late Triassic, Newark Supergroup). *American Museum Novitates*, No.3334: 24pp.
- Damiani, R. J., Neveling, J. & Hancox, P. J. 2001. First record of a mastodontosaurid (Temnospondyli, Stereospondyli) from the Early Triassic *Lystrosaurus* Assemblage Zone (Karoo Basin) of South Africa. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (1): 133-144.
- Danelian, T. & Robertson, A. H. F. 2001. Neotethyan evolution of eastern Greece (Pagondas Mélange, Evia island) inferred from radiolarian biostratigraphy and the geochemistry of associated extrusive rocks. *Geological Magazine*, 138 (3): 345-363.
- Darby, B. J., Wyld, S. J. & Gehrels, G. E. 2000. Provenance and paleogeography of the Black Rock terrane, northwestern Nevada: implications of U-Pb detrital zircon geochronology. *Geologi-*

- cal Society of America, Special Paper 347: 77-87.
- De, C. 2001. Marine influence in the Indian Gondwana basins: a review of ichnological findings. *Journal of the Geological Society of India*, 57: 411-415.
- Dickinson, W. R. 2000. Geodynamic interpretation of Paleozoic tectonic trends oriented oblique to the Mesozoic Klamath-Sierran continental margin in California. *Geological Society of America, Special Paper 347*: 209-245.
- Dickinson, W. R. & Gehrels, G. E. 2000. Sandstone petrofacies of detrital zircon samples from Paleozoic and Triassic strata in suspect terranes of northern Nevada and California. *Geological Society of America, Special Paper 347*: 151-171.
- Dickinson, W. R. & Lawton, T. F. 2001. Carboniferous to Cretaceous assembly and fragmentation of Mexico. *Geological Society of America Bulletin*, 113 (9): 1142-1160.
- Dill, H. G. & Dultz, S. 2001. Chemical facies and proximity indicators of continental – marine sediments (Triassic to Liassic, SE Germany). *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (3): 289-324.
- DiPietro, J.A., Isachsen, C.E., 2001. U-Pb zircon ages from the Indian plate in northwest Pakistan and their significance to Himalayan and pre-Himalayan geologic history. *Tectonics*, 20: 510-525.
- Dolenec, T. & Lojen, S. 2000. Ce anomaly at the Permian-Triassic boundary in the Idrijca Valley as evidence of changing redox conditions at the P/Tr transition in the western Tethys (Slovenia). *Geologija*, 43 (1): 103-107.
- Dolenec, T., Lojen, S. & Ramovs, A. 2001. The Permian-Triassic boundary in western Slovenia (Idrijca Valley section): magnetostratigraphy, stable isotopes and elemental variations. *Chemical Geology*, 175: 175-190.
- Dorka, M. 2001. Shark remains from the Triassic of Schöningen, Lower Saxony, Germany. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (2): 219-247.
- Dornsiepen, U. F., Manutsoglu, E. & Mertmann, D. 2001. Permian-Triassic palaeogeography of the external Hellenides. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 172: 327-338.
- Dypvik, H., Hankel, O., Nilsen, O., Kaaya, C. & Kilembe, E. 2001. The lithostratigraphy of the Karoo Supergroup in the Kilombero Rift Valley, Tanzania. *Journal of African Earth Sciences*, 32 (3): 451-470.
- Dzik, J., 2001. A new Paleorhinus fauna in the early Late Triassic of Poland. *Journal of Vertebrate Paleontology*, 21 (3): 625-627.
- Evans, S. E. 2001. The early Triassic 'lizard' *Colubrifer campi*: a reassessment. *Palaeontology*, 44 (5): 1033-1041.
- Ermakova, S. P., 2001. Description and taxonomy of a new Early Triassic Genus *Eovavilovites* (Ammonoidea, Ceratitida). *Paleontological Journal*, 35: 259-261.
- Fodor, L. & Koroknai, B. 2000. Ductile deformation and revised lithostratigraphy of the Martonyi subunit (Torna unit, Rudabánya Mts.), northeastern Hungary. *Geologica Carpathica*, 51 (6): 355-369.
- Franzese, J. R. & Spalletti, L. A. 2001. Late Triassic-Early Jurassic continental extension in southwestern Gondwana: tectonic segmentation and pre-break-up rifting. *Journal of South American Earth Sciences*, 14: 257-270.
- Frixa, A., Bertamoni, M., Catrullo, D., Trincianti, E. & Miuccio, G. 2000. Late Norian-Hettangian paleogeography in the area between wells Noto 1 and Polpo 1 (S-E Sicily). *Memorie della Società Geologica Italiana*, 55: 279-284.
- Gehrels, G. E. 2000. Introduction to detrital zircon studies of Paleozoic and Triassic strata in western Nevada and northern California. *Geological Society of America, Special Paper 347*: 1-17.
- Gehrels, G. E., Dickinson, W. R., Darby, B. J., Harding, J. P., Manuszak, J. D., Riley, B. C. D. & Spurlin, M. S. 2000. Tectonic implications of detrital zircon data from Paleozoic and Triassic strata in western Nevada and northern California. *Geological Society of America, Special Paper 347*: 133-150.
- Ghosh, P., Ghosh, P. & Bhattacharya, S. K. 2001. CO<sub>2</sub> levels in the Late Palaeozoic and Mesozoic atmosphere from soil carbonate and organic matter, Satpura basin, Central India. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170: 219-236.



- Gialanella, P. R., Heller, F., Mietto, P., Incoronato, A., De Zanche, V., Gianolla, P. & Roghi, G. 2001. Magnetostratigraphy and biostratigraphy of the Middle Triassic Margon section (Southern Alps, Italy). *Earth and Planetary Science Letters*, 187: 17-25.
- Gierlowski-Kordesch, E. H. & Kelts, K. R. (eds). 2000. *Lake basins through space and time*. AAPG Studies in Geology, 46.
- Gillhaus, A. & Richter, D. V. 2001. Polyphase Dolomitgenese in oberpermischen und obertriassischen Sabkha-Kleinzyklen von Hydra (Griechenland). *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, 2001 (7): 399-422.
- Góczán, F. 2001. The *Classopollis visscherii* sp.nov. *Földtani Közlöny*, 131 (1-2): 97-106.
- Göncüoğlu, M. C., Tekin, U. K. & Turhan, N. 2001. Late Carnian radiolarite-bearing basalt blocks within the Late Cretaceous Central Sakarya ophiolitic melange, NW Anatolia: geological constraints (Abstract). 54<sup>th</sup> Geological Congress of Turkey, Ankara, May 7-10, 2001. Abstracts and Proceedings: 9-10.
- Gower, D. J. & Sennikov, A. G. 2000. Early archosaurs from Russia. Pp.140-159 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Grabowski, J. 2000. Palaeo-and rock magnetism of Mesozoic carbonate rocks in the sub-Tatric Series (central West Carpathians) – palaeotectonic implications. *Polish Geological Institute Special Papers*, 5: 1-8
- Groenewald, G. H., Welman, J. & MacEachern, J. 2001. Vertebrate burrow complexes from the Early Triassic Cynognathus Zone (Dreikoppen Formation, Beaufort Group) of the Karoo Basin, South Africa. *Palaios*, 16: 148-160.
- Gu, L., Dai, T.G., Deng, C.H., Fan, W.M., 2000. Petrochemical characteristics of the Mesozoic-Cenozoic volcanic rock in Huanghua basin. *J. Centr. South Univ. Techn.*, 7: 75-80.
- Gungor, T., Erdogan, B., 2001. Emplacement age and direction of the Lycian nappes in the Soke-Selcuk region, western Turkey. *International Journal of Earth Sciences*, 89: 874-882.
- Haas, J., Budai, T., Dosztály, L., Oraveczné Scheffer, A. & Tardiné, F. 2000. Upper Ladinian-Lower Carnian platform-slope facies in the Veszprém area, Transdanubian Range, Hungary. *Földtani Közlöny* 130 (4): 725-758.
- Haas, J. & Kovács, S. 2001. The Dinaridic-Alpine connection – as seen from Hungary. *Acta Geologica Hungarica*, 44 (2-3): 345-362.
- Hallam, A., 2001. A review of the broad pattern of Jurassic sea-level changes and their possible causes in the light of current knowledge. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 167: 23-37.
- Hannigan, P. K., Dietrich, J. R., Lee, P. J. & Osadetz, K. G. 2001. Petroleum resource potential of sedimentary basins on the Pacific margin of Canada. *Geological Survey of Canada Bulletin* 564, 72pp.
- Haubold, H., Klein, H., 2000. Die dinosauroiden Fährten *Parachirotherium* - *Atreipus* - *Grallator* aus dem unteren Mittelkeuper (Obere Trias: Ladin, Karn, ?Nor) in Franken. *Hallesches Jahrb. Geowiss.*, B 22: 59-85.
- Hauser, M., Martini, R., Burns, S., Dumitrica, P., Krystyn, L., Matter, A., Peters, T. & Zaninetti, L. 2001. Triassic stratigraphic evolution of the Arabian-Greater India embayment of the southern Tethys margin. *Eclogae Geologicae Helveticae*, 94 (1): 29-62.
- Hauschke, N., Wilde, V., 2000. Limulidenreste aus dem Unteren Butsandstein (Bernburg-Formation) von Beesenlaubingen (Sachsen-Anhalt). *Hallesches Jahrb. Geowiss.*, B 22: 87-90.
- Hautmann, M., 2001. Taxonomy and phylogeny of cementing Triassic bivalves (families Prospondylidae, Plicatulidae, Dimyidae and Ostreidae). *Palaeontology*, 44: 339-373.
- Henriksen, N., Higgins, A. K., Kalsbeek, F. & Pulvertaft, T. C. R. 2000. Greenland from Archaean to Quaternary. Descriptive text to the Geological Map of Greenland, 1:2 500 00. *Geology of Greenland Survey Bulletin* 185, 93pp.
- Hervé, F., Fanning, C.M., 2001. Late Triassic detrital zircons in meta-turbidites of the Chonos Metamorphic Complex, southern Chile. *Revista Geologica de Chile*, 28: 91-104.
- Heydari, E., Wade, W. J. & Hassanzadeh, J. 2001. Diagenetic origin of carbon and oxygen isotope

- compositions of Permian-Triassic boundary strata. *Sedimentary Geology*, 143, 191-197.
- Holmes, W.B.K., 2000. The Middle Triassic megafossil flora of the Basin Creek Formation, Nymboida Coal Measures, NSW, Australia. Part 1: Bryophyta, Sphenophyta. *Proceedings of the Linnean Society of New South Wales*, 122: 43-68.
- Letto, F. & Romano, R., 2001. Middle Triassic age and litho-stratigraphy of the Diplopora marbles from the San Donato Unit auctt. (Catena Costiera, Calabria, Italy). *Bollettino della Società Paleontologica Italiana*, 40 (1): 17-24.
- Irnstorfer, O. 2001. Trias-Ammoniten aus den Hallstätter Kalken (3. Teil). *Fossilien*, 2001(3): 144-150.
- Ishida, K. & Hirsch, F. 2001. Taxonomy and faunal affinity of Late Carnian-Rhaetian conodonts in the southern Chichibu Belt, Shikoku, SW Japan. *Rivista Italiana di Paleontologia e Stratigrafia*, 107 (2): 227-250.
- Jaacks, G.S., Carlson, S.J., 2001. How phylogenetic inference can shape our view of heterochrony: examples from thecideid brachiopods. *Paleobiology*, 27: 205-225.
- Jin Hui-Juan & Lu Yu-Ci. 2001. A study of Middle Triassic flysch facies in West Qinling orogenic belt. *Acta Sedimentologica Sinica*, 19 (3): 321-326.
- Kaiho, K., Kajiwara, Y., Nakano, T., Miura, Y., Kawahata, H., Tazaki, K., Ueshima, M., Chen, Z. & Shi, G. R. 2001. End-Permian catastrophe by bolide impact: evidence of a gigantic release of sulfur from the mantle. *Geology*, 29 (9): 815-818.
- Kamoun, F., Peynèrnès, B., Ciszak, R. & Calzada, S. 2001. Triassic palaeogeography of Tunisia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 172: 223-242.
- Kattenhorn, S. A. & Pollard, D. D. 2001. Integrating 3-D seismic data, field analogs, and mechanical models in the analysis of segmented normal faults in the Wytch Farm oil field, southern England, United Kingdom. *American Association of Petroleum Geologists Bulletin*, 85 (7): 1183-1210.
- Keim, L. & Schlager, W. 2001. Quantitative compositional analysis of a Triassic carbonate platform (southern Alps, Italy). *Sedimentary Geology*, 139: 261-283.
- Kemkin, I. V. & Filippov, A. N. 2001. Structure and genesis of the lower structural unit of the Samarka Jurassic accretionary prism (Sikhote-Alin, Russia). *Geodiversitas*, 23 (3): 323-339.
- Kerr, R.A., 2001. Whiff of gas points to impact mass extinction. *Science*, 291: 1469-1470.
- Kiessling, W. 2001. Paleoclimatic significance of Phanerozoic reefs. *Geology*, 29 (8): 751-754.
- Kidder, D.L., Erwin, D.H., 2001. Title: Secular distribution of biogenic silica through the phanerozoic: Comparison of silica-replaced fossils and bedded cherts at the series level. *Journal of Geology*, 109: 509-522.
- Kim, J. 2001. New fossil plants from the Nampo Group (Lower Mesozoic), Korea. *Geosciences Journal*, 5 (2): 173-180.
- Koralay, O. E., Satir, M. & Dora, O. Ö. 2001. Geochemical and geochronological evidence for Early Triassic calc-alkaline magmatism in the Menderes Massif, western Turkey. *International Journal of Earth Sciences*, 89: 822-835.
- Krause, T., Aktuelle Zahnfunde der Lungenfische *Ptychoceratodus* und *Ceratodus* aus der Thuringer Triasmulde. *Veröffentlichungen des Naturhistorischen Museum Schleusingen*, 15: 41-45.
- Krzeminski, W. & Lombardo, C. 2001. New fossil Ephemeroptera and Coleoptera from the Ladinian (Middle Triassic) of Canton Ticino (Switzerland). *Rivista Italiana di Paleontologia e Stratigrafia*, 107 (1): 69-78.
- Kurushin, N. I. & Truschelev. A. M., 2001. *Magnolobia*: A new bivalve genus from the Boreal Ladinian. *Paleontological Journal*, 35: 243-248.
- Lago San José, M, Galé Bornao, C, Arranz Yaguë, Vaquer Navarro, R., Gil Imaz, A., Povoci Juan, A., 2000. Triassic tholeiitic dolerites (\*ophites\*) of the El Grado diapir (Pyrenees, Huesca, Spain): emplacement and composition. *Estudios Geol.*, 65: 3-18.
- Lai, X.L., Wignall, P., Zhang, K.X., 2001. Palaeoecology of the conodonts *Hindeodus* and *Clarkina* during the Permian-Triassic transitional period. *Palaeogeography Palaeoclimatology Palaeoecology*, 171: 63-72.
- Landra, G., Longoni, R. & Fantoni, R., 2000. Seis-

- mic models of two middle Triassic carbonate platforms from the Southern Alps. *Mem. Sci. Geol.*, 52 (1): 139-163.
- Lehrman, D. J., Yang Wan, Jiayong Wei, You Yi Yu & Jiafei Xiao. 2001. Lower Triassic peritidal cyclic limestone: an example of anachronistic carbonate facies from the Great Bank of Guizhou, Nanpanjiang Basin, Guizhou province, South China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 173: 103-123.
- Le Roy, P., Pique, A., 2001. Triassic-Liassic western Moroccan synrift basins in relation to the central Atlantic opening. *Marine Geology*, 172: 359-381.
- Leven, E.J., Korchagin, O.A., 2001. Permian-Triassic biotic crisis and foraminifers. *Stratigraphy and Geological Correlation*, 9: 364-372.
- Li Wen-Ben. 2000. The fossil megaspore *Tricristatispora tricristata* (LI, 1974) comb. nov. *Acta Micropalaeontologica Sinica*, 17 (4):451-455.
- Liu Yi-Qun, Yuan Ming-Sheng, Li Hua-Ming & Wei Jian-Jing. 2001. Formation conditions of the oil-bearing sandbodies of Triassic and Jurassic in Turpan-Hami Basin, Xinjiang. *Acta Sedimentologica Sinica*, 19 (3): 399-404.
- Liu Zhao-Sheng. 2000. Palynological assemblage of the Sangonghe Formation from Tuziakeneigou, Xinjiang, NW China. *Acta Micropalaeontologica Sinica*, 17 (4):443-450.
- Liu Zhi-Li & Tong Jin-Nan. 2001. The Middle Triassic stratigraphy and sedimentary paleogeography of South China. *Acta Sedimentologica Sinica*, 19 (3): 327-332.
- Li, Z.X., Powell, C.M., 2001. An outline of the palaeogeographic evolution of the Australasian region since the beginning of the Neoproterozoic. *Earth Science Reviews*, 53: 237-277.
- Lindhorst, D., 2000. Biometrische Untersuchungen an der Gattung *Stolleyites* Arkhipov, 1974 (Nathorstitidae, Ceratitinae) aus der Oberen Trias (Unter-Karn) Spitzbergens. *Mitt. Geol.-Paläont. Inst. Univ. Hamburg*, 84: 1-64.
- Litvinovsky, B.A., Yarmolyuk, V.V., Vorontsov, A.A., Zhuravlev, D.Z., Posokhov, V.F., Sandimirova, G.P., Kuz'min, D.V., 2001. Late Triassic stage of formation of the Mongolo-Transbaikalian alkaline-granitoid province: Data of isotope-geochemical studies. *Geologiya i Geofizika* 42: 445-455.
- Liu, H.T., 2001. Qimantage terrestrial volcanics: petrologic evidence of active continental margin of Tarim Plate during late Indo-China epoch. *Acta Petrologica Sinica*, 17: 337-351.
- Lott, G. K. 2001. Geology and building stones in the East Midlands. *Mercian Geologist*, 15 (2): 97-122.
- Looy, C.V., Twitchett, R.J., Dilcher, D.L., Konijnenburg-Van Cittert, J.H.A., Visscher, H., 2001. Life in the end-Permian dead zone. *Proceedings of the National Academy of Sciences of the United States of America*, 98: 7879-7883.
- Lou, Z., Zhu, R., Tang, Z., Liu, Z., Zhao, X. & Chen, Z. 2000. Sedimentary environments of the Cangfanggou Group in Junggar basin, Xinjiang, in response to climate and tectonic regime. *Acta Geologica Sinica*, 74 (4): 795-806.
- Lucas, S. G., Heckert, A. B., Harris, J. D., Seegis D., & Wild, R.. 2001. Mammal-like tooth from the Upper Triassic of Germany: *Journal of Vertebrate Paleontology*, 21: 397-399.
- Lucas, S. G. & Heckert, A. B. 2001. Theropod dinosaurs and the Early Jurassic age of the Moenave Formation, Arizona-Utah, USA: *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*, 2001: 435-448.
- Lugli, S. 2001. Timing of post-depositional events in the Burano Formation of the Secchia valley (Upper Triassic, Northern Apennines), clues from gypsum-anhydrite transitions and carbonate metasomatism. *Sedimentary Geology*, 140: 107-122.
- Luo, Z., Compton, A. W. & Sun, A. 2001. A new mammaliaform from the Early Jurassic and evolution of mammalian characteristics. *Science*, 292: 1535-1540.
- Lutz, A., Crisafulli, A. & Herbst, R., 2001. Contribución al estudio xiloflorístico de la Formación La Ternera, Triásico Superior (Chile). *Amerghiniana*, 38: 119-127.
- MacIntyre, D. G., Villeneuve, M. E. & Schiarizza, P. 2001. Timing and tectonic setting of Stikine Terrane magmatism, Babine-Takla lakes area, central British Columbia. *Canadian Journal of Earth Sciences*, 38: 579-601.

- Maisch, M.W., 2001. Observations on Triassic ichthyosaurs. Part VII. New data on the osteology of *Chaohusaurus geishanensis* Young & Dong, 1972 from the Lower Triassic of Anhui (China). N. Jb. Geol. Paläont., Abh., 219: 305-327.
- Maisch, M. W. 2001. Observations on Karoo and Gondwana vertebrates. Part 2: a new skull-reconstruction of *Stahleckeria potens* von HUENE, 1935 (Dicynodontia, Middle Triassic) and a reconsideration of kannemeyeriiform phylogeny. Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 220 (1):127-152.
- Maisch, M. W. & Matzke, A. T. 2000. The mixosaurid ichthyosaur *Contectopalatus* from the Middle Triassic of the German Basin. Lethaia, 33 (2): 71-74.
- Maisch, M. W., & Matzke, A. T. 2001. Observations on Triassic ichthyosaurs. Part VIII. A redescription of *Phalarodon major* (von HUENE, 1916) and the composition and phylogeny of the Mixosauridae. Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 220 (3): 431-447.
- Maliszewskiej, A. (ed.) 1999. Diagenesis of Upper Permian and Mesozoic deposits of the Kujawy region (central Poland). Prace Panstwowego Instytutu Geologicznego, 167: 154pp.
- Mángano, M. G., Buatois, L. B., Wu Xiantao, Sun Junmin & Zhang Guocheng. 2000. Triassic lacustrine sedimentation from the Tanzhuang Formation, Jiyuan-Yima Basin, southeastern China. AAPG Studies in Geology, 46: 133-140.
- Manuszak, J. D., Satterfield, J. I. & Gehrels, G. E. 2000. Detrital zircon geochronology of Upper Triassic strata in western Nevada. Geological Society of America, Special Paper 347: 109-118.
- Marchal-Papier, F., Nel, A. & Grauvogel-Stamm, L., 2000. Nouveaux Orthoptères (Ensifera, Insecta) du Trias des Voges (France). Acta Geologica Hispanica, 35 (1-2): 5-18.
- Matura, A., Császár, G., Kröll, A., Vozár, J. & Wessely, G., 2000. Map of the pre-Tertiary basement. Danube region environmental geology programme DANREG – Explanatory notes (G. Császár, ed.). Jahrbuch der Geologischen Bundesanstalt, 142 (4): 465-482.
- McArthur, J.M., Howarth, R.J., Bailey, T.R., 2001. Strontium isotope stratigraphy; LOWESS Version 3; best fit to the marine Sr-isotope curve for 0-509 Ma and accompanying look-up table for deriving numerical age. Journal of Geology, 109: 155-170.
- McKinley, J. M., WORDEN, R. H. & RUFFELL, A. H. 2001. Contact diagenesis: the effect of an intrusion on reservoir quality in the Triassic Sherwood Sandstone Group, Northern Ireland. Journal of Sedimentary Research, 71 (3): 484-495.
- McLoughlin, S., 2001. The breakup history of Gondwana and its impact on pre-Cenozoic floristic provincialism. Australian Journal of Botany, 49: 271-300.
- McNeill, G. W., CRONIN, A. A., YANG, Y., ELLIOT, T. & KALIN, R. M. 2000. The Triassic Sandstone aquifer in Northern Ireland: constraint of a groundwater flow model for resource management. Pp.179-190 in Robins, N. S. & Misstear, B. D. R. (eds) Groundwater in the Celtic regions: studies in hard rock and Quaternary hydrogeology. Geological Society, London, Special Publication 182.
- McRoberts, C. A. 2001. Triassic bivalves and the initial marine Mesozoic revolution: a role for predators? Geology, 29 (4): 359-362.
- Melchor, R.N., Herbst, R. 2000. Sedimentology of the El Puquen Formation (Upper Triassic, central Chile) and the new plant *Mollesia melendeziae* gen. et sp nov (Pteridophylla, incertae sedis). Ameghiniana, 37: 477-485.
- Menning, M. 2001. A Permian time scale 2000 and correlation of marine and continental sequences using the Illawarra Reversal (265MA). Natura Bresciana, Monografia 25: 355-362.
- Mezzalira, S. 2000. Os Fósseis do Estado de São Paulo, Parte II. Período 1987 (parcial) – 1996. Boletim do Instituto Geológico, São Paulo, 15.
- Milani, E. J., Faccini, U. F., Scherer, C. M., Araújo, L. M. & Cupertino, J. A. 1998. Sequences and stratigraphic hierarchy of the Paraná Basin (Ordovician to Cretaceous), southern Brazil. Boletim Instituto de Geociências – Universidade de São Paulo 29: 125-173.
- Miller, M. F., Hasiotis, S. T., Babcock, L. E., Isbell, J. L. & Collinson, J. W. 2001. Tetrapod and large burrows of uncertain origin in Triassic high paleolatitude floodplain deposits, Antarctica. Palaios, 16: 218-232.
- Mojica, J. & Prinz-Grimm, P. 2000. La fauna de Amonitas del Triásico Tardío en el Miembro

- Chicalá (=parte baja de la Formación Saldaña) en Payandé, Tolima, Colombia. *Geologia Colombiana*, 25: 13-23.
- Mundil, R., Metcalfe, I., Ludwig, K. R., Renne, P. R., Oberli, F. & Nicoll, R. S. 2001. Timing of the Permian-Triassic biotic crisis: implications from new zircon U/Pb age data (and their limitations). *Earth and Planetary Science Letters*, 187: 131-145.
- Musashi, M., Isozaki, Y., Koike, T. & Kreulen, R. 2001. Stable carbon isotope signature in mid-Panthalassa shallow-water carbonates across the Permo-Triassic boundary: evidence for <sup>13</sup>C-depleted superocean. *Earth and Planetary Science Letters*, 191 (1-2): 9-20.
- Muttoni, G., Garzanti, E., Alfonsi, L., Cirilli, S., Germani, D. & Lowrie, W. 2001. Motion of Africa and Adria since the Permian: paleomagnetic and paleoclimatic constraints from northern Libya. *Earth and Planetary Science Letters*, 192: 159-174.
- Nedjari, A., Mesbah, M., Benhammouche, A. & Berraki, A. 2001. La province triasique saharienne: une approche de sa structure par l'analyse du réseau hydrographique sub actuel et actuel. *Bulletin du Service Géologique de l'Algérie*, 12 (1): 77-95.
- Nicholls, E. L. & Manabe, M. 2001. A new genus of ichthyosaur from the Late Triassic Pardonet Formation of British Columbia: bridging the Triassic-Jurassic gap. *Canadian Journal of Earth Sciences*, 38 (6): 983-1002.
- Nokleberg, W. J., Parfenov, L. M., Monger, J. W. H., Norton, I. O., Khanchuk, A. I., Stone, D. B., Scotese, C. R., Scholl, D. W. & Fujita, K. 2000. Phanerozoic tectonic evolution of the circum-North Pacific. U.S. Geological Survey Professional Paper 1626, vii+122pp.
- Novikov, I. V., Shishkin, M. A. & Golubev, V. K. 2000. Permian and Triassic anthracosaurs from Eastern Europe. Pp.60-70 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Novokshonov, V. G., 2001. New Triassic Scorpionflies (Insecta, Mecoptera) from Kyrgyzstan. *Paleontological Journal*, 35: 281-288.
- Ochev, V. G. & Surkov, M. V. 2000. The history of excavation of Permo-Triassic vertebrates from Eastern Europe. Pp.1-16 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Orchard, M. J., Cordey, F., Rui, L., Bamber, E. W., Mamet, B., Struik, L. C., Sano, H. & Taylor, H. J. 2001. Biostratigraphic and biogeographic constraints on the Carboniferous to Jurassic Cache Creek Terrane in central British Columbia. *Canadian Journal of Earth Sciences*, 38: 551-578.
- Ottone, E.G., Amenabar, C.R., 2001. A new disaccate pollen grain from the Triassic of Argentina. *Ameghiniana*,38: 157-161.
- Ouarhache, D., Charriere, A., Chalot-Prat, F., El-Wartiti, M., 2000. Late Triassic to infra-Liassic continental detrital sedimentation synchronous with an explosive volcanic event in the Atlas area [High Moulouya, Morocco]. *J. Afr. Earth Sci.*, 31: 555-570.
- Ouarhache, D., Charriere, A., Chalot-Prat, F., El-Wartiti, M., 2001. Late Triassic to infra-Liassic continental detrital sedimentation synchronous with an explosive volcanic event in the Atlas area [High Moulouya, Morocco] (vol 31, pg 560, 2001). *Journal of African Earth Sciences*, 32: 1-1.
- Panko, L.J., 2000. How to distinguish the basal Triassic cynodonts *Galesaurus* and *Thrinaxodon* from one another using rib morphology. *American Zoologist*, 40: 1162-1162
- Papier, F. & Nel, A. 2001. Les Subioblattidae (Blattodea, Insecta) du Trias d'Asie Centrale. *Paläontologische Zeitschrift*, 74 (4): 533-542.
- Parrish, J. T., Droser, M. L. & Bottjer, D. J. 2001. A Triassic upwelling zone: the Shublik Formation, Arctic Alaska, U.S.A. *Journal of Sedimentary Research*, 71 (2): 272-285.
- Peeva, N., Stoyanova, G. & Belivanova, V. 1999. Investigation of the organic matter in the Triassic rocks from the Golo Bardo Mountain, southwestern Bulgaria. *Review of the Bulgarian Geological Society*, 60 (1-3): 49-53.
- Peng Yuanqiao, Tong Jinnan & Gao Yongqun. 2001. Griesbachian cyclostratigraphy at Majiashan section, Anhui Province, China. *Journal of China University of Geosciences*, 12 (1): 22-26.

- Radwanski, A. & Summesberger, H. 2001. A new species of inarticulate brachiopods, *Discinisca zapfei* sp.n., from the Upper Triassic Zlambach Formation (Northern Calcareous Alps, Austria), and a discussion of other Triassic disciniscans. *Annalen des Naturhistorischen Museums in Wien*, 102A: 109-129.
- Rein, S., 2000. Zur Lebensweise von *Ceratites* und *Germanonutilus* im Muschelkalkmeer. *Veröffentlichungen Naturhistorisches Museum Schleusingen*, 15: 25-40.
- Renesto, S., Dalla Vecchia, F.M., 2000. The unusual dentition and feeding habits of the prolacertiform reptile *Langobardisaurus* (Late Triassic, northern Italy). *Journal of Vertebrate Paleontology*, 20: 622-627.
- Repin, S. Y., 2001. New Taxa of Late Triassic Bivalve Mollusks from Central Iran. *Paleontological Journal*, 35: 233-242.
- Rieppel, O. 2001. A new species of *Tanystropheus* (Reptilia: Protorosauria) from the Middle Triassic of Makhtesh Ramon, Israel. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (2): 271-287.
- Rieppel, O. & Dalla Vecchia, F. M. 2001. Marine reptiles from the Triassic of the Tre Venezie area, northeastern Italy. *Fieldiana, Geology new series*, No.44: iii+25pp.
- Rodland, D. L. & Bottjer, D. J. 2001. Biotic recovery from the end-Permian mass extinction: behaviour of the inarticulate brachiopod *Lingula* as a disaster taxon. *Palaios*, 16: 95-101.
- Samankassou, E., Bernecker, M., Flügel, E., 2001. Facies and small-scale geometrics of the Late Triassic Kawr Platform, Sultanate of Oman. *Géologie Méditerranéenne*, 28: 155-157.
- Sari, A., Sonel, N. & Albayrak, M. 2001. Geochemistry and thermal maturation of Triassic to Eocene source rocks in the Central Taurus Belt, SW Turkey. *Petroleum Geoscience*, 7: 191-199.
- Sashida, K., Nakornsri, N., Ueno, K. & Sarsud, A. 2000. Carboniferous and Triassic radiolarian faunas from the Saba Yoi area, southernmost part of Peninsular Thailand and their palaeogeographic significance. *Science Reports of the Institute of Geoscience, University of Tsukuba, Section B – Geological Sciences* 21: 71-99.
- Scalera, G., 2001. The global paleogeographical reconstruction of the Triassic in the Earth's dilatation framework and the paleoposition of India. *Annali di Geofisica*, 44: 13-32.
- Schatz, W., 2001. Taxonomic significance of biometric characters and the consequences for classification and biostratigraphy, exemplified through moussoneliform daonellas (*Daonella*, Bivalvia; Triassic). *Palaeontologische Zeitschrift*, 75 (1): 51-70.
- Schlirf, M., Uchman, A. & Kuemmel, M., 2001. Upper Triassic (Keuper) non-marine trace fossils from the Hassberge area (Franconia, SE Germany). *Palaeontologische Zeitschrift*, 75 (1): 71-96.
- Schneider, J.A., Carter, J.G., 2001. Evolution and phylogenetic significance of cardioidean shell microstructure (Mollusca, Bivalvia). *Journal of Paleontology*, 75: 607-643.
- Sciunnach, D. 2001. Heavy mineral provinces as a tool for palaeogeographic reconstruction: a case study from the Buntsandstein of Nurra (W Sardinia, Italy). *Eclogae Geologicae Helvetiae*, 94: 197-211.
- Seeling, M., 2000. Die Sunundatabank (Oberer Buntsandstein) bei Spielberg (Sachsen-Anhalt) unter Berücksichtigung eines Spurenfossils. *Hallesches Jahrbuch für Geowissenschaften*, B 22: 91-98.
- Sennikov, A. G., 2001. Discovery of a primitive Sauropterygian from the Lower Triassic of the Donskaya Luka (Don Basin) and the Range of Triassic Marine Reptiles in Russia. *Paleontological Journal*, 35: 301-309.
- Senowbari-Daryan, B. & Di Stefano, P. 2001. Middle Triassic dasycladales in Sicily: evidence of an Anisian?-Ladinian carbonate platform. *Acta Geologica Hungarica*, 44 (1): 95-109.
- Senowbari-Daryan, B., Stanley, G. D. & Gonzalez-Leon, C. 2001. A new Triassic sponge from the Antimonio terrane, Sonora, Mexico. *Journal of South American Earth Sciences*, 14: 447-452.
- Senowbari-Daryan, B., Flügel, E. & Preto, N., 2001. *Tethysocarnia cautica* n. gen. N. sp., a sessil foraminifer from Late Ladinian and Carnian reefs of the western and southern Thetys. *Geologica et Palaeontologica*, 35: 105-119.
- Senkowiczowa, H. 2000. The Roethian and

- Muschelkalk stratigraphy of southwestern Poland. *Biuletyn Państwowego Instytutu Geologicznego*, 390: 97-1666.
- Sharland, P. R., Archer, R., Casey, D. M., Davies, R. B., Hall, S. H., Heward, A. P., Horbury, A. D. & Simmons, M. D. 2001. Arabian plate sequence stratigraphy. *GeoArabia Special Publication 2*: 371pp.
- Shilong Mei & Henderson, C. M. 2001. Evolution of Permian conodont provincialism and its significance in global correlation and palaeoclimate implication. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170: 237-260.
- Shishkin, M. A. 2000. Mesozoic amphibians of Middle and Central Asia. Pp.297-308 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Shishkin, M. A., Novikov, I. V. & Gubin, Y. M. 2000. Permian and Triassic temnospondyls from Russia. Pp.35-59 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Shishkin, M. A., Ochev, V. G., Lozpvskii, V. R. & Novikov, I. V. 2000. Tetrapod biostratigraphy of the Triassic of Eastern Europe. Pp.120-139 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Siblik, M., 2001. On the find of Triassic brachiopod fauna in the variegated micrites on the Dachstein Plateau. *Slovak Geol. Mag.*, 7: 91-94.
- Sivell, W. J. & McCulloch, M. T. 2001. Geochemical and Nd-isotopic systematics of the Permo-Triassic Gympie Group, southeast Queensland. *Australian Journal of Earth Sciences*, 48: 377-393.
- Sneh, A., Bartov, Y. & Weissbrod, T. 2000. Stratigraphic chart of exposed rock units in Israel. *Geological Survey of Israel, Current Research* 12: 257-262.
- Soreghan, M. J. & Gehrels, G. E. (eds). 2000. Paleozoic and Triassic paleogeography and tectonics of western Nevada and northern California. *Geological Society of America, Special Paper* 347.
- Spencer, P. S. & Benton, M. J. 2000. Procolophonids from the Permo-Triassic of Russia. Pp.160-176 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Spurlin, M. S., Gehrels, G. E. & Harwood, D. S. 2000. Detrital zircon geochronology of upper Paleozoic and lower Mesozoic strata of the northern Sierra terrane, northeastern California. *Geological Society of America, Special Paper* 347: 89-98.
- Stiller, F. 2001. Echinoid spines from the Anisian (Middle Triassic) of Qingyan, south-western China. *Palaeontology*, 44 (3): 529-551.
- Stiller, F. 2001. Scaphopods from the Upper Anisian (Middle Triassic) of Qingyan, SW China. *Neues Jahrbuch fuer Geologie und Palaeontologie Monatshefte*, 10: 617-632.
- Storrs, G. W., Arkhangel'skii, M. S. & Efimov, V. M. 2000. Mesozoic marine reptiles of Russia and other former Soviet republics. Pp.187-210 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Struik, L. C., Schiarizza, P., Orchard, M. J., Cordey, F., Sano, H., Macintyre, D. G., Lapierre, H. & Tardy, M. 2001. Imbricate architecture of the upper Paleozoic to Jurassic Cache Creek Terrane, central British Columbia. *Canadian Journal of Earth Sciences*, 38: 495-514.
- Su, P., Fan, X.Z., Shi, R.P., 2001. Magnetostratigraphy of Lower Triassic in Shuiyuguan, Jiaocheng, Shanxi. *Chin. J. Geophys. Ch.*, 44: 219-227.
- Svanidze, Ts. I., Lebanidze, Z. A. & Jakobidze, E. B. 2000. Upper Triassic-Bajocian vegetation of the western Georgia and stratigraphical significance of the flora. *Geologica Balcanica*, 30 (1-2): 19-24.
- Tabyaoui, M., Ait Brahim, L., Tahiri, A., Chotin, P., 1999. Évolution tectono-sédimentaire du Nord-Est du Maroc au cours du Trias supérieur: apport de données microtectoniques. *Géologie Méditerranéenne*, 26: 231-243.
- Tanner, L. H. 2000. Triassic-Jurassic lacustrine deposition in the Fundy Rift Basin, eastern

- Canada. AAPG Studies in Geology, 46: 159-166.
- Tardy, M., Lapierre, H., Struik, L. C., Bosch, D. & Brunet, P. 2001. The influence of mantle plume in the genesis of the Cache Creek oceanic igneous rocks: implications for the geodynamic evolution of the inner accreted terranes of the Canadian Cordillera. *Canadian Journal of Earth Sciences*, 38: 515-534.
- Taylor, R. S., Schram, F. R. & Shen Yan-Bin. 2001. A new upper Middle Triassic shrimp (Crustacea: Lophogastrida) from Guizhou, China, with discussion regarding other fossil "mysidaceans". *Journal of Paleontology*, 75 (2): 310-318.
- Tekin, U. K., 1999. Biostratigraphy and systematics of late Middle to Late Triassic Radiolarians from the Taurus Mountains and Ankara Region, Turkey. *Geologische Palaeontologische Mitteilungen Innsbruck Sonderband*, 5: 296 pp.
- Tiwari, R.S. 1999. The Permian-Triassic boundary, mass extinction and pattern of pteridophytic occurrence across the boundary. *Indian Fern Journal*, 16: 75-86.
- Tiwari, R.S., 2000. Permian Triassic palynofloras from Eaurameican and Gondwanan regions: an overview. In: Mukerji, Chamola & Chandra (eds.), *Glimpses in Botany*, pp. 100-120. APH Publishing Co., New Delhi.
- Torsvik, T.H., Van der Voo, R., Meert, J.G., Mosar, J., Walderhaug, H.J., 2001. Reconstructions of the continents around the North Atlantic at about the 60th parallel. *Earth and Planetary Science Letters*, 187: 55-69.
- Troncoso, A. & Herbst, R. 1999. Ginkgoales del Triásico del norte de Chile. *Revista Geológica de Chile*, 26 (2): 255-273.
- Twitchett, R. J. 2001. Incompleteness of the Permian-Triassic fossil record: a consequence of productivity decline? *Geological Journal*, 36: 341-353.
- Twitchett, R.J., Looy, C.V., Morante, R., Visscher, H., Wignall, P.B., 2001. Rapid and synchronous collapse of marine and terrestrial ecosystems during the end-Permian biotic crisis. *Geology*, 29: 351-354.
- Unwin, D. M., Alifanov, V. R. & Benton, M. J. 2000. Enigmatic small reptiles from the Middle-Late Triassic of Kirgizstan. Pp.177-186 in Benton, M. J., Shishkin, M. A., Unwin, D. M. & Kurochkin, E. N. (eds), *The age of dinosaurs in Russia and Mongolia*. Cambridge, Cambridge University Press.
- Urlichs, M. 2000. *Germanonautilus* (Nautiloidea) aus dem Unterkarnium der Dolomiten (Obertrias, Italien). *Stuttgarter Beiträge zur Naturkunde, Ser.B, Nr.291*: 13pp.
- Urlichs, M., Tichy, G., 2000. Zur Alterstellung der Schlernplateau-Schichten (Mitteltrias) am Schlern (westliche Dolomiten, Italien). *Stuttgarter Beiträge zur Naturkunde, Serie B*, 297: 1-10.
- Vachard, D., Martini, R. & Zaninetti, L. 2001. Earliest Artinskian (Early Permian) fusulinids reworked in the Triassic Lecara Formation (NW Sicily). *Journal of Foraminiferal Research*, 31 (1): 33-47.
- Van Der Voo, R. & Torsvik, T. H. 2001. Evidence for late Paleozoic and Mesozoic non-dipole fields provides an explanation for the Pangea reconstruction problems. *Earth and Planetary Science Letters*, 187: 71-81.
- Villeneuve, M., Cornée, J.-J., Gunawan, W., Martini, R., Tronchetti, M.-C., Janin, M.C., Saint-Marc, P., Zaninetti, L., 2001. La succession lithostratigraphique du bloc de Banda dans la région de Kolonodale (Sulawesi central, Indonésie). *Bulletin de la Société Géologique de France*, 172: 59-68.
- Vörös, A. 2001. Paleobiological analysis: a tool for the reconstruction of Mesozoic Tethyan and Penninic basins. *Acta Geologica Hungarica*, 44 (2-3): 145-158.
- Waltham, A. 2001. Landmark of geology in the East Midlands: the explosion crater at Fauld. *Mercian Geologist*, 15 (2): 123-125.
- Wang Darui, Wang Xinping & Sun Xiaoting. 2001. The anomaly of carbon and oxygen isotopic composition across the Permian-Triassic boundary in northwest Guangxi, South China, and its significance. *Geological Review*, 47 (3): 225-228.
- Wang, E., Meng, Q., Chen, Z. & Chen, L. 2001. Early Mesozoic left-lateral movement along the Longmen Shan fault belt and its tectonic implications. *Earth Science Frontiers*, 8 (1): 375-384.
- Wang Shangyan & Yin Hongfu. 2001. Discovery of microspherules in claystone near the terrestrial Permian-Triassic boundary. *Geological Review*, 47 (4): 411-414



- Ward, P. D., Haggart, J. W., Carter, E. S., Wilbur, D., Tipper, H. W. & Evans, T. 2001. Sudden productivity collapse associated with the Triassic-Jurassic boundary mass extinction. *Science*, 292:1148-1151.
- Wignall, P.B., 2001. Large igneous provinces and mass extinctions. *Earth-Science Reviews*, 53: 1-33.
- Winchester, S. 2001. The Map that changed the World. The tale of William Smith and the birth of a Science. London: Viking, xiv+338pp.
- Worsley, D. & Mørk, A. 2001. The environmental significance of the trace fossil *Rhizocorallium jenense* in the Lower Triassic of western Spitsbergen. *Polar Research*, 20 (1): 37-48.
- Wu, X.C., Russell, A.P., 2001. Redescription of *Turfanosuchus dabanensis* (Archosauriformes) and new information on its phylogenetic relationships. *Journal of Vertebrate Paleontology*, 21: 40-50.
- Wyld, S. J. 2000. Triassic evolution of the arc and backarc of northwestern Nevada and evidence of extensional tectonism. *Geological Society of America, Special Paper* 347: 185-207.
- Xulong Lai, Wignall, P. & Zhang, K. 2001. Palaeoecology of the conodonts *Hindeodus* and *Clarkina* during the Permian-Triassic transitional period. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170: 63-72.
- Yang Qun, Matsuoka, A., Wang Yu-Jing, Kobayashi, K., Nagahashi, T. & Zeng Qing-Gao. 2000. A Middle Triassic radiolarian assemblage from Quxia, Lhaze County, southern Tibet. *Science Reports of Niigata University, Series E (Geology)* 15: 59-65.
- Yang Shi-Pu, Deng Xi-Guang, Zheng Xiang-Shen & Liu Xiao-Han. 2001. Trace fossils in flysch facies in the Miers Bluff Formation (Late Triassic), Livingston Island, Antarctica. *Acta Palaeontologica Sinica*, 40 (1): 111-115.
- Yang Shou-Ren, Hao Wei-Cheng & Jiang Da-Yong. 2001. Early Triassic conodonts from Luolou Formation in Luolou, Lingyin, Guangxi. *Acta Palaeontologica Sinica*, 40 (1): 91-92.
- Yang Zhenyu & Besse, J. 2000. Early Triassic paleomagnetic results from southern part of Sichuan Basin and its tectonic implications. *Scientia Geologica Sinica* 35 (1): 77-82.
- Yao, A. & Kuwahara, K. 2000. Permian and Triassic radiolarians from the southern Guizhou Province, China. *Journal of Geosciences, Osaka City University* 43: 1-19.
- Yates, A.M., 2000. A new tiny rhytidosteid (Temnospondyli : Stereospondyli) from the Early Triassic of Australia and the possibility of hidden temnospondyl diversity. *Journal of Vertebrate Paleontology*, 20: 484-489.
- Yelles-Chaouche, A. K., Ait-Ouali, R., Bracene, R., Derder, M. E. M. & Djellit, H. 2001. Chronologie de l'ouverture du bassin des Ksour (Atlas Saharien, Algérie) au début du Mésozoïque. *Bulletin de la Société géologique de France*, 172 (3): 285-293.
- Yin Hongfu, Zhang Kexin, Tong Jinnan, Yang Zunyi & Wu Shunbao. 2001. The Global Stratotype Section and Point (GSSP) of the Permian-Triassic Boundary. *Episodes*, 24 (2):102-114.
- Yümün, Z. U., Kiliç, A. M., Önder, F. & Yılmaz, A. 2001. Stratigraphic characteristics of the Mesozoic carbonates around the Deliktas (Kangol-Sivas) region (Abstract). 54<sup>th</sup> Geological Congress of Turkey, Ankara, May 7-10, 2001. Abstracts and Proceedings: 1-2.
- Zagorchev, I., Chatalov, A., Trifonova, E., Goranov, E. & Pemov, P. 1999. New data about Triassic stratigraphy of the Skakavitsa anticline (Kyustendil Krayshte, southwest Bulgaria). *Review of the Bulgarian Geological Society*, 60 (1-3): 73-81.
- Zagorchev, I., Trifonova, E., Budurov, K., Goranov, E. & Pemov, P. 1999. The Triassic of southwest Bulgaria. New data on the Upper Triassic in the Konyava Mountain. *Geologica Balcanica* 29 (3-4): 25-32.
- Zeeh, S. & Geng, A. 2001. Entstehung und diagenese von dolomit in der Muschelkalk-Gruppe (Mitteltrias) SW-Deutschlands. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (3): 359-395.
- Zhang Lujin. 2000. Upper Triassic palynoflora from northeast Guizhou, China. *Palynofloras and palynomorphs of China*, 41-52. Hefei Press of University of Science and Technology of China.
- Zhang Zhenfang, Feng Qinglai, Fang Nianqiao, Huang Junyi, Li Shixiong & Chen Xuefang. 2001.

Geochemical characteristics and sedimentary environment of Triassic siliceous rocks from Muyinhe Formation in Changning-Menglian Belt, southwest Yunnan. *Earth Science – Journal of China University of Earth Sciences*, 26 (5): 449-455.

Zharkov, M.A., Chumakov, N.M., 2001. Humid and arid state of the warm biosphere. *Geologiya i Geofizika*, 42: 668-677.

Zharkov, M.A., Chumakov, N.M., 2001. Paleogeography and sedimentation settings during Permian-Triassic reorganizations in biosphere. *Stratigraphy and Geological Correlation*, 9: 340-363.

Zhenyu Yang & Besse, J. 2001. New Mesozoic apparent wander path for South China: tectonic consequences. *Journal of Geophysical Research*, 106 (B5): 8493-8520.

Ziegler, M. A. 2001. Late Permian to Holocene paleofacies evolution of the Arabia Plate and its hydrocarbon occurrences. *GeoArabia*, 6 (3): 445-503.

**Acknowledgements:**

The continuous help of Dr. Zwier Smeenk (Utrecht) and Dipl. Geol. Sunia Lausberg and Anke Grewing (Muenster) in tracing relevant Triassic literature is gratefully acknowledged. Thanks are also due to all authors who sent information on their recent publications.

G. Warrington publishes with the approval of the Director, British Geological Survey (N.E.R.C.).

## British Triassic Palaeontology: Supplement 26

**G. Warrington**

Since the completion of the writer's previous supplement (No.25; ALBERTIANA, 25: 48) on British Triassic palaeontology, the following works relating to aspects of that subject have been published or have come to his notice:

Ambrose, K. 2001. The lithostratigraphy of the Blue Lias Formation (Late Rhaetian-Early Sinemurian) in the southern part of the English Midlands. *Proceedings of the Geologists' Association*, 112: 97-110.

Carpenter, S. 2001. Excavating a vertebrate-bearing horizon in the Upper Triassic sediments of north Mendip. *Geology Today*, 17 (1): 29-32.

Clark, N. D. L. 2001. 18. Using the fossil resource: a Scottish museum perspective. In: Bassett, M. G., King, A. H., Larwood, J. G., Parkinson, N. A. & Deisler, V. K. (eds). *A Future for Fossils*. 84-88. National Museum of Wales, Geological Series No.19, Cardiff.

Clarkson, E. N. K. 2001. 1. The palaeontological resource of Great Britain – our fossil heritage. In: Bassett, M. G., King, A. H., Larwood, J. G., Parkinson, N. A. & Deisler, V. K. (eds). *A Future for Fossils*. 10-17. National Museum of Wales, Geological Series No.19, Cardiff.

Duffin, C. J. 2001. Synopsis of the selachian genus *Lissodus* BROUGH, 1935. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, 221 (2): 145-218.

Godefroit, P. & Sigogneau-Russell, D. 1999. Kuehneotheriids from Saint-Nicholas-de-Port (Late Triassic of France). *Geologica Belgica*, 2 (3-4): 181-196.

Heckert, A. B. & Lucas, S. G. 2000. Taxonomy, phylogeny, biostratigraphy, biochronology, paleobiogeography, and evolution of the Late Triassic Aetosauria (Archosauria: Crurotarsi). *Zentralblatt für Geologie und Paläontologie, Teil I, Jahrgang 1998, Heft 11-12*: 1539-1587.

King, A. H. & Larwood, J. G. 2001. 3. Conserving our most 'fragile' fossil sites in England: the use of 'OLD25'. In: Bassett, M. G., King, A. H.,

Larwood, J. G., Parkinson, N. A. & Deisler, V. K. (eds). *A Future for Fossils*. 24-31. National Museum of Wales, Geological Series No.19, Cardiff.

Langer, M., Ferigolo, J. & Schultz, C. L. 2000. Heterochrony and tooth evolution in hyperodontine rhynchosaurs (Reptilia, Diapsida). *Lethaia*, 33 (2): 119-128.

Larwood, J. G. & King, A. H. 2001. 25. Conserving palaeontological sites: applying the principles of sustainable development. In: Bassett, M. G., King, A. H., Larwood, J. G., Parkinson, N. A. & Deisler, V. K. (eds). *A Future for Fossils*. 119-125. National Museum of Wales, Geological Series No.19, Cardiff.

Maisch, M. W. & Matzke, A. T. 2000. The Ichthyosauria. *Stuttgarter Beiträge zur Naturkunde, Serie B, Nr.298*: 159pp.

Thomas, B. A. & Batten, D. J. 2001. The Triassic palaeobotany of Great Britain. In: *Mesozoic and Tertiary Palaeobotany of Great Britain*, Geological Conservation Review Series, No.22, (C. J. Cleal, B. A. Thomas, D. J. Batten & M. E. Collinson), Joint Nature Conservation Committee, Peterborough, pp.17-27.

Thompson, D. B. 2001. An enigmatic borehole core with Triassic trace fossils displayed in the entrance foyer of the Earth Sciences Department of Keele University. *Bulletin of the North Staffordshire Group of the Geologists' Association*, 58, Appendix 14: 1-9.

Tresise, G. & Radley, J. D. 2000. Triassic footprints: the first English finds. *Geological Curator*, 7 (4): 135-140.

Webber, M. 2001. 23. The sustainability of a threatened fossil resource: Lower Jurassic *Caloceras* beds of Doniford Bay, Somerset. In: Bassett, M. G., King, A. H., Larwood, J. G., Parkinson, N. A. & Deisler, V. K. (eds). *A Future for Fossils*. 108-113. National Museum of Wales, Geological Series No.19, Cardiff.

This contribution is published with the approval of the Director, British Geological Survey (N.E.R.C.).

### Contributor's address:

G. Warrington, British Geological Survey, Keyworth, Nottingham NG12 5GG, Great Britain.

## New Books

### THE RECONSTRUCTION OF LATE PALEOZOIC AND MESOZOIC MARINE ENVIRONMENTS FROM ISOTOPIC DATA (evidence from north Eurasia)

Y.D. ZAKHAROV, N.G. BORISKINA, A.M.  
POPOV

Vladivostok: Dalnauka, 2001. 112 p.  
(in Russian, with English abstract and  
explanation of figures and tables)

<http://www.fegi.ru/FEGL/reconst/>

Some recently discovered events at the Tethys, respectively occurring at (1) early Permian (Artinskian?), (2) Wordian, (3) early Midian, (4) Midian-Dzhulfian transition, (5) early Dorashamian, (6) middle Olenekian, (7) early Anisian, (8) latest Ladinian, (9) late Carnian, (10) early Norian and (11) early Rhaetian intervals are characterized by anomalously high values of  $\delta^{13}\text{C}$  in organogenic carbonates («carbon-isotope macrorhythms»). We can consider the four brightest Phanerozoic events, which were probably reflected by the greatest phytoplankton heyday and, accordingly, by an intensive photosynthesis: Late Carboniferous ( $\delta^{13}\text{C}=6.2\text{‰}$ ), Midian-Dzhulfian  $6.5\text{-}7.3\text{‰}$ ), middle Olenekian ( $6.9\text{‰}$ ) and early Aptian ( $6.8\text{‰}$ ).

The geochemical variations appear explainable by high bioproductivity of data on isotopic composition of organic carbonates testify that the carbon-isotopic anomalies in many periods of the Phanerozoic appreciably are reflection of climatic fluctuation. Their positive maxima during the end of Paleozoic and Mesozoic fell, probably, on warm epochs caused by several factors, main of which seem to be: (1) rise of solar activity, (2) macropulsation of an energy of the sun and (3) number of astronomical variations. About eight solar cycles distinguishing on their duration are known now. Carbon-isotopic macrorhythms of Late Palaeozoic and Mesozoic demonstrated here may testify to the existence of cycles of solar activity (solar macrocycles) by duration from 1.5 up to 12 Ma, less often up to 15-18 Ma.

Based on oxygen-isotope and Ca-Mg analysis of invertebrate shells and limestones in the Tethys

the highest Late Paleozoic temperatures in North Hemisphere were obtained for the Midian-Dzhulfian transition (till  $27.9^{\circ}\text{C}$ ). In general, at low latitudes temperatures are thought to begin a recurrent warming trend in the Artinskian-Dorashamian, reaching temperature maxima in the Artinskian of Early Permian, early Murgabian and particularly Midian-Dzhulfian transition time of Middle Permian, and in the late early Dorashamian ( $23.8^{\circ}\text{C}$ ) – early late Dorashamian ( $23.8^{\circ}\text{-}24.2^{\circ}\text{C}$ ) time of the Late Permian, which agrees with a placing of the positive carbon isotope anomalies and taxonomic diversity of Permian invertebrates.

It seems justified to assume that through the course of the early Induan temperature dropped of about three times (twice within the beginning of the Lytophicerias medium Zone and once just at the beginning of the Gyronites Zone, with a visible warming in the middle part of the latter. A short-term fall of temperature at the beginning of the Induan (volcanic winter) soon followed by warm period (greenhouse summer) seem to be caused by eruption of the Siberian Traps. A similar situation apparently occurred at the end of the Cretaceous, when extensive flood basalts, known as Decan Traps, erupted.

Temperatures during early Olenekian seem to be significantly lower than during late Olenekian or late Anisian ( $8.8^{\circ}\text{C}$ ,  $16.2^{\circ}\text{C}$  and  $15.4^{\circ}\text{C}$ , respectively for Arctic Siberia).

Two main trends can be recognized in Late Cretaceous temperature conditions in the shallow-water Hokkaido-south Sakhalin Basin. (1) In general, a recurrent warming trend is thought to have begun in the Turonian-Campanian, reaching temperature maxima in the early Late Santonian ( $19.6^{\circ}\text{C}$ ) and perhaps early Campanian. (2) During the Maastrichtian, temperatures dropped sharply (till  $7.1^{\circ}\text{C}$ ), with only a slight warming (till  $11.2^{\circ}\text{C}$ ) in early Late Maastrichtian. The existence of a Santonian-transition thermal maximum has previously been expected, but is not confirmed by new isotopic results. The influence of some basic factors: drop of temperature, oxygen deficit, enormous sea level fluctuation and volcanic activity seems to be the main reason for the destruction of epicontinental sea ecosystems both at the end of the Permian and the end of the Cretaceous.

For reconstruction of late Palaeozoic and early Mesozoic environments the data from reef distribution seem to be very important, because the reefs consider to be very sensible indicator for marine environment changes. As an example of a prospering reef is that of the end-Permian strata

of the Urushtenskaya Suite in North Caucasus. It is known that at the start of the Triassic, reefs disappeared from the face of the earth and a reef formation was not renewed in any region of the world in both the middle Olenekian climatic optimum (transgression) and the similar condition of the beginning of Middle Triassic. After the Permian-Triassic boundary ecological crisis they have arisen in the tropical zone only in Late Triassic (although scleractinian corals made their first appearance in Middle Triassic). Lack of reefs in the low latitudes during the beginning of the Triassic more logically to connect with O<sub>2</sub> deficit of that time as consequence of the anoxic event across the Permian-Triassic boundary. The absence of visible signs of organic SiO<sub>2</sub>-accumulation just in the Permian-Triassic transition time and the low rates in reconstruction of the radiolarian taxonomic diversity through the Induan and Olenekian to the early Anisian seem to be caused by the same reason.

## Future Meetings

### Shallow Tethys 6, Budapest, Hungary, 26-31st August, 2002.

See <http://pangea.elte.hu/paleo/tethys/>

This is the venue for an official STS meeting with a thematic session on "Middle Triassic boundaries". It is hoped that substantial progress towards agreement on the GSSPs for the O-A, A-L, and L-C boundaries will result from this meeting. Also of interest will be a meeting of IGCP project 458 on Triassic-Jurassic boundary events. A full program of lectures and field trips will accommodate those interested in other Triassic topics.

The STS will also be independently arranging a pre-conference field trip from Italy to Hungary to view GSSP candidates at Bagolino (A-L, Italy), Prati di Stuares (C-N, Italy), and Felsoors (A-L, Hungary). The option of an additional post-meeting field trip to Delsi Caira (O-A, Romania) is possible depending on demand. Space for all these field trips will be limited and priority will be given to voting and working group members. Those interested in these trips should inform the Chair asap.

### Paleontological & Biostratigraphic Congress, Corrientes Province, Argentina, Oct., 2002.

A Triassic Symposium is being organized at this congress by South American colleagues. Contact Andrea Arcucci at email [arcucci@satlink.com](mailto:arcucci@satlink.com)

### Geological Ass. Canada, Vancouver, British Columbia, Canada, 26-28th May, 2003.

This will be a joint meeting of the Geological Association of Canada (GAC), Mineralogical Association of Canada (MAC) and the Society of Economic Geologists (SEG). A thematic session on Upper Triassic standards and correlations will be organized. A field trip to Williston Lake, British Columbia will include a visit to the Carnian-Norian boundary GSSP prospect and several Triassic-Jurassic contacts. In addition, the non-marine auxiliary C-N candidate in the Chinle

Group of New Mexico, USA, will be organized by Spencer Lucas..

### 32<sup>nd</sup> International Geological Congress, Florence, Italy, August 20-28th, 2004.

"From the Mediterranean toward a Global Renaissance" is the banner of this IGC. We plan an STS workshop and at least one Symposium topic has been proposed: "Triassic of the Tethys Realm" (M. Gaetani).

## BOUNDARY WORKING GROUPS

In order to make quicker progress toward GSSP definition in the Triassic, the Chair, on behalf of the executive, has now created additional boundary working groups (WGs) and nominated some new Chairpersons. In line with ICS recommendation, these WGs are to be more formally constituted and the onus will be upon the Chairs (plus in some cases a secretary) to convene a representative group of experts to nominate and consider GSSP candidates. Their recommendation will form the basis of any subsequent vote undertaken by the STS voting slate so much of the essential scientific work falls upon the members of the WGs.

Whereas Chairs may invite researchers who they know to be active in particular boundary studies, those STS members who feel they have a contribution to make on specific boundaries may write directly to the contact people below and make their interests and views known. It is important that we do not encumber the working groups with numerous peripherally interested persons so please consider carefully your potential contribution. For your information, the recently revised guidelines for GSSP proposals are reproduced below.

P-T*	-	Yin Hongfu
I-O	-	Yuri Zacharov
O-A**	-	Eugen Gradinaru
A-L	-	Aymon Baud
L-C	-	Maurizio Gaetani
C-N	-	Mike Orchard
N-R	-	Leo Krystyn
Non-marine	-	Spencer Lucas

\*The P-T boundary GSSP is now fixed at Meishan, China. \*\*There is currently only one candidate for the O-A boundary GSSP (Desli

Caira, Romania) for which the responsible proponent is E. Gradinaru.

## **Global Boundary Stratotype Section and Point (GSSP)**

### **Guidelines for GSSP proposals**

The Global Boundary Stratotype Section and Point is a permanent reference for geological time. The GSSP is a specific point in a stratotype, clearly marked and documented.

1. Stratigraphic rank of boundary
2. Proposed GSSP — geographic & physical geology
  - A. Geographic location
  - B. Geological location
  - C. Location of Level and Specific Point
  - D. Stratigraphic completeness
  - E. Adequate thickness and stratigraphic extent
  - F. Provisions for conservation and protection
3. Primary and Secondary Markers
  - A. Principal correlation event (marker) at GSSP level
  - B. Other stratigraphy — Biostratigraphy, Magnetostratigraphy, Chemical stratigraphy, Sequence stratigraphy, Cycle stratigraphy, Other event stratigraphy, Marine-Land correlation potential, Amenability to geochronometry
  - C. Demonstration of regional and global correlation.
4. Selection process
  - A. Relation of the GSSP to historical usage
  - B. Other candidates and reasons for rejection
  - C. Votes
  - D. Selected publications
5. Other useful sections

## STS MEMBERS LIST 2001

### Voting members 2001

**Yoshiaki Aita**, Utsunomiya, JAPAN

[aida@cc.utsunomiya-u.ac.jp](mailto:aida@cc.utsunomiya-u.ac.jp)

*Radiolarians in the southern high latitudes, mainly from New Zealand and Russia; P-T boundary succession in New Zealand.*

**Andrea B. Arcucci**, San Luis, ARGENTINA

[arcucci@satlink.com](mailto:arcucci@satlink.com)

*Vertebrate paleontology of the continental Triassic of western Argentina (faunal evolution, taphonomic, stratigraphic, and sedimentological studies).*

**Darioush Baghbani**, Tehran, IRAN

[baghbanid@nioc-ripi.org](mailto:baghbanid@nioc-ripi.org)

*Biostratigraphy and chronostratigraphy of Triassic Foraminifera, especially the Zagros Basin, SW Iran.*

**Aymon Baud**, Lausanne, SWITZERLAND

[Aymon.Baud@sst.unil.ch](mailto:Aymon.Baud@sst.unil.ch)

*Permian-Triassic in Oman and Turkey; Late Permian-Early Triassic physical stratigraphy and palynofacies, and correlation based on carbon isotope curves from carbonates and organic matter; Blind Fiord transgression in the high Canadian Arctic; Anisian-Ladinian WG Chair.*

**Hugo Bucher**, Lyon, FRANCE

[Hugo.Bucher@univ-lyon1.fr](mailto:Hugo.Bucher@univ-lyon1.fr)

*Lower Triassic ammonoids, western Guangxi, China; Spiti, North America. Middle Triassic ammonoids of North America.*

**Hamish Campbell**, Dunedin, NEW ZEALAND

[H.Campbell@gns.cri.nz](mailto:H.Campbell@gns.cri.nz)

*Molluscs (especially bivalves) and brachiopods; Stratigraphy, correlation, terrane relations, and biogeography.*

**Yoram Eshet**, Jerusalem, ISRAEL

[yoram@mail.gsi.gov.il](mailto:yoram@mail.gsi.gov.il)

*Palynology of the P-T boundary, focused on the „Fungal Spike“ and reconstruction of the floral succession, Israel and the Karoo Basin, South Africa.*

**Maurizio Gaetani**, Milano, ITALY

[maurizio.gaetani@unimi.it](mailto:maurizio.gaetani@unimi.it)

*Middle Triassic boundaries; Ladinian-Carnian WG Chair.*

**P. John Hancox**, Wits, SOUTH AFRICA

[065PJH@cosmos.wits.ac.za](mailto:065PJH@cosmos.wits.ac.za)

*Non-marine Triassic basins of southern and eastern Africa, particularly the Induan-Olenekian and Olenekian-Anisian boundaries.*

**Dennis Kent**, Palisades, U.S.A.

[dvk@rci.rutgers.edu](mailto:dvk@rci.rutgers.edu)

*Paleomagnetism of nonmarine (Newark basins, East Greenland, Argana) and marine (esp. Tethys); U. Triassic, astronomically-calibrated geomagnetic polarity time scale and global correlation; magnetostratigraphic correlation of M-U. Triassic Tethyan marine sections (with G. Muttoni) and calibration of biochronologies; marine-nonmarine correlation.*

**Sandor Kovács**, Budapest, HUNGARY

[skovacs@IRIS.geobio.elte.hu](mailto:skovacs@IRIS.geobio.elte.hu)

*Middle-Late Conodont biostratigraphy, Balaton Highland; Circum-Pannonian terrane maps and Late Triassic environments; Triassic correlation, Carpatho-Balkans area (Austria to Greece); Triassic of northeast Hungary; A-L and L-C boundaries.*

**Heinz W. Kozur**, Budapest, HUNGARY

[kozurh@helka.iif.hu](mailto:kozurh@helka.iif.hu)

*Conodonts, and their calibration with radiolarian zones and marine ostracodes, esp. Tethys; U. Triassic biostratigraphic scales (conodonts, holothurian sclerites, radiolarians, ostracods) in Silicka Brezova; Germanic continental-marine correlation using conchostracans and ostracods.*

**Leopold Krystyn**, Vienna, AUSTRIA

[leopold.krystyn@univie.ac.at](mailto:leopold.krystyn@univie.ac.at)

*Ammonoids and conodonts; Triassic boundary definitions and resolutions. Norian-Rhaetian WG Chair.*

**Jinling Li**, Beijing, CHINA

[pljinling@263.net](mailto:pljinling@263.net)

*Marine reptiles from Guizhou, China; Environmental change at the P-T boundary; P-T boundary of the Tianshan section using vertebrates, palynology, paleomagnetism, and geochemistry, and correlation with Dalongkou.*

**Vladlen R. Lozovsky**, Moscow, RUSSIA

[vlozovsky@mtu-net.ru](mailto:vlozovsky@mtu-net.ru)

*Stratigraphy of continental red beds of the Russian platform and surrounding areas, and the position of the P-T, O-A, and A-L boundaries.*

**Spencer Lucas**, Albuquerque, U.S.A.

[SLucas@nmmnh.state.nm.us](mailto:SLucas@nmmnh.state.nm.us)

*Tetrapod-based timescale; tetrapods and stratigraphy in the western United States; Triassic of*



northern Mexico; Stratigraphic distribution of tetrapods in the type German Triassic; tetrapod assemblages of the Parana Basin, Brazil; Non-marine WG Chair.

**Manfred Menning**, Potsdam, GERMANY  
[menne@gfz-potsdam.de](mailto:menne@gfz-potsdam.de)  
Magnetostratigraphic reference scale of the Buntsandstein and Keuper (with M. Szurlies); global correlations. German Subcommission for P-T stratigraphy Chair.

**Alda Nicora**, Milano, ITALY  
[aldanic@e35.gp.terra.unimi.it](mailto:aldanic@e35.gp.terra.unimi.it)  
Conodont studies; Middle-Upper Triassic of the southern Alps; Upper Triassic of Sicily, Hydra Island, and Spiti; boundary studies of the O-A boundary of Romania and Greece; A-L of the Southern Alps (with Brack), and L-C of the Himalaya (with Krystyn and Balini).

**Michael J. Orchard**, Vancouver, CANADA  
[morchard@nrcan.gc.ca](mailto:morchard@nrcan.gc.ca)  
Conodont biostratigraphy and biochronology in North America; conodont taxonomy, apparatus structure, and evolution; Cordilleran terranes and paleogeography; O-A boundary in Romania (with Nicora); A-L boundary in Nevada (with Bucher); C-N boundary WG Chair; STS Chair.

**George D. Stanley**, Missoula, U.S.A.  
[fossil@selway.umt.edu](mailto:fossil@selway.umt.edu)

Early Mesozoic reefs; Paleocology and taxonomy of corals and sponges; Cordilleran tectonics and paleogeography of terranes.

**Vijaya**, Lucknow, INDIA  
[bsip@bsip.sirnetd.ernet.in](mailto:bsip@bsip.sirnetd.ernet.in)  
Lower Triassic palynology, Raniganj Godwana Basin, West Bengal.

**Henk Visscher**, Utrecht, THE NETHERLANDS  
[h.visscher@bio.uu.nl](mailto:h.visscher@bio.uu.nl)  
Triassic palynology in chronostratigraphic correlation and classification, Europe; Correlation of Alpine and continental Germanic facies; P-T boundary terrestrial ecosystems (with C. Looy).

**Geoffrey Warrington**, Nottingham, U.K.  
[gwar@bgs.ac.uk](mailto:gwar@bgs.ac.uk)  
Palynology U.K., N. Africa, Middle East; U.K. stratigraphy. STS Secretary. Convenor, Triassic-Jurassic Boundary Working Group.

**Wolfgang Weitschat**, Hamburg, GERMANY  
[weitschat@geowiss.uni-hamburg.de](mailto:weitschat@geowiss.uni-hamburg.de)

Triassic stratigraphy and ammonoids of Svalbard, Spitsbergen; High paleolatitude correlations.

**Hongfu Yin**, Hubei, CHINA  
[hfyin@cug.edu.cn](mailto:hfyin@cug.edu.cn)  
Terrestrial stratotype of the P-T boundary, western Guizhou; P-T boundary events; I-O boundary in China; STS Vice-Chair; P-T boundary WG Chair.

**Yuri D. Zakharov**, Vladivostok, RUSSIA  
[zakharov@fegi.ru](mailto:zakharov@fegi.ru)  
Uppermost Permian and I-O ammonoids from South Primorye; Paleomagnetic studies in the Abrek Bay; STS vice-Chair; Induan-Olenekian WG Chair.

### Corresponding members

**Viorel Atudorei**, Albuquerque, USA  
[atudorei@unm.edu](mailto:atudorei@unm.edu)  
Chemostratigraphy, carbon, sulfur and oxygen stable isotope geochemistry; Triassic sediments, Triassic paleo-oceanography, strontium isotope stratigraphy.

**Marco Balini**, Milan, ITALY  
[marco@e35.gp.terra.unimi.it](mailto:marco@e35.gp.terra.unimi.it)  
Ammonoids of the Tethyan realm; Anisian (S. Alps, Hydra, Spiti, Zanskar); L.-C. (S. Alps, Spiti); L. Triassic (Kçira, Albania; Mangyshlak, Kazakhstan); taxonomy, integrated biostratigraphy/ ammonoids-conodonts-pelagic bivalves, biostratigraphy, palaeobiogeography.

**Michael J. Benton**, Bristol, ENGLAND  
[mike.benton@bristol.ac.uk](mailto:mike.benton@bristol.ac.uk)  
Vertebrate [primarily reptilian] paleontology, particularly the rhynchosaurs of the UK; Russian and Mongolian faunas (with Shishkin).

**Robert B. Blodgett**, Corvallis, USA  
[blodgetr@bcc.orst.edu](mailto:blodgetr@bcc.orst.edu)  
Gastropod faunas from para-autochthonous or cratonic North American rocks, including Sonora, Nevada, British Columbia, and east-central Alaska; accreted terrane faunas from western North America.

**Peter Brack**, Zuerich, SWITZERLAND  
[brack@erdw.ethz.ch](mailto:brack@erdw.ethz.ch)  
A-L boundary (with Hans Rieber) at Bagolino, Italy; correlation of S. Alpine successions with the Germanic Muschelkalk; multidisciplinary studies of platform carbonate cycles and the evolution of the platform-basin system; quantifi-

cation of processes in the Middle Triassic.

**Nikita Bragin**, Moscow, RUSSIA

<mailto:bragin@geo.tv-sign.ru>

*Radiolaria: morphology, taxonomy, biostratigraphy, and palaeogeography; zonal stratigraphy in E. Siberia, E. Mediterranean; high latitude faunas in NE Siberia, their origin and history; geology of deep-water sediments and P/T boundary in these environments.*

**Kiril Budurov**, Sofia, BULGARIA

[budurov@geology.bas.bg](mailto:budurov@geology.bas.bg)

*Triassic stratigraphy, palaeogeography and palaeogeodynamics of the Balkan Peninsula; Definition of Middle Triassic stages and sub-stages and their boundaries based on conodonts; CAI maps for several Triassic substages.*

**Galina I. Buryi**, Vladivostok, RUSSIA

[buryi@mail.ru](mailto:buryi@mail.ru)

*Conodonts from Far East Asia (Sikhote-Alin); biostratigraphy of silicious and carbonate formations; conodont affinities and paleobiology.*

**Elisabeth S. Carter**, Portland, U.S.A.

[escarter@coinet.com](mailto:escarter@coinet.com)

*Dynamics of radiolarian extinction and recovery across the T-J boundary, Queen Charlotte Islands, Baja California, and Japan; radiolarians across the C-N boundary, Queen Charlotte Islands; intercalibration of radiolarian zonation with conodont and ammonoid zones.*

**Jinghua Chen**, Nanjing, CHINA

[jhchen@nigpas.ac.cn](mailto:jhchen@nigpas.ac.cn)

*Bivalves - biostratigraphy, bioevents, paleobiogeography; Lower Triassic recovery and Middle Triassic bivalve radiation in south China.*

**Zhengwu Cheng**, Beijing, CHINA

[ljiivpp@263.net](mailto:ljiivpp@263.net)

*Non-marine P-T boundary at the Dalongkou section of Jimusar and neighbouring areas of Xinjiang; Biostratigraphic and palaeomagnetic studies of the T-J boundary in Lufeng, Yunnan Province.*

**Chonpan Chonglakmani**, Nakhon Ratchasina, THAILAND

[chongpan@ccs.sut.ac.th](mailto:chongpan@ccs.sut.ac.th)

*Stratigraphy, sedimentology, and tectonic evolution of Thailand; systematics of bivalve and ammonoids.*

**Simonetta Cirilli**, Perugia, ITALY

[simocir@unipg.it](mailto:simocir@unipg.it)

*Sequence-stratigraphy and correlation in the*

*Tethys; Sedimentary organic matter (palynofacies, palynostratigraphy, organic facies) and sedimentary facies; marine ecosystems - stratigraphy, evolution, palaeoclimatology, palaeogeography.*

**Dang Tran Huyen**, Hanoi, VIETNAM

*Fauna and stratigraphy of Vietnam.*

**G. Demathieu**, Dijon, FRANCE

*Vertebrate ichnotaxonomy, Middle-Upper Triassic, Spain and France.*

**Jim M. Dickins**, Turner, AUSTRALIA

*Global climates, faunas, events, and correlations; Burrowing bivalves (with N. Morris); T-J boundary, UK.*

**Paulian Dumitrica**, Bern, SWITZERLAND

c/o [Ruth.Dumitrica@bfs.admin.ch](mailto:Ruth.Dumitrica@bfs.admin.ch)

*Radiolarian biostratigraphy in Oman, esp. the Zulla Formation (Upper Anisian-Lower Norian); radiolarian taxonomy, Oman, northern Italy, and Romania.*

**Ashton Embry**, Calgary, CANADA

[AEmbry@nrca.gc.ca](mailto:AEmbry@nrca.gc.ca)

*Stratigraphy and sedimentology of the Sverdrup Basin, northern Canada; sequence boundaries and global tectonics.*

**Douglas H. Erwin**, Washington, USA

[erwin.doug@nmnh.si.edu](mailto:erwin.doug@nmnh.si.edu)

*Early Triassic biotic recovery; high-resolution geochronology (U/Pb) of the Early Triassic (with Sam Bowring); gastropod systematics and diversity patterns (with Alex Nutzel).*

**Jean-Claude Gall**, Strasbourg, FRANCE

[jcgall@illite.u-strasbg.fr](mailto:jcgall@illite.u-strasbg.fr)

*Biological recovery following the Permian mass extinction event, emphasis on strategies developed by the flora and fauna (crustaceans, terrestrial arthropods) to ensure colonization of disturbed habitats; Contributions of microbial communities to taphonomy and sediment genesis.*

**Oscar Gallego**, Corrientes, ARGENTINA

[ofgallego@hotmail.com](mailto:ofgallego@hotmail.com)

*Conchostracans, insects and other continental invertebrates, especially taxonomy of conchostracans; evolution, stratigraphical and palaeogeographical distribution of these groups; comparisons between faunas from Gondwana and Laurasia.*

**Yves Gallet**, Paris, FRANCE

[gallet@ipgp.jussieu.fr](mailto:gallet@ipgp.jussieu.fr)

*Magnetostratigraphy of the O/A and C/N boundaries.*

**Andrzej Gazdzicki**, Warszawa, POLAND

[gazdzick@twarda.pan.pl](mailto:gazdzick@twarda.pan.pl)

*Triassic foraminifera of Tethys (West Carpathians, Himalaya, Papua New Guinea) and the Polish and Spanish Muschelkalk; Rhaetian conodonts from the Tatra Mountains; Early Triassic conodonts from Spitsbergen; Microcoprolites from the Carpathians. Norian-Rhaetian boundary.*

**Piero Gianolla**, Ferrara, ITALY

[glr@unife.it](mailto:glr@unife.it)

*Sequence stratigraphy and paleoclimate of Middle and Upper Triassic successions of Tethys and the correlations with different areas as Arctic and the Germanic basin.*

**Brian F. Glenister**, Iowa City, U.S.A.

[brian-glenister@uiowa.edu](mailto:brian-glenister@uiowa.edu)

*Extinction patterns in key Permian biostratigraphic indicators (ammonoids, fusulinaceans, conodonts) as they relate to the P-T extinction and subsequent recovery.*

**Spela Gorican**, Ljubljana, SLOVENIA

[spela@alpha.zrc-sazu.si](mailto:spela@alpha.zrc-sazu.si)

*Ladinian-Carnian Radiolaria from Oman; Middle-Late Triassic Radiolaria from northern Croatia.*

**Eugen Gradinaru**, Bucharest, ROMANIA

[egradin@geo.edu.ro](mailto:egradin@geo.edu.ro)

*Biostratigraphy (ammonoids, conodonts) and magnetostratigraphy of Desli Caira, Romania; O-A boundary; lithostratigraphy and biostratigraphy of North Dobrogea; Upper Anisian ammonoids of the Brasov and Sasca areas, S. Carpathians.*

**Bruno R.C. Granier**, Abu Dhabi, UAE

[bgranier@emirates.net.ae](mailto:bgranier@emirates.net.ae)

*Algae, taxonomy and stratigraphic range; carbonate sedimentology and stratigraphy.*

**Jack A. Grant-Mackie**, Auckland, NEW ZEALAND

[j.grant-mackie@auckland.ac.nz](mailto:j.grant-mackie@auckland.ac.nz)

*Palynofloras across our T-J boundary (with Zhang Wang-Ping); paleobiogeography of Australasia; Mesozoic macrofauna of Misool Island (with Fauzie Hasibuan).*

**Jean Guex**, Lausanne, SWITZERLAND

[Jean.Guex@igp.unil.ch](mailto:Jean.Guex@igp.unil.ch)

*Lower (Spathian) and Upper Triassic (Rhaetian) ammonoids, western USA; T-J boundary.*

**János Haas**, Budapest, HUNGARY

[haas@ludens.elte.hu](mailto:haas@ludens.elte.hu)

*Sedimentology, Stratigraphy; T-J boundary.*

**Hans Hagdorn**, Ingelfingen, GERMANY

[Encrinus@t-online.de](mailto:Encrinus@t-online.de)

*Crinoid faunas - revision and palaeobiology of families Holocrinidae and Traumatocrinidae; palaeoecology of German Muschelkalk and Lower Keuper (e.g. cephalopod and echinoderm taphonomy, reptile palaeobiogeography and stratigraphy); Muschelkalk stratigraphy.*

**Charles M. Henderson**, Calgary, CANADA

[henderson@geo.ucalgary.ca](mailto:henderson@geo.ucalgary.ca)

*Conodonts and sequence biostratigraphy, western and arctic Canada; L-M Triassic in the subsurface and surface section of western Alberta and northeastern British Columbia; Secretary of the SPS.*

**Francis Hirsch**, Jerusalem, ISRAEL

[Francis.Hirsch@mail.gsi.gov.il](mailto:Francis.Hirsch@mail.gsi.gov.il)

*Late Triassic conodonts from the Chichibu accretionary belt (with K. Ishida); relationship between Tethys and Circum-Pacific faunal assemblages.*

**Mark Hounslow**, Lancaster, ENGLAND

[m.hounslow@lancaster.ac.uk](mailto:m.hounslow@lancaster.ac.uk)

*Magnetostratigraphy of: Lower-Upper Triassic of Svalbard; the proposed T-J GSSP at St Audries Bay; the terrestrial Triassic of the UK and North Sea basin; P-T boundary strata and lower Triassic in eastern Greenland.*

**Yukio Isozaki**, Tokyo, JAPAN

[isozaki@chianti.c.u-tokyo.ac.jp](mailto:isozaki@chianti.c.u-tokyo.ac.jp)

*Permo-Triassic boundary mass extinction event in Japan and in China; conodont, radiolarian, and fusulinacean biostratigraphy, and chemostratigraphy of the boundary beds.*

**Daria Ivanova**, Sofia, BULGARIA

[dariaiv@geology.bas.bg](mailto:dariaiv@geology.bas.bg)

*Foraminifers, calcareous algae - taxonomy, biostratigraphy, microfacies, event stratigraphy, sequence stratigraphy, database, palaeobiogeography, palaeoecology.*

**Volker Jacobshagen**, Berlin, GERMANY

[vojac@zedat.fu-berlin.de](mailto:vojac@zedat.fu-berlin.de)

*Lower Triassic ammonoids, Chios sections (with D. Mertmann).*

**Marji J. Johns**, Victoria, CANADA

[johnsm@telus.net](mailto:johnsm@telus.net)

*Ichthyolith biostratigraphy, paleoenvironments, sequences, and boundaries of the Middle and Late Triassic of NE British Columbia and Canadian Cordillera.*

**Hans Kerp**, Münster, GERMANY

[kerp@uni-muenster.de](mailto:kerp@uni-muenster.de)

*Antarctic palynology; Rhaetian floras, particularly seed ferns.*

**Tatyana V. Klets**, Novosibirsk, RUSSIA

[fossil@lab.nsu.ru](mailto:fossil@lab.nsu.ru)

*Conodonts of northeast Asia (North of Siberia and Far East Russia).*

**Toshio Koike**, Yokohama City, JAPAN

[koike@ed.ynu.ac.jp](mailto:koike@ed.ynu.ac.jp)

*Reconstruction of conodont apparatuses of the Gondolellidae and Ellisonidae in Japan.*

**Tea Kolar-Jurkovsek**, Ljubljana, SLOVENIA

[tea.kolar@geo-zs.si](mailto:tea.kolar@geo-zs.si)

*Biostratigraphy of Slovenia with special interest in Smithian, Carnian, and Norian faunas, including the Raibl beds; Conodonts (stratigraphy, CAI) of Croatia.*

**Galina V. Kotlyar**, St. Petersburg, RUSSIA

[gkotlyar@mail.wplus.net](mailto:gkotlyar@mail.wplus.net)

*Permian brachiopods and the P/T boundary transition.*

**Wolfram M. Kuerschner**, Utrecht,  
THE NETHERLANDS

[w.m.kuerschner@bio.uu.nl](mailto:w.m.kuerschner@bio.uu.nl)

*Palaeobotany and palynology in the Carnian and Norian of Southern Germany and Austria; Editor of *Albertiana*.*

**Jochen Lepper**, Hannover, GERMANY

*Red Beds of the Germanic Triassic; co-ordinator of Buntsandstein Monograph; Dinocephalian fauna from Zimbabwe.*

**Jean Marcoux**, Paris, FRANCE

[marcoux@ipgp.jussieu.fr](mailto:marcoux@ipgp.jussieu.fr)

*Magnetostratigraphy (Austria, Turkey; Saudi Arabia); Paleoenvironments and P-T Pangaea configuration; Neotethyan geodynamic evolution of Oman; Biotic response to mass extinction: the lowermost Triassic microbialites facies; Exotic limestone blocks of Crimea.*

**Claudia Marsicano**, Buenos Aires, ARGENTINA

[claumar@gl.fcen.uba.ar](mailto:claumar@gl.fcen.uba.ar)

*Triassic continental tetrapod faunas, especially temnospondyl amphibians; correlation of continental basins in Argentina; Triassic Stratigraphic Lexicon of Argentina.*

**Adelaide Mastandrea**, Modena, ITALY

[mastandrea@unical.it](mailto:mastandrea@unical.it)

*Late Norian-Rhaetian conodonts from Calabria (Southern Italy); Ladinian-Carnian conodont biostratigraphy from the Dolomites.*

**Christopher R. McRoberts**, Cortland, U.S.A.

[mcroberts@cortland.edu](mailto:mcroberts@cortland.edu)

*Bivalve molluscs - systematics, biochronology and paleoecology; particularly taxic and ecologic diversity through the North American Triassic.*

**Selam Meço**, Tirana, ALBANIA

[smeco\\_2001@yahoo.com](mailto:smeco_2001@yahoo.com)

*Conodonts and stratigraphy of the deposits of the Korabi Zone, Albania; Conodonts and pelagic Triassic stratigraphy of the Cukali Zone, Albania; Anisian conodonts in Albania.*

**Dorothy Mertmann**, Berlin, GERMANY

[mertmann@zedat.fu-berlin.de](mailto:mertmann@zedat.fu-berlin.de)

*Olenekian ammonoids of Chios Island (Greece) from collections of Assereto and Gaetani-Jacobshagen; Sequence-stratigraphy of the Salt Range and Trans Indus Ranges (facies development, biofacies, depositional environments).*

**Ian Metcalfe**, Armidale, AUSTRALIA

[imetcalf@metz.une.edu.au](mailto:imetcalf@metz.une.edu.au)

*Conodonts from the P-T boundary in China (with Bob Nicoll) - multidisciplinary study of the boundary and mass extinction (marine/ non-marine) integrating isotope geochronology, biostratigraphy, chemostratigraphy, and magnetostratigraphy; faunas from Malaysia and Indonesia, and the tectonic evolution of E and SE Asia.*

**Jozef Michalik**, Bratislava, SLOVAKIA

[geolmich@savba.sk](mailto:geolmich@savba.sk)

*Coprolites from the uppermost Triassic, Western Carpathians; Ladinian brachiopods; Lower Carnian orbital rhythms in laminated Hauptdolomit; T-J boundary studies via sequence stratigraphic eustatic signal in Rhaetian Fatra Fm.*

**Paolo Mietto**, Padova, ITALY

[mietto@epidote.dmp.unipd.it](mailto:mietto@epidote.dmp.unipd.it)

*Anisian-Carnian ammonoid (and partially also conodont) biostratigraphy in the Tethys Realm (Southern Alps, Italy), particularly the A-L and L-C boundaries; tetrapod footprints - paleontology and biostratigraphy.*

**Roberto S. Molina Garza**, Querétaro, MEXICO

[rmolina@unicit.unam.mx](mailto:rmolina@unicit.unam.mx)

*Magnetostratigraphy and correlation of non-marine Triassic sequences, southwest USA - Moenkopi Fm., Chinle Gp., Glen Canyon Gp.; Magnetostratigraphy of non-marine P-T boundary in southeastern New Mexico, and Sonora.*

**Atle Mørk**, Trondheim, NORWAY

[atle.mork@iku.sintef.no](mailto:atle.mork@iku.sintef.no)

*Lower Triassic succession of Svalbard,*

*Spitsbergen and the Norwegian Shelf; P-T boundary of Svalbard.*

**Helfried Mostler**, Innsbruck, AUSTRIA  
[helfried.mostler@uibk.ac.at](mailto:helfried.mostler@uibk.ac.at)

*M. Triassic holothurians, Alpine area; lower M-U. Triassic and T-J boundary radiolarians; M. Triassic siliceous sponges, S. Alps; siliceous sponges (Porifera) of Tethys, especially the T-J boundary; P-T boundary in Sicily and Iran, M. Triassic of Turkey), and U. Triassic of Columbia.*

**Giovanni Muttoni**, Milan, ITALY  
[giovanni.muttoni1@unimi.it](mailto:giovanni.muttoni1@unimi.it)

*Magnetostratigraphy aimed at the construction of a Polarity Time Scale; P-T paleogeography from paleomagnetism, esp. Pangea configurations and Tethyan paleogeography; P-T climate analysis: testing the zonal vs monsoonal pangea climate by means of paleolatitude curves correlated with climatic indicators in rocks.*

**K. Nakazawa**, Kyoto City, JAPAN  
*Ammonites and bivalves, Spitsbergen; Salt Range; biostratigraphy of Japan.*

**Norman D. Newell**, New York City, U.S.A.  
[newell@amnh.org](mailto:newell@amnh.org)

*Upper limit of the Paleozoic as revealed by mass extinction and introduction of new Bivalvia; Texan and western American collections from the American Museum of Natural History and the U.S. National Museum.*

**Robert S. Nicoll**, Flynn, AUSTRALIA  
[bnicoll@goldweb.com.au](mailto:bnicoll@goldweb.com.au)  
*Conodonts from the P-T boundary at Meishan and Shangsi, China (with Ian Metcalfe).*

**Omer Faruk Noyan**, Manisa, TURKEY  
[noyan@nil.com.tr](mailto:noyan@nil.com.tr)  
*Conodonts - biostratigraphy, morphogenesis, paleoecology and extinctions.*

**Gilles S. Odin**, Paris, FRANCE  
[gilodin@ccr.jussieu.fr](mailto:gilodin@ccr.jussieu.fr)  
*Radiometric dating and refinement of the time scale.*

**Jim Ogg**, West Lafayette, U.S.A.  
[jogg@purdue.edu](mailto:jogg@purdue.edu)  
*Early Triassic magnetic polarity time scale; integrated magnetic-ammonite-sequence scale for Canadian Arctic sections and correlation to a composite magnetostratigraphy and sequence stratigraphy of the Dolomites region. Secretary General of the ISC.*

**Paul Olsen**, Palisades, U.S.A.  
[polsen@ldeo.columbia.edu](mailto:polsen@ldeo.columbia.edu)  
*Upper Triassic magnetostratigraphy, Newark basins; Carnian-Norian boundary and correlation between terrestrial and marine successions.*

**Qiqing Pang**, Hebei, CHINA  
*Continental P-T boundary and Ostracoda, northern China; Lower and Middle Triassic stratigraphy and ostracodes, Henan Province.*

**Gary Pattemore**, Mt. Nebo, AUSTRALIA.  
*Gymnosperms and stratigraphy, Queensland and other areas of Gondwanaland, e.g. the Molteno Formation of the Karoo Basin in South Africa; habitats and environmental conditions.*

**Rachel A. Paull**, Littleton, U.S.A.  
[rocdox@worldnet.att.net](mailto:rocdox@worldnet.att.net)  
*P-T boundary to Smithian conodont biostratigraphy and depositional history, Rocky Mountains, eastern Great Basin, and Colorado Plateau of the western USA.*

**Maria Cristina Perri**, Bologna, ITALY  
[perri@geomun.unibo.it](mailto:perri@geomun.unibo.it)  
*Permian-Triassic conodonts in the Southern Alps; biostratigraphy and biofacies analysis.*

**Lyudmila Petrounova**, Sofia, BULGARIA  
[lyudmila@geology.bas.bg](mailto:lyudmila@geology.bas.bg)  
*Norian palynomorphs from East Bulgaria; Ladinian-Carnian boundary based on Bulgarian palynomorphs; conodonts.*

**Jozef Pevny**, Bratislava, SLOVAKIA.  
[c/o\\_bystriec@gssr.sk](mailto:c/o_bystriec@gssr.sk)  
*Anisian-Norian conodonts from pelagic facies in central Slovakia; Middle-Upper Triassic ichthyoliths (with Dr. Salaj).*

**Renato Posenato**, Ferrara, ITALY  
[psr@unife.it](mailto:psr@unife.it)  
*Brachiopods and molluscs at the P-T boundary of the S. Alps; bivalves of Anisian plant bearing beds of the Dolomites; biostratigraphy/paleoecology of M. Carnian benthic mollusc & brachiopod assemblages, Dolomites; paleoclimatic changes v. the mid Carnian extinction.*

**Roberto Rettori**, ITALY  
[rrettori@unipg.it](mailto:rrettori@unipg.it)  
*Systematics, biostratigraphy, paleoecology, and evolution of Triassic foraminifers of the Tethyan domain; foraminiferal evolution and extinction across P-T and T-J boundaries; intercalibration of foraminifer ranges with other fossil groups.*

**Hans Rieber**, Zürich, SWITZERLAND  
[rieber@im.unizh.ch](mailto:rieber@im.unizh.ch)  
*Ammonoids and other fossils and stratigraphy of the middle Triassic of the Southern Alps; Definition of the A-L boundary (with P. Brack).*

**John Rigby**, Brisbane, AUSTRALIA  
[j.rigby@qut.edu.au](mailto:j.rigby@qut.edu.au)  
*Megafloras of Gondwanaland; P-T boundary events; tectonic evolution of south east Asia in relationship to distribution of non-marine biotas.*

**Guido Roghi**, Padova, ITALY

[guido@geol.unipd.it](mailto:guido@geol.unipd.it)

*Palynological and paleobotanical analysis of Middle and Upper Triassic sequences of the Julian Alps and Dolomites.*

**Bruce Rubidge**, Wits, SOUTH AFRICA

[106gar@cosmos.wits.ac.za](mailto:106gar@cosmos.wits.ac.za)

*Temnospondyl and Therapsid phylogeny; Biostratigraphy and basin analysis of Karoo-aged rocks from southern Africa.*

**Franco Russo**, Consenza, ITALY

[f.russo@unical.it](mailto:f.russo@unical.it)

*Carbonate diagenesis and biomineralization, especially in the Dolomites; Ladinian/Carnian conodont biostratigraphy of Dolomites; Norian/Rhaetian sequences in Calabria and in Basilicata.*

**Michael R. Sandy**, Dayton, U.S.A.

[Michael.Sandy@notes.udayton.edu](mailto:Michael.Sandy@notes.udayton.edu)

*Triassic brachiopod taxonomy, biogeography, and evolution, North America, southern Europe, and Middle East.*

**Baba Senowbari-Dryan**, Erlangen, GERMANY

[basendar@pal.uni-erlangen.de](mailto:basendar@pal.uni-erlangen.de)

*Algae, foraminifers, palynomorphs, and reefs in Iran; systematic paleontology of coralline sponges in Turkey; Norian reefs in Greece; Carnian Inozoid sponges of Hydra, Greece; biotic studies in Austria, Sicily, and Hungary.*

**Kazem Seyed-Emami**, Tehran, IRAN

[k.seyedemami@kavosh.net](mailto:k.seyedemami@kavosh.net)

*Stratigraphy and ammonoid biozonation.*

**Yasunari Shigeta**, Tokyo, JAPAN

[shigeta@kahaku.go.jp](mailto:shigeta@kahaku.go.jp)

*Systematics, biostratigraphy and evolution of Triassic ammonoids and nautiloids; Induan-Olenekian boundary.*

**Michael A. Shishkin**, Moscow, RUSSIA

[schsz@orc.ru](mailto:schsz@orc.ru)

*Biostratigraphy of the continental Triassic based on tetrapod evidence; Evolution and biogeography of Triassic tetrapods.*

**Milos Siblík**, Suchdol, CZECH REPUBLIC

[inst@gli.cas.cz](mailto:inst@gli.cas.cz)

*Upper Triassic brachiopods, especially Alpine Kössen Beds and the Dachstein Limestone of Hochschwab; Complement to Fossilium Catalogus Austriae: Brachiopoda triadica.*

**Norman J. Silberling**, Lakewood, USA

[slbrlng@email.msn.com](mailto:slbrlng@email.msn.com)

*Ammonoids and Stratigraphy, USA*

**Evgeny S. Sobolev**, Novosibirsk, RUSSIA

[alfalfa@cgi.nsk.su](mailto:alfalfa@cgi.nsk.su)

*Rhaetian deposits in Eastern Taimyr; Triassic nautiloids, North Dobrogea (Romania), Primor'ye (Far East), and North Siberia; Lower Anisian ammonoids, Eastern Taimyr; Paleobiogeographic differentiation of nautiloids.*

**Maureen B. Steiner**, Laramie, U.S.A.

[magnetic@uwyo.edu](mailto:magnetic@uwyo.edu)

*The sequence of magnetic polarity changes near the P-Triassic boundary; Late Carnian through Rhaetian sequence of magnetic reversal changes; comprehensive summary of the Late Permian-Early Triassic magnetic reversal history.*

**Milan N. Sudar**, Belgrade, YUGOSLAVIA

[sudar@EUnet.yu](mailto:sudar@EUnet.yu)

*Triassic conodonts and foraminifers; Sedimentological-paleontological investigations of carbonate platform deposits and red nodular limestones in the Yugoslav Dinarides; Ophiolitic melanges; Correlation within the Mediterranean region; boundary resolution.*

**David Taylor**, Portland, USA

[blitz124@home.com](mailto:blitz124@home.com)

*Norian-Rhaetian ammonoid biochronology and evolution, Western USA; Early Triassic Thaynes Formation and its ammonoid fauna in Nevada, Idaho and Utah.*

**Kagan Tekin**, Ankara, TURKEY

[uktekin@yahoo.com](mailto:uktekin@yahoo.com)

*Biostratigraphy and systematics of late Middle to Late Triassic radiolarians from Turkey.*

**R.S. Tiwari**, Bhopal, INDIA

c/o [shailendra\\_singh\\_75@yahoo.com](mailto:shailendra_singh_75@yahoo.com)

*P-T boundary in the nonmarine sequence of the Indian peninsula; Tagging of palynozones with marine datum of Tethyan Himalayan and creation of a palyno-chronological scale for nonmarine Gondwana sequence of India; intra-Gondwanaland correlations.*

**Jinnan Tong**, Hubei, CHINA

[jntong@cug.edu.cn](mailto:jntong@cug.edu.cn)

*Lower Triassic stratigraphy and paleontology.*

**E. Timothy Tozer**, Vancouver, CANADA

[timozer@nrcan.gc.ca](mailto:timotozer@nrcan.gc.ca)

*Canadian ammonoids and global correlations.*

**John Utting**, Calgary, CANADA

[JUtting@NRCan.gc.ca](mailto:JUtting@NRCan.gc.ca)

*Palynology of the Permian -Triassic boundary and Lower Triassic of Canada.*

**Carmina Virgili**, Barcelona, SPAIN

*Triassic of Spain and its relations with Occidental Europe; Continental deposits of the Lower and Middle Triassic; Problems of the P-T boundary in the continental series.*

**Attila Voros**, Budapest, HUNGARY

[voros@zoo.zoo.nhmus.hu](mailto:voros@zoo.zoo.nhmus.hu)

*Anisian-Ladinian ammonoid stratigraphy, Hungary; A-L boundary.*

**Bruce Waterhouse**, Christchurch, NEW ZEALAND

c/o [loris@xtra.co.nz](mailto:loris@xtra.co.nz)

*E. Triassic Claraia from Nepal; Anisian ammonoids from south NZ; mapping M. Triassic in the S. Alps; reinterpreting E. Triassic of NZ as M. Triassic from ammonoids; E. & M. Triassic ammonoid succession, western and central Nepal (7 part Palaeontographica series).*

**Jobst Wendt**, Tübingen, GERMANY

[jobst.wendt@uni-tuebingen.de](mailto:jobst.wendt@uni-tuebingen.de)

*Carnian mud mounds in Sichuan, China - biostratigraphy, sedimentology, palaeoecology, and diagenesis (stable isotopes) of the mounds and off-mound lithologies.*

**Paul B. Wignall**, Leeds, ENGLAND

[P.Wignall@earth.leeds.ac.uk](mailto:P.Wignall@earth.leeds.ac.uk)

*Boundary studies: P-T boundary in western Canada, E Greenland, S. Tibet; Scythian stage boundaries; T-J boundary in England and S. Tibet.*

**Zunyi Yang**, Beijing, CHINA

[zunyi@public.bta.net.cn](mailto:zunyi@public.bta.net.cn)

*Compilation of the Lexicon of the Chinese Triassic System, and the Chinese Triassic System.*

**Jin Yugan**, Nanjing, CHINA

[ygjin@public1.ptt.js.cn](mailto:ygj@public1.ptt.js.cn)

*Stratigraphy and events of the basal Triassic; Brachiopods from western China.*

**Ivan Zagorchev**, Sofia, BULGARIA

[zagor@geology.bas.bg](mailto:zagor@geology.bas.bg)

*Triassic stratigraphy, palaeogeography, tectonics and palaeogeodynamics of the Balkan Peninsula.*

**Ana Maria Zavattieri**, Mendoza, ARGENTINA

[amz@lab.cricyt.edu.ar](mailto:amz@lab.cricyt.edu.ar)

*Palynology and biostratigraphy of Triassic Basins of Argentina.*

**John-Paul Zonneveld**, Calgary, CANADA

[jzonneve@nrcan.gc.ca](mailto:jzonneve@nrcan.gc.ca)

*Ichthyology and sequence biostratigraphic evolution of the Triassic of western Pangea; in Western Canada: Griesbachian-Dienerian and Ladinian trace fossil assemblages; brachiopod-echinoderm biostromes (Ladinian); coral patch reefs (Carnian); new taxa of lobster (Ladinian).*

## GUIDELINES FOR THE SUBMISSION OF MANUSCRIPTS TO ALBERTIANA

Albertiana is published twice a year. Contributions should be sent to the editor. In order to facilitate the production of this newsletter and reduce typing errors, authors are kindly requested to submit their contributions electronically, preferably by email or on 3½ inch MS-DOS formatted floppy discs together with a printed hard copy. You may also use the Utrecht Albertiana FTP server to upload larger files that cannot be sent as email attachments. The user name is "Alb", even so the password is "Alb". Please DO NOT use BinHex-encoded files (MacIntosh) as these cannot be read! Those who are unable to submit a manuscript in electronic format are kindly requested to send flat (unfolded), clearly typed manuscripts in a 12-point typeface (sans serif) with single line spacing.

Text files can be submitted formatted as \*.wpd, \*.doc or \*.rtf files and illustrations as pixel based graphics (e.g: \*.bmp, \*.tif, \*.gif or \*.jpeg) or vector based graphics (e.g: \*.ai, \*.cdr) that can be directly imported into Adobe PageMaker. Please provide good, clean, flat, printed copies (NOT xerox copies) of any illustrations, which MUST be designed to fit on an A4 page (centered, with at least 2.54 cm wide margins left and right, and 4 cm margins at the top and bottom).

Special attention should be paid to grammar and syntax - linguistic corrections will be minimal. In case of doubt, send your manuscript to a colleague for proof reading. References should be in the format used in the 'New Triassic Literature' section in issue 25 of Albertiana. Please write all Journal titles in full length. The use of names of biostratigraphic units should be in accordance with the International Stratigraphic Guide:

The formal name of a biostratigraphic unit should be formed from the names of one, or preferably no more than two, appropriate fossils combined with the appropriate term for the kind of unit in question."

The writing and printing of fossil names for stratigraphic units should be guided by the rules laid down in the International Code of Zoological Nomenclature and in the International Code of Botanical Nomenclature. The initial letter of generic names should be capitalized; the initial letter of the specific epithets should be in lowercase; taxonomic names of genera and species should be in italics. The initial letter of the unit-term (Biozone, Zone, Assemblage Zone) should be capitalized; for example, Exus albus Assemblage Zone."

The name of the fossil or fossils chosen to designate a biozone should include the genus name plus the specific epithet and also the subspecies name, if there is one. Thus Exus albus Assemblage Zone is correct. After the first mention, the genus name may be abbreviated to its initial letter if there is no danger of confusion with some other genus beginning with the same letter; for example, Exus albus may be shortened to E. albus. On the other hand, the use of the specific epithet alone, in lowercase or capitalized, in italics or not (albus Assemblage zone, Albus Assemblage zone, albus Assemblage zone, or Albus Assemblage zone), is inadvisable because it can lead to confusion in the case of frequently used species names. However, once the complete name has been cited, and if the use of the specific epithet alone does not cause ambiguous communication, it may be used, in italics and lowercase, in the designation of a biozone; for example, uniformis Zone."

From: Salvador, A. (ed.), 1994. International Stratigraphic Guide. Second Edition. International Commission on Stratigraphic Classification of IUGS International Commission on Stratigraphy. IUGS/GSA, Boulder, Co, p. 66.



## Subcommission on Triassic Stratigraphy

**STS Chairman**

Dr. Mike Orchard, Geological Survey of Canada, 101-605 Robson Street, Vancouver, British Columbia, Canada.

**Vice Chairman**

Dr. Yuri D. Zakharov, Far Eastern Geological Institute, Far Eastern Branch, Russian Academy of Sciences, Prospect Stoletiya Vladivostoka 159, 690022 Vladivostok, Russia.

**Vice Chairman**

Dr. Ying Hongfu, Office of the President, China University of Geosciences, Yujiashan, Wuhan, Hubei, 430074, People's Republic of China

**STS Secretary-General**

Dr. Geoffrey Warrington, British Geological Survey, Keyworth, Nottingham NG12 5GG, United Kingdom.

ALBERTIANA is published twice a year by the Subcommittee on Triassic Stratigraphy. Individuals can obtain ALBERTIANA for the sum of US \$ 20,- or EURO 20,- per year. Readers are kindly requested to pay their annual contribution timely.

European readers can send a Eurocheque made payable in Euro to Dr. Zwieterman, Laboratory of Palaeobotany and Palynology, Utrecht University, Budapestlaan 4, 3584 CD Utrecht, The Netherlands. Everyone else is kindly requested to send cash in a closed non-transparent envelope to the above Utrecht address. Because of the high provision costs of other cheques/currencies, other methods of payment cannot be accepted. Institutions can receive ALBERTIANA on an exchange basis.

All correspondence regarding the distribution of ALBERTIANA should be sent to Dr. Z. Zwieterman.

**DEADLINE FOR ALBERTIANA 27  
CONTRIBUTIONS IS:  
June 1<sup>st</sup> 2002**