





DECEMBER 2014 ISSN 0619-4324

ALBERTIANA 42 · DECEMBER 2014

CONTENTS

- The global marine Permian-Triassic boundary: over a century of adventures and controversies (1880–2001) 1 Aymon Baud
- Marine Lariosaurus (Sauropterygia) records from the Middle Triassic of the Germanic Basin: stratigraphic22and paleobiogeographic importance for Tethyan and Germanic Basin correlation22Cajus G. Diedrich23
 - A summary of new conodont biostratigraphy and correlation of the Anisian (Middle Triassic) strata in British Columbia, Canada. *Martyn L. Golding, Michael J. Orchard and John-Paul Zonneveld*
 - Time calibrated Early Triassic $\delta^{13}C_{carb}$, $\delta^{18}O_{apatite}$ and SST curves from South China: an update 41 Nicolas Goudemand
 - Note on the conodonts from the Induan-Olenekian boundary. *Nicolas Goudemand* 49
- New and corrected evidences on early Triassic biostratigraphy in South Primorye, Siberia, Kazakhstan and Vietnam (report on the IOBWG activity in 2014). *Yuri D. Zakharov* 52
 - Obituary: Carmina Virgili (1927–2014) 54
 - New Triassic literature. *Geoffrey Warrington* 56
 - Albertiana Forum 86
 - Publication announcement: Stratigraphie von Deutschland XI: Buntsandstein 88
 - Meeting announcements 89

Editor Christopher McRoberts State University of New York at Cortland, USA

Editorial Board

Aymon Baud Université de Lausanne, Switzerland

Piero Gianolla Università di Ferrara, Italy *Arnaud Brayard* Université de Bourgogne, France

Mark Hounslow Lancaster University, United Kingdom

ral History, Geological Survey of Canada, Vancouver Canada

Spencer Lucas New Mexico Museum of Natural History, USA

Yuri Zakharov Far-Eastern Geological Institute, Vladivostok, Russia

Albertiana is the international journal of Triassic research. The primary aim of Albertiana is to promote the interdisciplinary collaboration and understanding among members of the I.U.G.S. Subcommission on Triassic Stratigraphy. Albertiana serves as the primary venue for the dissemination of orignal research on Triassic System. Albertiana also serves as a newsletter for the announcement of general information and as a platform for discussion of developments in the field of Triassic stratigraphy. Albertiana thus encourages the publication of contributions in which information is presented relevant to current interdisciplinary Triassic research and at provides a forum for short, relevant articles including. original research articles, reports on research and works in progress, reports on conferences, news items and conference announcements, Albertiana Forum: letters, comment and reply, and literature reviews. Albertiana is published bianually by SUNY Cortland's Paleontological Laboratory for the Subcommission on Triassic Stratigraphy. Albertiana/

Cover Image: Waagen's (1895) lithographic plate of the Ceratite limestone from Khoora section, Salt Range. See Baud, p. 3

Marco Balini Università di Milano, Italy

Margaret Fraiser University of Wisconson Milwaukee, USA

> *Wolfram Kürschner* Univseristy of Oslo, Norway

THE GLOBAL MARINE PERMIAN-TRIASSIC BOUNDARY: OVER A CENTURY OF ADVENTURES AND CONTROVERSIES (1880–2001)

Aymon Baud

BGC, Parc de la Rouvraie 28, CH-1018 Lausanne, Switzerland. Email: aymon.baud@unil.ch

FOREWORD

Different views have been recounted on the Permian-Triassic boundary story. Here are considered only some aspects of a long story and for a more full account, especially of the early development of the Permian-Triassic boundary, I refer to some parts the Tozer's (1984) book. As researcher in IGCP projects from 1974, vice-chairman, chairman and past-chairman of the Subcommission on Triassic Stratigraphy (1984–2000), I provide here an insider point of view.

INTRODUCTION FROM GRIESBACH TO DIENER: THE DISCOVERY OF THE OTOCERAS AMMONOID IN THE HIMALAYA

Carl Ludloph Griesbach

The story starts in 1880, when Carl Ludloph Griesbach (Fig. 1) was publishing a note on the Paleontology of the Lower Triassic of the Himalaya: during his Survey of the Central Himalaya, he discovered the *Otoceras* ammonoid (Fig. 1) and he was the first who understood the importance of this fossil to mark the base of the Triassic Series and the Permian-Triassic boundary. His Central Himalaya's studies were published in 1891.



Figure 1 - Carl Ludloph Griesbach with its Otoceras original plate (Griesbach, 1880).

Edmond von Mojsisovics, Wilhelm Waagen and Carl Diener

Shortly after Griesbach's publication came out the Mojsisovics, Waagen and Diener 1895 paper with the Lower Triassic subdivision proposal made by Waagen and Diener (Fig. 2) in which they strongly advised the use of the *Otoceras* Beds described in the Central Himalaya to define the base of the Triassic. In the same year, Wilhelm Waagen published an impressive paleontological work on the Salt Range fossils (Fig. 3). This was soon followed by the contributions of Carl Diener who explored the Central Himalaya and published his survey in 1895 (Fig. 4). Like Waagen before him, Diener visited the Salt Range, then the Permian-Triassic sections in Kashmir. He discovers also *Otoceras* specimens in Spiti Valley sections and made correlations with the Shalshal Cliff section in the Central Himalaya (Diener, 1912). All these extensive paleontological collections have been carefully stored at the Geological Survey of India in Calcutta.

The first controversy starts with von Krafft (1901), Frech (1902) and Noetling (1905) who assumed the Permian age of *Otoceras* with references to the Transcaucasia sections.

	Stufen	Unterstufen	Mediterrane Triasprovinz		Indische Triasprovinz		
Serien			Zone (der pelagischen Facies) Schichtbezeichn (verschiedenarti örtl. Entwicklu		Zone (der pelagischen Facies)	Schichtbezeichnung (verschiedenartiger örtlicher Entwicklung)	
Dinarisch	Anisisch	Bosnisch	10. Z. des Ceratites trinodosus	Oberer Muschelkalk	Z, des Ptychites rugifer	Muschelkalk des Himalaya	
		Balatonisch	9. Z. des Ceratites binodosus	Unterer Muschelkalk	Z. des Sibirites Prahlada	Brachiopoden - Schichten mit <i>Rhynchonella Gries-</i> bachi (Himalaya)	
	Hydaspisch				8. Z. des Stephanites superbus	Obere Ceratiten-Kalke der Salt Range	
Skythisch	Jakutisch		Z. des Tirolites Cassianus	Werfener Schichten	7, Z. des Flemingites Flemin- gianus 6, Z, des Flemingites radiatus	Ceratiten- Sandstein der Salt Range	
					5. Z. des Ceratites normalis		
	Brahmanisch			der Ostalpen	4. Z. des Proptychites trilo- batus	Ceratite Marls der Sult Range Untere Ceratiten-Kalke der Sult Range	
		Gandarisch			3. Z. des Proptychites Law- rencianus		
					2. Z. des Gyronites frequens		
		Gangetisch			1. Z. des Otoceras Woodwardi	Otoceras Beds des Himalaya	

Untere Trias nach W. Waagen und C. Diener.

Figure 2 - The Lower Triassic scale according to Waagen and Diener in Mojsisovics et al. (1895).



Figure 3 – Left, cover of the Salt Range fossils by Waagen (1895); right: plate XL of Waagen (1895) showing a limestone slab of the Lower *Ceratite* limestone, Khoora section.





Figure 4 - The cover of the Central Himalaya paper of C. Diener (1895).

THE ARCTIC ADVENTURES, AMMONOIDS AND MIXED FAUNA

Carl Diener

The Permian-Triassic succession and boundary of three main areas of the Western Arctic have been intensely studied (Fig. 5, left): Jameson Land on East Greenland with the expeditions of Lauge Koch in the 1930's (Fig. 5, right), of R. Trümpy in the 1950's, and of C. Teichert and B. Kummel at the end of the 1960's; Axel Heiberg and Ellesmere Islands surveyed by E.T. Tozer; and Spitsbergen (Svalbard) Islands studied during three expedition of a Japanese team leaded by Nakamura (1984) and by K. Nakazawa in 1986.



Figure 5 – Left, Western Arctic map: 1– Jameson Land, 2- Axel Heiberg and Ellesmere Islands, 3- Spitsbergen (Svalbard) Islands; right: the "Gustave Holmes" boat of the Lauge Koch Expedition close to Jameson Land (photo from Augusto Gansser, a young geologist of the expedition).

Lauge Koch collections and Leonard Frank Spath

Between 1930 and 1950, large Cephalopod collections from Eastern Greenland made by Lauge Koch (1931) were carefully studied by Leonard Frank Spath (Fig. 6), allowing him to publish a new and very fine subdivision of the Lower Triassic from the Arctic region. Working on ammonoid biochrones, his basal Triassic was called Otoceratan (Spath, 1934, Spath et al., 1951).

Rudolf Trümpy

At the end of the 1950s, Rudolf Trümpy, Professor in Lausanne and Zürich, Switzerland went to East Greenland (Jameson Land, Kap Stosch), studied the late Permian to basal Triassic macrofossils and, in 1960, he wrote an interesting study about the problematic of the mixed fauna at the Permian-Triassic transition. Later, in 1969, Trümpy published his work on the Lower Triassic Ammonites from Jameson Land (Fig. 7) with refined subdivisions of the local basal Triassic.



Figure 7 - Cover of the 1969 Jameson Land Triassic paper with the Trümpy's ammonites study.

Edward Timothy Tozer

In the sixties, a Canadian paleontologist ammonoid specialist, Edward Timothy Tozer (Fig. 8, left), started his famous Triassic studies in the High Arctic, for the Geological Survey of Canada (Fig. 8, right). As only geographic names are allowed for stage name, Tozer gave the name of the pioneers of the Triassic studies to small gullies or creeks, both on Axel Heiberg and on Ellesmere Island (Fig. 9). That is why, in 1965, he took the opportunity to subdivide the Lower Triassic Series in four stages, named from these creeks: Griesbachian, Dienerian, Smithian and Spathian.

With Norman Silberling, the ammonoid specialist from Denver (USA), Tozer extended the Arctic lower Triassic subdivisions to the whole of North America and published (Tozer and Silberling, 1968), as point of reference, the biochronological scale and the subdivisions of the Western North America Triassic (Fig. 10).



Figure 8 – Left, Tim Tozer surprised at the PTB in Nammal Gorge (Salt Range, Pakistan, December 1987); right, cover of the Tozer's (1967) Triassic book.



Figure 9 – The four remote gullies (Creeks) used by E.T. Tozer as lower Triassic stages stratotypes.



Figure 10 – Left, cover of the Silberling and Tozer (1968) basic Triassic paper; right, Tozer's (1965, table 2) Lower Triassic comparative classification with: 1 = Silberling and Tozer (1968), 2= Kiparisova & Popov (1956, 1961), 3 = Kiparisova & Popov (1964), 4 = Spath (1934), 5 = Kummel (1957).

Curt Teichert and Bernhard Kummel

During the 1967 Summer, Curt Teichert specialist of Permian Cephalopods and Bernhard Kummel specialist of Triassic ammonoids get the opportunity to go and work on the Permian Triassic transition in the famous Kap Stosch area. They finally went to the conclusion that the Permian fossils with brachiopods

> BULLETIN OF CANADIAN PETROLEUM GEOLOGY VOL. 20, NO. 4 (DECEMBER, 1972) P. 659-675

PERMIAN-TRIASSIC BOUNDARY IN THE KAP STOSCH AREA, EAST GREENLAND

CURT TEICHERT¹ AND BERNHARD KUMMEL²

ABSTRACT

Study of a number of Permian-Triassic sections in the Kap Stosch region in the summer of 1967 has provided a number of new observations. First, the Permian-Triassic sections southwest of Kap Stosch are of homogeneous shale, silty shale, and siltsone facies. None of the rock units are highly indurated, but all are markedly "soft." Solifluction has so badly affected all outcrops that meaningful stratigraphic sections are next to impossible to obtain. The lowest Triassic beds in these regions do contain thin (1-5 cm) hard bands consisting of coquinas of anmonoids (*Glyptophiceras* and *Otoceras*) and containing fragments of productids, bryozoans, and other fossils of "Permian" affinities.

Southeast of Kap Stosch (e.g., between Rivers 6 and 14) the lowest Triassic strata encompassing the *Gluptophiceras* Zone are about 200 m thick. They are predominantly arkote sandstone and conglomerate. A number of horizons yield fragmentary and whole specimens of productid brachlopods, fragments of crinold stems, bryozoans and other fossils in coarse sandstone and conglomerate matrix, occurring as much as 100 m above the base of the Triassic sequence.

These strata, containing mixed as not in above the base of the Thassic sequence. These strata, containing mixed associations, by their thickness, sedimentary structures, and composition, clearly indicate very rapid rates of deposition. These environmental considerations lead to the conclusion that the "Permian" faunal elements almost certainly did not actually live and form part of the benthos during earliest Triassic time. The underlying Permian formations are of diverse facies among which richly-fossiliferous biohermal banks are present elsewhere. Some of these banks weather easily, yielding nearly perfectly preserved fossils free of matrix. We consider it most probable that some of the Permian faunal elements in the lowest Triassic formations have been brought into that environment as argillaceous boulders, that once coming to rest, dissolved, leaving well-preserved fossils that were rapidly buried in the coarse sediment and in a free state were transported very little. The majority of fossils, however, were washed out of soft rocks and were badly broken during transportation.

We conclude that the Permian-Triassic sequence in the area encompasses a break equivalent to at least the Dzhulfian Stage.

and bryozoans are reworked within the basal Triassic *Ophiceras* beds (so called mixed fauna). The first paper was published in 1972 (Fig. 11), and the detailed description of the sections appear in their 1976 paper with the conodont description in an appendix written by Walter C. Sweet.



Figure 11 - Left, first page of the Teicher and Kummel's (1973) Kap Stosch paper; right, their plate with Ophiceratids accumulation.

Koji Nakamura, Keiji Nakazawa and the Japanese Group

As reported by Nakamura et al. (1987), a cooperative study between Norsk Polarinstitutt and Hokkaido University, Japan was undertaken during the year 1984 in West Central Spitsbergen. The main objective of which was to obtain detailed stratigraphical and palaeontological informations on the Permian and the Permian-Triassic boundary (Fig. 12). During the Summer 1986, Keiji Nakazawa leaded a Japanese geological expedition to Svalbard with a report of *Otoceras boreale* finding in 1987. Further studies were made in 1990, and the final Report published by Tatsumi (ed.) in 1990 with the Nakamura et al. paper on Permian-Triassic boundary.

THE PERMIAN-TRIASSIC BOUNDARY

Keiji Nakazawa

Introduction

There are two problems with regard to the Permian-Triassic boundary. One is the correlation of the Permian-Triassic transition beds, or mixed fauna beds. The other is the location of boundary itself. An accurate correlation of the transition beds needs first to be cleared before the second problem can be discussed. The correlation of the *Otoceras* Zone is of special importance, because it has classically been defined as the base of the Triassic System. However, its distribution is limited to the arcto-boreal and peri-Gondwana provinces and the correlation with the Tethyan province is still in dispute. The correlation of the so-called mixed fauna beds is also of importance, but these two problems are intimately related with each other.



Figure 12 – Left, Introduction of the PTB paper of Nakazawa (1992); right, a Permian-Triassic transition section in Spitsbergen

THE CENTRAL TETHYS

Research on Permian-Triassic transition along the central Tethys margins concerned Transcaucasia (Armenia and Azerbaidjan), Iran, Pakistan, Northern India and Tibet.

In the Salt Range, they started again in the 1950s, with Schindewolf short survey published in 1954 and were followed in the sixties by Curt Teichert, Bernhard Kummel and Walter Sweet, and later by Keij Nakazawa and his Japanese team.

Transcaucasia; from Hermann Abich (1878) to Tim Tozer (1969)

Due to the highly fossiliferous late Paleozoic, the Transcaucasia attracted paleontologists as Abich (1878) and Mojsisovics (1879). For Frech and Arthaber (1900) the Himalayan *Otoceras* were Permian in age as the Armenian one's. Bonnet (1919, Fig.13) found large similarities between the *Otoceras joulfense* Abich and the Himalayan *Otoceras woodwardi*, and fighting against Diener, he strongly claimed a same Eo-Triassic age of the Armenian red "*Otoceras*" beds. Due to the abundance of Xenodiscidae and early Otoceratidae, the area became source of biochronological problems and misplaced correlation. The story continued and later Ruzhentsev and Sarycheva (1965), Russian paleontologists incorrectly determined a basal Triassic *Tompophiceras* zone below the *Dzhulfites* and *Paratirolites* beds. Influenced by this, D.L. Stepanov, F. Golshani and J. Stöklin (1969) published, on the Iranian Julfa sections, a wrong correlation with a Permian-Triassic boundary in the middle of the red ammonoid limestone with *Productus*. Fortunately, after carefully studying Armenian specimens, Tozer, the same year (1969), reinterpreted the data and went to the right conclusion that the Permian-Triassic boundary is to be placed above the red ammonoid limestone, at the top of the *Paratirolites* beds. A long controversy was resolved.

Curt Teichert and Bernhard Kummel in association with Walter C. Sweet

In the 1960s, Curt Teichert specialist of Permian Cephalopods, Bernhard Kummel specialist of Triassic ammonoids and Walter C. Sweet conodont specialist started to work together on Permian-Triassic boundary sections of the Tethys: first in the Salt-Range, then in Kashmir and finally in the Julfa section of NW Iran. Both, separately or together, published very detailed reports on examined sections and their macrofossil contents.

Curt Teichert and Bernhard Kummel undertook field work in the Salt Range sections, Pakistan, in 1961 up to1964, and Kummel published a first report in 1966, followed by an edited book (Teichert and Kummel, 1970) that contains the description of 12 main sections with the Permian-Triassic transition (Fig. 14). Walter C. Sweet (1970) wrote a section on the lower Triassic conodont succession.



Pierre Bonnet



Figure 13 - Cover of the book on the story of the Pierre Bonnet geological explorations in Transcaucasia, from 1909 to 1914.



Figure 14 – Left, C. Teichert photo and B. Kummel portrait; middle, cover of the Kummel and Teichert book (1970) on the Salt Ranges; right: slide of the Central part of the Salt Range with, in the foreground, the Lower Triassic strata worked by Teichert and Kummel.



Figure 15 - Left, view on the Guryul Ravine section; right: the Guryul Ravine section drawn by Teichert & Kummel (1970).

In 1968, Curt Teichert and Bernhard Kummel spend 2 days at the Guryul Ravine section, Kashmir, India (Fig. 15), collected samples and wrote a paper (Teichert & Kummel, 1970). The same year, Sweet gave an account on the conodont zonation from late Permian to early Triassic at Guryul Ravine (Sweet, 1970).

After short reconnaissance on Iranian side of the Transcaucasia in 1966, Teichert went back with Bernhard Kummel in Julfa area in 1968 and studied four sections along the slope. The Teichert et al. paper was printed in 1973 (Fig. 16, left) and took account of the Tozer's views for interpreting the PTB position. The "upper Julfa beds" was sampled in their locality 4 (Fig. 16, right), packed, sent and stored at the Museum of Comparative Zoology, but probably wrongly labeled with the name of the overlying unit "Ali Bashi Formation". This mistake has been the starting point of controversies on the late Permian conodont succession and determination, explained by Henderson et al., 2008 (this story is related by Baud, 2008 and recently by Ghaderi et al., 2013).



Figure 16 – Left, cover of the Kummel and Teichert (1973) booklet on Julfa sections; right, view on the locality 4 of Kummel and Teichert.

Keiji Nakazawa and the Japanese Group

In 1969 and during the seventies, a Japanese Group leaded by Keiji Nakazawa, in cooperation with local Survey, started extensive studies of three main Permian–Triassic marine and fossiliferous area of the central Tethys and published bed by bed description and fossil contents.

In 1972, detailed paleontological and sedimentological studies have been carried out in Abadeh, Central Iran (Fig. 17, right) and in Elburz Mountain by geologists of the Geological and Mineral Survey of Iran and by a research staffs of Japanese universities leaded by Keiji Nakazawa. Supplementary fieldwork was done in 1975 and the extensive results (Fig. 17, left) were published in the Kyoto University Memoirs (Iranian-Japanese Research Group, IJRG, 1981)

With Hari Mohan Kapoor from Indian Survey, the Japanese

Group leaded by Keiji Nakazawa studied the Guryul Ravine section, (Fig. 18, right) and surrounding sections of the Indian Kashmir, during 1971 and 1972. The extensive results were published in the Kyoto University Memoirs (Nakazawa et al., 1975), the fauna in Nakazawa and Kapoor, 1981 (Fig. 18, left), with the ammonoids systematic description by Bando, and during the 80's, Tetsuo Matsuda (1981, 1982, 1983 and 1984) wrote detailed conodont papers on the Kashmir Permian-Triassic succession.

With colleagues from the Pakistani Geological Survey, the Japanese Group leaded by Keiji Nakazawa studied in 1975, 1976 and 1979 the Nammal Gorge and seven surrounding sections in the Salt and the Surghar Ranges, Pakistan (Fig. 19, right). The first results were published by Pakistanese-Japanese Research Group (PJRG), first in 1981 (Fig. 19, left) and later in a 90 pages detailed report (PJRG, 1985).



Figure 17 - Left, cover of the Kyoto University Memoirs on Abadeh (IJRG, 1981); right: view of the Abadeh section.



Figure 18 – Left, Cover of the Kashmir book (1981); right: slide of the Guryul Ravine Permian-Triassic section with the Nakazawa beds numbering and PTB (red line).



Figure 19 – Left, cover of the first Salt Range paper (PJRG, 1981); right: view on the Surgar Range with upper Permian and lower Triassic in the ground.

THE INTERNATIONAL GEOLOGICAL CORRELATION PROGRAM (IGCP), CONFERENCES AND FIELD WORKSHOPS (1974–2001)

Three years prior to the beginning of the International Geological Correlation Program, a turning point event in the Permian-Triassic boundary studies was the first International Permian-Triassic Conference held August 23-26, 1971 in Calgary (Canada) with over fifty contributions on the paleontology, stratigraphy, sedimentology, radiometric chronology, paleoclimatology and paleomagnetism of this time interval and of the boundary. A. Logan and L. V. Hills edited

the Proceeding entitled "The Permian and Triassic Systems and their Mutual Boundary" in 1972.

For nearly a century, the research of this boundary mainly dealt with macrofossils and associated biochronology as tool for correlation. Microfossils as conodont systematic and geochemical analyses were only at the opening as magnetostratigraphy.

Heller at al. (1988) gave a first study on marine PTB magnetic time scale on Shangsi section (Sichuan, China), a candidate for the Permian-Triassic boundary section. Later on, Ogg and Steiner (1991) published their data on the Lower Triassic stratotype of Ellesmere and Axel Heiberg Islands, showing a normal polarity within the *Otoceras* zones.

IGCP Projects 4 and 106 (1974-1984), Figs. 20 and 21 left.

Sponsored by UNESCO, in 1974 started the International Geological Correlation Program with the Project 4 on "Triassic of the Tethys Realm", leaded by Prof. H. Zapfe from Vienna University and Austrian Academy. The Project 106 "Permo-Triassic stage of geological evolution" followed it up to 1984, leaded by a Russian team with Dr. A.N. Oleynikov.

Two successful fieldtrips on Permian-Triassic boundary of new areas, the Setorym Creek in Verkoyansk mountains (East Siberia) and the Vedi and Sovetachen sections in Transcaucasia (Armenia) were organized during the 1984 IGC in Moscow. It was a unique opportunity for participants to collect samples of these remote sections (Fig. 20).



Figure 20 – Left, E.T. Tozer between Y. Arkhipov and A. Dagis, field trip leaders, at the lower *Otoceras* beds, Setorym Creek (E Siberia); right: Field trip participants in front of the snowy Ararat summit (Armenia), Zun-Yi Yang is on the right and G. Kotlyar in the middle.

Progress of IGCP Projects*

No. 4/106 TRIASSIC OF THE TETHYS REALM

H. Zapfe, Erdwissenschaftliche Kommissionen der Österreichischen Akademie d. Wissenschaften, A-1010 Wien, Postgasse 7, Austria.

<u>Description</u>. Revision of the biostratigraphy in the Tethyan Triassic. Redefinition of classical stages and substages, their emendation for a worldwide application. Comparison of the scheme with other major faunal realms (North America, Eurasia, Indo-Pacific) alming at a global Triassic time scale. Once this stage has been achieved, other important topics are to be studied, such as palaeogeography, faunal provinces and palaeomagnetism during Triassic time. An original study was made of the Upper Triassic and during recent years the work has expanded more and more to cover the entire Triassic, and the Project has been expanded to Asia in order to include the Triassic of the Himalayas, China and the Far East.

NO. 203 - PERMO-TRIASSIC EVENTS OF EASTERN TETHYS AND THEIR INTERCONTINENTAL CORRELATION

Yang Zunyi, Beijing Graduate School, Wuhan College of Geology, Beijing 100083, China.

J.M. Dickins, BMR, P.O.B. 378, Canberra City, ACT 2601, Australia.

W.C. Sweet, Department of Geology, Ohio State University, Ohio 43210, USA.

Description: The main objectives of Project 203 include: (1) a detailed study of the Permo-Triassic strata in the Eastern Tethys region and their correlation with equivalent stratigraphic units in Gondwana and Circum-Pacific realms,



Figure 21 – left, IGC Project. 4. 106 and 203 short description; right: Zun-Yi Yang and Norman Newell at the PT boundary, Shangsi section, March 1984.

IGCP Project 203: "Permo-Triassic events in the Eastern Tethys", 1984-1989

In 1984 began the first Chinese IGCP project 203 (Fig. 21, left) proposed and leaded by Prof. Zun-Yi Yang from Beijing Graduate School, on "Permo-Triassic events in the Eastern Tethys".

With this project, the large opening of China to Western scientists became operative and a successful conference took place in Beijing, March 1984, followed by the fieldtrip to Shangsi section in northern Sichuan province (Fig. 21, right). This is the start of an active collaboration on Permian-Triassic boundary studies with Chinese colleagues.

In July 1986 at Brescia (northern Italy), our Italian colleagues organized a field conference on Permian and Permian-Triassic boundary of Western Tethys sponsored by the IGCP Project 203. The Proceedings of the Field Conference were published at the "Memorie della Societa Geologica Italiana" (Cassinis, ed., 1988, Fig. 22, left) with new views and aspects on the Permian-Triassic boundary. An invitation was addressed to the members of the International Sub-commission on Triassic stratigraphy (SST) and to the members of the Permian-Triassic Boundary working group to participate to a post conference field trip (Fig. 22, right) on Permian and Triassic of Western and Southwestern Turkey (Visscher, 1986) with the Kemer Gorge and the Curuk Dagh PTB sections and a guidebook was published (Marcoux, ed., 1986).

IGCP Project 199: "Rare events in Geology", 1984

The IGCP project 199 on "Rare events in Geology" started in 1984 under the leadership of Prof. Ken Hsu (ETH Zurich, Switzerland).

Among "Rare events", the Permian-Triassic largest extinction was the subject of a Conference in Beijing in 1987, followed by a visit to the Nanjing Geological Institute (Fig. 23) and by an international field workshop to the Meishan section, with the first opportunity to sample the PT Boundary in the Meishan quarries.

Boosted during the 1980's by the IGCP Project 199, Bill Holser (Fig 24), from Oregon University in Eugene (USA), directed geochemical studies towards Permian and Triassic successions. With him and Mordekai Magaritz from Rehovot University (Israel), we started in 1985, the first database on stable carbon isotopes of more than twenty Permian-Triassic boundary sites from Southern Alps to South China, from studied sample collections stored in Lausanne Geological Museum. We published (Baud et al., 1989) twelve well dated sections with C isotope curve from bulk rock marine carbonate samples showing, for the first time, the global shift (Fig. 24 right) at the boundary: a new tool for marine Permian-Triassic boundary correlation was open and two main PTB candidate sections, Meishan and Shangsi were concerned and illustrated.

With Viorel Atudorei and Halil Sharp (Lausanne University), we provided a detailed C isotope data on a third PTB candidate, the Guryul Ravine section, Indian Kashmir (Baud et al., 1996). Some years later in Lausanne, Viorel Atudorei was the first to publish a complete C isotope curve of the marine lower Triassic, showing large positive anomalies within the well dated marine succession of the Spiti Valley (Atudorei, 1999) and, Sylvain Richoz (2004) first reported very detailed C isotope curves from PTB sections in Turkey, Iran and Oman.

IGCP Project 272: Late Paleozoic and Early Mesozoic Circum-Pacific Bio-Geological Events, 1988-1992

Succeeding to the IGCP Project 203, the Project 272: Late Paleozoic and Early Mesozoic Circum-Pacific Bio-Geological Events was leaded by J.M. Dickins from Australian Geological Survey.

Just after the Kyoto IGC (end of August 1992), Profs. Yuri Zakharov and Galina Kotlyar organized with my help the first Permian-Triassic conference (Fig. 25, right) and field workshop in Vladivostok (Russia). It was sponsored by the IGCP Project 272 and supported by the STS. The participants get the opportunity to visit four main Permian-Triassic sections of this Far-East Russian area, just open to foreigners' scientists. A first report was published by Dickins (1993). The proceedings of the field workshop went out in Lausanne (Baud et al., ed., 1997, Fig. 25, left) and the main Project 272 results were published by Dickins et al. (1997).



LATE PERMIAN AND TRIASSIC WORKSHOP MEETING IN WESTERN TURKEY (July 12-23, 1986) - REPORT

Four institutions joined their efforts to organize a workshop and field meeting on the Late Permian and Triassic of western Turkey: (1) the Technical University of Istanbul (with Prof.Dr. K. Kafali, rector of the University), (2) the Faculty of Mines, Istanbul (with Prof.Dr. E. Yüzer, Dean of the Faculty, Dr. A.M.C. Sengör, Dr. N. Ozgül, Dr. R. Akkök), (3) the "Laboratoire de Tectonique" of Paris VII University (with Dr. J. Marcoux) and (4) the Subcommission on Triassic Stratigraphy (with Dr. A. Baud. Vice Chairman).

The meeting was attended by 22 specialists from 11 countries and 4 continents; it was small enough to enable warm and durable contact between the participants.

Figure 22 - Left, cover of the Brescia Conference Proceedings; right: report on the PT West Turkey field workshop, 1986.

NO. 199 - RARE EVENTS IN GEOLOGY

K.J. Hsü, Geological Institute, ETH-Zentrum, Sonneggstrasse 5, 8006 Zürich, Switzerland. The Second International Meeting of the project was held in Beijing, China, 3-5 March 1987, with participants coming coming from ten countries: Austria, Canada, People's Republic of China, France, Italy, Spain, Switzerland, the United Kingdom, U.S.A. and USSR. Some 30 papers were presented fol-lowed by two sessions of general discussion on various aspects of rare events and mass extinctions. Chinese geologists reported that biological mass extinctions, stable isotope perturbations, geochemical anomalies and microspherules have been found not only across the Cretaceous/Tertiary boundary, but also the Permian/Triassic, Devonian/Carboniferous, Frasnian/Famennian, Ordovician/Silurian, and Precambrian/Cambrian boundaries in China. However, much work is needed to confirm the preliminary results. Professor K.J. Hsü gave a paper on the use of carbon isotope as a monitor for biomass. D. McLaren emphasized that the mass extinctions in geological history are not only distinguished by the disappearance of many taxa, but more impressive is the nearly total destruction of the biomass by the boundary event. Many participants took part in a five-day post-conference field trip to Changxing in Zheijang province.



Figure 23 – Left, IGC Project 199 Beijing meeting report; right: the participants of the "rare events" meeting on the steps of the Nanjing Geological Institute, March 1987; in black and in middle front, Ken Hsü, leader of IGC Project 199.Institute, March 1987; in black and in middle front, Ken Hsü, leader of IGC Project 199.Institute, March 1987; in black and in middle front, Ken Hsü, leader of IGC Project 199.



William T. Holser



Figure 24 – Left, W.T. Holser photo; middle: cover of the 1989 Baud, Holser and Magaritz Carbon isotope studies paper; right: the Carbon isotope global shift as shown in the last figure.



Figure 25 – Left, cover of the Vladivostok Permian-Triassic conference Proceedings; right: round table on IGCP Project 272 during the Vladivostok Permian-Triassic conference with Y. Zakharov, G. Kotlyar, J. M. Dickins, chairman, and the author.

IGCP project 359: Correlation of Tethyan, Circum-Pacific and marginal Gondwanan Permo-Triassic (1993-1998)

Led by Prof. Hongfu Yin (Fig. 26), this project, starting in 1993 and continued through 1997, gave to the participants the

opportunity to participate to specific international meetings with group discussion on the PTB and to visit Permian-Triassic sections in South China, in North Vietnam and in North Caucasus. For the PTB, one of the major achievements was the publication of the Yin's edited book (1996) on the four PTB candidates (Fig. 26, right).



Figure 26 - Cover of Yin's edited book on the four PTB candidates (1985)

THE INTERNATIONAL SUBCOMMISSION ON TRIASSIC STRATIGRAPHY (SST) AND THE PERMIAN-TRIASSIC BOUNDARY WORKING GROUP (PTBWG)

The history of the Subcommission on Triassic stratigraphy up to 1984 has been well described by Tozer (1985). Albertiana, the STS newsletter and platform of discussion publishing preliminary notes, played a very important role in promoting agreement among scientists about criteria fixing the Permian-Triassic boundary, as did Permophiles, the Subcommission on Permian Stratigraphy newsletter (Fig. 27).

Divergent opinions to define the boundary: ammonoid versus conodonte

During the 1980's two main camps rose out, the ammonoid workers with Tim Tozer and the new conodont worker group supported by Prof. H. Yin, former student of Prof. Z.-Y. Yang.

In 1984, Tim Tozer asked the PTBWG members about how to define the base of the Triassic. The great majority answered the *Otoceras* beds. But Normam Newell expressed his choice with the base of the Dienerian as published in his 1988 paper. Galina Kotlyar (1990) was in favor of the *Ophiceras* zone and correlative *Isarcica* conodont zone as Walter Sweet (1988, 1992). The Tozer's proposal was supported by the papers of A. Dagis (1988) and of Nakazawa (1992, 1993).

The conodont specialists get their defender with the H. Yin proposals (Yin, 1988): the first occurrence of the *Hineodus parvus* conodont to define the base of the Triassic, -and the top of boundary clay of Chinese sections as boundary stratotype.

The Lausanne Conference (1991)

Elected STS Chairman during the Washington ICS in 1989, I prepared with my colleagues Jean Guex (Lausanne), Maurizio Gaetani (Milano), Jean Marcoux (Paris) and Hans Rieber (Zurich) a Triassic Conference in Lausanne, Switzerland, for October 1991 with the participation of most of the Subcommission voting members. This conference was followed by a one day fieldtrip in the Triassic succession of the nearby Prealps (Fig. 28, right). The Triassic Conference proceedings were published 3 years later (Guex and Baud, eds., 1994, Fig. 28 left). Urged by the International Commission on Stratigraphy (ICS) we had, during this conference, to decide how to subdivide the Early Triassic period among four proposals (subdivision in 1, 2, 3 or 4 stages). After a long discussion, the majority of the



Figure 27 - Left, cover of Albertiana 17 with Ophiceras tibeticum; left: cover of Permophiles 31 with PTB view in Meishan quarry (up and right).



Figure 28 – Left, cover of the 1991 Triassic conference Proceedings (Guex and Baud, Eds., 1994); right: the Triassic ammonite specialists W. Weitschat, A. Dagis, H. Rieber and Y. Zakharov in the Saint-Triphon quarry (post-Triassic conference fieldtrip in Western Switzerland).

voting members proposed the two stage subdivision, named Induan and Olenekian. A detailed report was made by Gaetani (1992). I forwarded the choice to the International Commission on Stratigraphy (ICS) and the new stage names were officially adopted in 1992 during the International Geological Congress (IGC) in Kyoto.

The Permian-Triassic Boundary Working Group (PTBWG) reactivated

The next step of the STS was the setting up of stage boundary working groups on Global Stratigraphic Section and Point (GSSP). As the discussion at the PTBWG was, from 1984, deadlocked, I proposed to H. Yin to reactivate it. Early in 1993, a majority of the voting members elected Yin Hongfu as new chairman. Efficiently, he submitted four GSSP candidates for the boundary: the Meishan section (South China), the Shangsi section (West China), the Selong section (South Tibet) and the Guryul Ravine section (Kashmir, Northern India).

Very active, with the support of the Chinese geological community, he answered point by point to the 1988 Tozer's views, showing the main difficulties when using the ammonoid *Otoceras* to fix the boundary, and the great advantage to move to conodont study for the correlations (Yin, 1994). This was debated during the 1994 PTBWG meeting in Calgary. Yin Hongfu received also a strong support in a 30 pages paper leaded by H. Kozur, with A. Ramovs, C.Y. Wang and Y. Zakharov, to use *Hindeodus parvus* (Kozur & Pjatakova, 1976) for the PTB definition and to set up the Meishan quarry for the stratotype (Kozur et al., 1994).

In 1995, a majority of the working group approved the Meishan section proposal for Permian-Triassic GSSP. This was

forwarded to the STS members with the new Yin's edited book (Fig. 26) on the four candidates (1996).

Return to High Arctic lower Triassic stratotype

Invited by Benoit Beauchamp I started collaborative work on the Permian-Triassic transition on Ellesmere Island during the summer 1992. Two years later, with Charles Henderson, we sampled High Arctic localities with *Otoceras* for conodont studies, to solve the controversy about the lower *Otoceas* zone correlation. The results were published in the Proceedings of the Beijing IGC 1996 (Henderson and Baud, 1997, Fig. 29, left) showing for the first time that overlying Changxingian conodonts, the *Hindeodus parvus* species appears in the middle of the upper *Otoceras* zone (*Otoceras boreale*).

ACHIEVEMENTS: THE NEW GSSP IN THE MEISHAN QUARRY

At the International Geological Congress (IGC) in Beijing (1996), Yin Hingfu proposed the base of bed 27C at Meishan section D with the first occurrence of the conodont *Hindeodus parvus* for the base of the Triassic GSSP (Yin et al., 1996). But I made the following warning of the very condensed succession of the Permian-Triassic transition at the Meishan locality with a frequent facies changes just below the boundary (not recommended for a stratotype, Baud, 1996)). And the long range, with numerous morphotypes of the selected *H. parvus* conodont genus erected by Heinz Kozur on the finding of Pjatakova in Transcaucasia (Kozur and Pjatakova, 1976) can

Proc. 30^a Int'l. Geol. Congr., Vol. 11, pp. 143-152 Wang Naiwen and J. Remane (Eds) © VSP 1997 CODDEL ATION OF THE PERMIAN.TRIAS

CORRELATION OF THE PERMIAN-TRIASSIC BOUNDARY IN ARCTIC CANADA AND COMPARISON WITH MEISHAN, CHINA

CHARLES M. HENDERSON Department of Geology and Geophysics, University of Calgary, Calgary, Alberta, CANADA and AYMON BAUD Geological Museum, UNIL-BFSH 2, CH-1015, Lausanne, SWITZERLAND.

Geological Museum, UNIL-BESH 2, CH-1013, Lausanne, SWI12

ABSTRACT

Correlation of the Permian-Triassic boundary (P-T) is impeded, in part, by the lack of biostratigraphic correlations between tethyan and boreal settings and the presumed unconformity at many P-T sections in the world. Sequence stratigraphic and biostratigraphic data were collected from six sections of the basal part of the Blind Fiord Formation (Confederation Point Member) on northwestern Ellesmere and Axel Heiberg islands; sequence biostratigraphic data are presented in this paper for only the Griesbach Creek and South Otto Fiord sections. These data address some of these impediments, and allow comparisons to be made with the possible Global Stratotype Section and Point (GSSP) at Meishan, China.



Figure 29 – Left, first page of Henderson and Baud (1997); right: the Permian-Triassic team with the author between Charles Henderson (left) and Benoit Beauchamp at Otto fiord, Ellesmere Island, 1994.

likely brought problems in the determination and consequently in the correlation. Also a detailed discussion on the diachronic *H. parvus* conodont first occurrence was published in Orchard and Krystyn (1998).

Following a positive vote in 1999 by the STS voting members on the Meishan GSSP, the proposal was sent by Maurizio Gaetani, new STS Chairman to the ICS and ratified by the ICS general Assembly, during the Rio de Janeiro IGC in 2000. And the new GSSP final report appears in Episode (Yin et al, 2001). Officially, the Meishan section became the new Erathem boundary with the base of the Triassic Period fixed in the Meishan quarry (Fig. 30, left).

This was unveiled in this quarry by an opening Ceremony of the GSSP Monument, August 11, 2001 (Fig. 30, right), during the International Symposium on "The Global Stratotype of the Permian-Triassic Boundary and the Paleozoic- Mesozoic Events" held in Changsing City and leaded by H. Yin.

Thus, after a long time, a successful achievement were done and now the Meishan quarry houses a huge Geopark including an Earth History Museum with giant sculptures and educational exhibitions.



Figure 30 – Left, cover Albertiana 26; right the red ribbon cut ceremony of the Erathem boundary monument, in the Meishan quarry, August 11, 2001 with Charles Henderson (left) and the author.

SOME CONCLUSIONS

During more than a Century were a rash of stage names for the underlying late Permian as Saxonian, Araxian, Chhidruan, Amarassian, Tatarian, Djoulfian / Dorashamian, Changhsingian. Same of plenty stage names happen to the following lower Triassic with successively Buntsandstein, Scythian, Brahamian, Gangetian, Werfenian, Induan or Griesbachian. Finally, the ICS adopted the Changhsingian for the latest Permian stage and the Induan for the Earliest Triassic stage.

With the adoption of the Permian-Triassic GSSP in the Meishan quarry by the ICS and the IUGS, started a boom in Permian-Triassic transition studies, special volumes and publications. Large opening of South China to foreign scientists, new IGCP projects on extinction and recovery were encouraging teams of young researchers in paleontology, biostratigraphy, magnetostratigraphy, geochemistry and absolute dating, with the coming out of hundreds of new papers each year.

But now, with the discover of extended latest Permian sections in South China and extended earliest Triassic sections in Artic areas, the actual Meishan highly condensed GSSP section shows his borderline. As an extremely short duration of the Induan have been established recently (see Burgess et al., 2014, Ovtcharova et al., 2007), sure a new subdivision of the lower Triassic will be proposed and a new locality for the GSSP will come in the next future.

ACKNOWLEDGEMENTS

Thanks to Prof. Christopher A. McRoberts for his advice and help to improve a first version of this illustrated historical study.

REFERENCES

- Abich, O.W. H. 1878. Geologische Forschungen in den kaukasischen Landern, I: Eine Bergkalkfauna aus der Araxesenge bei Djoulfa in Amenien. A. Hölder, Wien, 126 pp.
- Atudorei, V. 1999. Constraints on the upper Permian to upper Triassic marine carbon isotope curve. Case studies from the Tethys. (PhD Dissertation) Lausanne University, 155 pp.
- Baud, A. 1996. The Permian-Triassic Boundary: recent developments, discussion and proposals: Albertiana, 18: 6–9.
- Baud, A. 2008. Correlation of Upper Permian localities in the Kuh-e-Ali Bashi area, NW Iran: Old collections, old and new data. Permophiles, 52: 6–11.
- Baud, A., Atudorei, V. & Sharp, Z. 1996, Late Permian and Early Triassic Evolution of the Northern Indian Margin: Carbon Isotope and Sequence Stratigraphy. Geodinamica Acta, 9: 57–77.
- Baud, A., Holser, W.T. & Magaritz, M. 1989. Permian-Triassic of the Tethys: Carbon isotope studies. Geologische Rundschau, 78(2): 649–677.
- Baud, A., Popova, I., Dickins, J.M., Lucas, S. & Zakharov, Y. (eds.). 1997. Late Paleozoic and Early Mesozoic Circum-Pacific Events: Biostratigraphy, Tectonic and Ore Deposits of

Primoryie (Far East Russia), Mémoire de Géologie (Lausanne), 30: 202 pp.

- Bonnet, P. 1919. Sur les relations entre les couches a *Otoceras* de l'Armenie (Transcaucasie meridionale) et celles de l'Himalaya. C.R. Acad . Sc. Paris, 169: 288–291.
- Burgess, S.D., Bowring, S. & Shen, S.-Z. 2014. High-precision timeline for Earth's most severe extinction. Proceedings of the National Academy of Sciences, 111: 3316–3322.
- Cassinis, G. (ed.). 1988. Proceedings of the Field Conference on Permian and Permian-Triassic boundary in the South-Alpine segment of the Western Tethys. Memorie della Societá Geologica Italiana, 34: 366 pp.
- Dagys, A.S. & Dagys, A.A. 1986. Biostratigraphy of the Lowermost Triassic and the boundary between Paleozoic and Mesozoic. Memorie della Societá Geologica Italiana, 34: 313–320.
- Dickins, J.M. 1993. IGCP Project 272. Late Paleozoic and Early Mesozoic Circum-Pacific Events. Working Group Meeting, Vladivostok, 6-13 September 1992. Albertiana, 11: 2–4.
- Dickins, J.M., Yang Zunvi, Yin Hongfu, Lucas, S.G. & Acharrya, S.K. (eds.). 1997. Late Palaeozoic and Early Mesozoic Circum-Pacific Events and their Global Correlation. World and Regional Geology Series, 10, Cambridge University Press, 245 pp.
- Diener, C. 1897. The Cephalopoda of the Lower Trias. Palaeontologia Indica, ser. 15, Himalayan Fossils 2: 181 pp.
- Diener, C. 1912. The Trias of the Himalayas. Memoirs of the Geological Survey of India, 36(3): 176 pp.
- Frech, F. 1902. Die Dyas (Schluss). Lethaea geognostica 1, Lethaea palaeozoica, 2(4): 453–664. Stuttgart.
- Frech, F. & Arthaber, G. V. 1900. Über das Paläozoicum in Hocharmenien und Persien: Mit einem Anhang über die Kreide von Sirab in Persien. Beiträge zur Paläontologie Österreich-Ungarns und des Orient, 12: 161–308.
- Gaetani, M. 1992. Report on the Symposium on Triassic Stratigraphy, Lausanne 1991, and report on the vote of the Subcommission. Albertiana, 10: 6–10.
- Ghaderi, A., Ashouri, A. R., Kozur, H. & Korn, D. 2013. Age assignment of section 4 of Teichert et al. (1973) at Ali Bashi Mountains (Julfa, NW Iran). Permophiles 58: 36–39.
- Griesbach, C.L. 1880. Paleontological notes on the Lower Trias of the Himalayas. Records of the Geological Survey of India, 13: 94–113.
- Griesbach, C.L. 1891. Geology of the Central Himalayas, Memoirs of the Geological Survey of India, XXIII: 232 pp.
- Guex, J. & Baud, A. (eds.). 1994. Recent developments on Triassic Stratigraphy Mémoires de Géologie, Lausanne, Switzerland, 22: 182 pp.
- Heller, F., Lowrie, W., Huamei, L. & Junda, W. 1988. Magnetostratigraphy of the Permo-Triassic boundary section at Shangsi (Guangyuan, Sichuan Province, China). Earth and Planetary Science Letters, 88/3: 348–356.
- Ogg, J.G. & Steiner, M.B. 1991. Early Triassic magnetic polarity time scale—integration of magnetostratigraphy, ammonite zonation and sequence stratigraphy from stratotype sections (Canadian Arctic Archipelago). Earth and Planetary Science Letters, 107/1: 69–89.

- Henderson, C. & Baud, A. 1997. Correlation of the Permian-Triassic boundary in Arctic Canada and comparaison with Meishan, China. *In*, Naiwen, W. & Remane, J. (eds.), Stratigraphy, 11, Proceedings of the 30th IGC: Beijing, VSP: 143–152.
- Henderson, C.M., Mei, S.L., Shen, S.Z. & Wardlaw, B.R. 2008. Resolution of the reported Upper Permian conodont occurrences from northwestern Iran. Permophiles, 51: 2–9.
- Iranian-Japanese Research Group, 1981. The Permian and the Lower Triassic Systems in Abadeh Region, Central Iran. Memoirs of the Faculty of Sclence, Kyoto University, Series of Geology & Mineralogy, 47: 61–133.
- Koch, L. 1931. Carboniferous and Triassic stratigraphy of East Greenland. Meddelelser om Grønland 83(2): 1–100.
- Kotlyar, G.V. 1991. Permian-Triassic boundary in Tethys and the Pacific Belt and its correlation. Proceedings of the International Symposium on Shallow Tethys, 3: 387–391.
- Kozur, H. & Pjatakova, M. 1976. Die conodontenart Anchignathodus parvus n. sp., eine wichtige Leitform der basalen Trias. Koninklijke Nederlandse Akademie van Wetenschappen Proceedings, Series B, 79(2): 123–127.
- Kozur, H.W., Ramovs, A., Wang, Cheng-Yuan & Zakharov, Y.D., 1994/95. The importance of *Hindeodus parvus* (Conodonta) for the definition of the Permian-Triassic boundary and evaluation of the proposed sections for a global stratotype section and point (GSSP) for the base of the Triassic. Geologija, Ljubljana, 37–38: 173–213.
- Kraft, A. von. 1901. Ueber das Permische Alter der *Otoceras*-Stufe des Himalaya. Zentralblatt fur Geologie und Palaontologie, 1901: 275–279.
- Kummel, B. 1966. The lower Triassic formations of the Salt Range and Trans-Indus Ranges, West Pakistan. Bulletin of the Museum of Comparative Zoology, 134(10): 361–429.
- Kummel, B. 1972. The Lower Triassic (Scythian) ammonoid *Otoceras*. Bulletin of the Museum of Comparative Zoology, 143(6): 365–417.
- Kummel, B. & Teichert, C. (eds.). 1970. Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan. Department of Geology, University of Kansas Special Publication, 44. University of Kansas Press, Lawrence. 330 pp.
- Kummel, B. & Teichert C. 1970. Stratigraphy and Paleontology of the Permian-Triassic boundary beds, Salt Range and Trans-Indus Range, West Pakistan. *In*, Kummel, B. & Teichert, C. (eds.), Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan. Department of Geology, University of Kansas Special Publication, 44: 1–109. University of Kansas Press, Lawrence.
- Logan, A., & Hills, L.V. (eds.) 1973. The Permian and Triassic Systems and their mutual boundary. Canadian Society of Petroleum Geologists, Memoir 2. Canadian Society of Petroleum Geologists Calgary. 766 p.
- Mojsisovics, E. von, 1879. Vorläufige kurze Übersicht der Ammoniten-Gattungen der mediterranen und juvavischen Trias. Verhandl. Reichesanst. Wien, Jg. 1879, 133–143.
- Mojsisovics, E. von, Waagen, W.H. & Diener, C. 1895. Entwurf einer Gliederung der pelagischen Sediments des Trias-Systems. Akademie Wissenschaft Wien, Mathematische-

naturwissenschaftliche Klasse Sitzungsberichte, 104: 1279–1302.

- Matsuda, T. 1981. Early Triassic conodonts from Kashmir, India, Pt. 1, *Hindeodus* and *Isarcicella*, Journal of Geosciences, Osaka City University, 24(3): 75–108.
- Matsuda, T. 1982. Early Triassic conodonts from Kashmir, India, Pt. 2, *Neospathodus* 1. Journal of Geosciences, Osaka City University, 25(6): 87–103.
- Matsuda, T. 1983. Early Triassic conodonts from Kashmir, India, Pt. 3, *Neospathodus* 2. Journal of Geosciences, Osaka City University,, 26(4): 87–110.
- Matsuda, T. 1984. Early Triassic conodonts from Kashmir, India, Pt. 4, *Gondolella* and *Platyvillosus*. Journal of Geosciences, Osaka City University, 27(4): 114–119.
- Nakazawa, K. 1992. The Permian-Triassic boundary. Albertiana, 10: 23–30.
- Nakazawa, K. 1993. Stratigraphy of the Permian-Triassic transition and the Paleozoic/Mesozoic boundary: Bulletin of the Geological Survey of Japan, 44: 425–445.
- Nakazawa, K., Ishii, K., Kato, M., Okimura, Y., Nakamura, K. & Haralambous, D. 1975. Upper Permian fossils from island of Salamis, Greece. Memoirs of the Faculty of Science. Kyoto University, Series of Geology and Mineralogy, 41: 21–44.
- Nakazawa, K., Kapoor, H. M., Ishii, K., Bando, Y., Okimura, Y. & Tokuoka, T. 1975. The Upper Permian and Lower Triassic in Kashmir, India. Memoirs of the Faculty of Science. Kyoto University, Series of Geology and Mineralogy, 47: 1–106.
- Nakazawa, K. & Kapoor, H.M.,(eds.) 1981. The Upper Permian and Lower Triassic Faunas of Kashmir. Palaeontologica Indica New Series, 46: 1–191.
- Nakazawa, K., Suzuki, H., Kumon, F. & Winsnes, T.S. 1990. Scientific results of the Japanese Geological Expedition to Svalbard 1986. *In*, Tatsumi, T. (ed.), The Japanese Scientific Expedition to Svalbard 1983–1988. Kyoikusha Press, Tokyo.: 179–214.
- Newell, N.D. 1986. The Paleozoic/ Mesozoic erathem boundary: Memorie della Societa Geologica Italiana, 34: 303–312.
- Noetling, F. 1905. Die Asiatische Trias. Lethaea geognostica 2, Lethaea mesozoica, Stuttgart 1 (3): 107–221.
- Ogg, J. G. & Steiner, M.B. 1991. Early Triassic magnetic polarity time scale—integration of magnetostratigraphy, ammonite zonation and sequence stratigraphy from stratotype sections (Canadian Arctic Archipelago). Earth and Planetary Science Letters, 107(1): 69–89.
- Orchard, M.J. & Krystyn, L. 1998. Conodonts of the lowermost Triassic of Spiti, and new zonation based on *Neogondolella* successions. Rivista Italiana di Paleontologia e Stratigrafia, 104(3): 341–367.
- Ovtcharova M., Bucher, H., Schaltegger, U., Galfetti, T., Brayard, A. & Guex, J. 2006. New Early to Middle Triassic U–Pb ages from South China: calibration with ammonoid biochronozones and implications for the timing of the Triassic biotic recovery. Earth Planetary Science Letters, 243(3-4): 463–475.
- Pakistanese-Japanese Research Group (PJRG). 1981. Stratigraphy and correlations of the marine Permian-Lower Triassic in the Surghar Range and Salt Range, Pakistan. Kyoto University

Press, 25 pp.

- Pakistanese-Japanese Research Group (PJRG). 1985. Permian and Triassic Systems in the Salt Range and Surghar Range, Pakistan. *In*, Nakazawa K.D. & Dickins J.M. (eds.), The Tethys. Tokai University Press, Tokyo.: 221–312.
- Ruzhentsev, V.E. & Sarycheva, T.G. 1965. Evolution and change in the marine organisms at the boundary between Paleozoic and Mesozoic. Transactions, Academy of Sciences of the USSR, Paleontological Institute, 108: 1-431.
- Schindewolf, O.H. 1954. Über die Faunenwende vom Palaozoikum zum Mesozoikum. Deutsch. Geol. Ges., Zeitschr., 105: 154–183.
- Silberling, N.J. & Tozer, E.T. 1968. Biostratigraphic classification of the marine Triassic in North America. Geological Society of America Special Paper, 110: 1–63.
- Spath, L.F. 1934. Part 4: The ammonoidea of the Trias, catalogue of the fossil cephalopoda in the British Museum (Natural History). The Trustees of the British Museum, London. 521 pp.
- Stepanov, D.L., Golshani, F. & Stöcklin, J. 1969. Upper Permian and Permian–Triassic boundary in North Iran. Report Geological Survey of Iran, 12: 1–72.
- Sweet, W.C. 1970a. Permian and Triassic conodonts from a section at Guryul Ravine, Vihi district, Kashmir. Paleontological Contributions, University of Kansas, 49: 1–10.
- Sweet, W.C. 1970b. Uppermost Permian and Lower Triassic conodonts of the Salt Range and Trans-Indus Ranges, West Pakistan. *In*, Kummel, B. & Teichert, C. (eds.), Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan. Department of Geology, University of Kansas Special Publication, 44: 207–275. University of Kansas Press, Lawrence.
- Sweet, W.C. 1976. Appendix. Conodonts from the Permian-Triassic boundary beds at Kap Stosch, East Greenland: Meddelelser om Grønland, 197: 51–54.
- Sweet, W.C. 1992. A conodont-based high-resolution biostratigraphy for the Permo-Triassic boundary interval. *In*, Sweet, W.C., Yang, Z., Dickins, J.M. & Yin, H. (eds.), Permo-Triassic events in the eastern Tethys: stratigraphy, classification, and relations with the western Tethys. Cambridge University Press, 120-133.
- Teichert, C., Kummel, B. & Kapoor, H.M. 1970. Mixed Permian-Triassic fauna, Guryul Ravine, Kashmir. Science, 167: 174–175.
- Teichert, C. & Kummel, B. 1972. Permian-Triassic boundary in the Kap Stosch area, East Greenland. *In*, Logan, A. & Hills, L.V. (eds.) The Permian and Triassic Systems and their mutual boundary. Canadian Society of Petroleum Geologists, Memoir 2: 269–285.
- Teichert, C., Kummel, B. & Sweet, W. 1973. Permian-Triassic strata, Kuh-e Ali Bashi, Northwestern Iran. Bulletin, Museum of Comparative Zoolology, 145: 359–472.

- Teichert, C. & Kummel, B. 1976. Permian-Triassic boundary in the Kap Stosch area, East Greenland. Meddelelser om Grønland, 197: 1–50.
- Tozer, E.T. 1965. Lower Triassic stages and ammonoid zones of Arctic Canada. Geological Survey of Canada Paper, 6–12: 1–45.
- Tozer, E.T. 1969. Xenodiscacean ammonoids and their bearing on the discrimination of the Permo-Triassic boundary. Geological Magazine, 106 (4): 348–361.
- Tozer, E.T. 1984. The Trias and its ammonoids: the evolution of a time scale. Geological Survey of Canada Miscellaneous Report, 35: 1–171.
- Tozer, E.T. 1985. Subcommission on Triassic Stratigraphy (STS): History 1968–1984. Albertiana, 3: 3–6.
- Tozer, E.T. 1986. Definition of the Permian-Triassic boundary: the question of the age of the *Otoceras* beds. Memorie della Societa Geologica Italiana, 36: 291–302.
- Tozer, E.T. 1988. Towards a definition of the Permian–Triassic boundary. Episodes, 11/3: 251–255.
- Trümpy, R. 1960. Über die Perm-Trias-Grenze in Ostgrönland und über die Problematik stratigraphischer Grenzen. Geologische Rundschau, 49: 97–103.
- Trümpy, R. 1969. Lower Triassic ammonoids from Jameson Land (East Greenland). Meddelelser om Grønland, 168: 77–116.
- Visscher H. 1986. Report 1986: Late Permian and Triassic workshop meeting in Western Turkey, July 12–23 1986. Albertiana, 5: 7–8.
- Waagen, W. 1895. Salt-Range fossils. Vol 2: fossils from the *Ceratite* Formation. Palaeontologia Indica, 13: 323 pp.
- Unesdoc at Unesco.org. 1972-2004. Geological Correlation Progress Report of the International Geological Correlation Programme (IGCP).
- Yin H.F., Yang F.Q., Zhang K.X. & Yang W.P. 1986. A proposal to the biostratigraphic criterion of the Permian-Triassic boundary. Memorie della Societa Geologica Italiana, 36: 329–344.
- Yin, H. 1996. The Paleozoic-Mesozoic Boundary. Candidates of the Global Stratotype Section and Point of the Permian-Triassic Boundary. *In*, Long, X. & Ding, M. (eds.), NSFC Project. China University of Geosciences Press, Wuhan, 137 pp.
- Yin, H. 1996. Recommendation of the Meishan section as the global stratotype and point (GSSP) for the Permian-Triassic Boundary (PTB), 30th International Geological Congress, Beijing, Abstract book, 57.
- Yin, H. & Tong, J. 2001. The Global Stratotype of the Permian-Triassic Boundary and the Paleozoic-Mesozoic Events, Changxing, China, August 10-13, 2001. Albertiana, 26: 34–41.
- Yin, H., Zhang, K., Tong, J., Yang, Z. & Wu, S. 2001. The Global Stratotype Section and Point (GSSP) of the Permian-Triassic Boundary. Episodes, 24: 102–114.

Research Article

MARINE *LARIOSAURUS* (SAUROPTERYGIA) RECORDS FROM THE MIDDLE TRIASSIC OF THE GERMANIC BASIN – STRATIGRAPHIC AND PALEOBIOGEOGRAPHIC IMPORTANCE FOR TETHYAN AND GERMANIC BASIN CORRELATION

Cajus G. Diedrich

PaleoLogic, Research Institute, Petra Bezruce 96, CZ-26751 Zdice, Czech Republic. Email: cdiedri@gmx.net

Abstract – The stratigraphically oldest *Lariosaurus* skeleton in Europe is of *L. teutonicus* (Diedrich, 1996) from the shallow marine lagoon deposits of the Lower Muschelkalk (Bithynian) of Borgholzhausen (NW Germany) in the Germanic Basin, possibly being synonymous with the (?)Pelsonian Chinese *L. xingyiensis* (Rieppel et al., 2003). Younger remains of *L. buzzii* Tschanz, 1989 were collected as isolated skulls and postcrania in the German Upper Muschelkalk (Illyrian/Fassanian) sites Bad Sulza and Bayreuth, mainly in the oolithic or shallow marine facies. *Lariosaurus buzzii* is also recorded in a few younger (Fassanian/Longobardian boundary) northern Tethys black-shale-lagoon deposits of the Swiss/Italian locality Monte San Giorgio by complete skeletons. The species *L. balsami* Curioni, 1841 seems to represent the stratigraphically youngest records in the late Longobardian of the northwestern Tethys (Austria, Italy, Switzerland). *Lariosaurus* is known from a complete Middle Triassic European record with possibly only three valid species and was distributed along the Pangaean coasts between Europe and Asia in lagoon and shallow marine paleoenvironments, where they probably smaller hunted fish and juvenile pachypleurosaurs.

INTRODUCTION

The Middle Triassic marine sauropterygian *Lariosaurus*, a close relative of the pachypleurosaurs and nothosaurs (Tschanz 1989), was thought to be present only in the Tethys Ocean and its coasts (Rieppel, 2000) and was never recorded or recognized from the Germanic Basin.

The remains of the Lower Muschelkalk-aged Chinese *Lariosaurus xingyiensis* (Rieppel et al., 2003), for which the exact stratigraphic position remains unclear (possibly Pelsonian: cf. Motani et al., 2008) extended the paleobiogeography of this genus. However, it was not compared to earlier mentioned material from the Germanic Basin, and possibly it represents a Pelsonian aged *Lariosaurus* being synonymous with the few younger, Pelsonian-aged "*P. teutonicus*" (cf. Diedrich, 1996; herein = *Lariosaurus teutonicus*). From the Lower Muschelkalk (Pelsonian) of Germany the oldest record of *Lariosaurus* is revised and further described, which was published preliminarily without detailed descriptions or any figures as "*Paranothosaurus teutonicus*" (Diedrich, 1996).

Already, Meyer (1847-55) figured from the Upper Muschelkalk (Illyrian-Fassanian) a skull and an ischium as "*Nothosaurus*" from the Bindlach locality near Bayreuth. In the Upper Muschelkalk (late Illyrian-Fassanian), *Lariosaurus* published skulls were misidentified as "*Nothosaurus marchicus*" (Rieppel and Wild, 1996) from the Bad Sulza and Bindlach sites and have been recently identified together with postcranial bones as *Lariosaurus* cf. *buzzii* remains (Diedrich, 2012a, 2013a). A few younger and similar-aged specimens assigned to *L. buzzii* were published by Tschanz (1989), which is best known by skeletons from the black shales of the Swiss Monte San Giorgio (Furrer, 2003) and Italian Perledo localities (Rieppel, 2000; Fig. 2B).

The Lower Keuper Lariosaurus balsami skeletal remains were published from the Tethys Ocean from the Partnach-Plattenkalk of the Gailtaler Alpen by Curioni (1847) in the early Longobardian (Ladinian) aged black shale deposits of Austria, which material was discussed (Rieppel, 1994, 1998). New finds of this species were further recorded on the northwestern Tethys from the Pyrenees in Spain (Sanz, 1976), France (Mazin, 1985) and Italy (Ticli, 1984), with different-aged individual skeletons. L. lavizzarii Kuhn-Schnyder 1979 represents, based on Rieppel (1994), only a juvenile of L. balsami, of which one skeleton from Perledo is discussed by Peyer (1933-34) and figured (Fig. 2A). Possibly, L. valceresii (Tintori and Renesto, 1990), which is recently the stratigraphically youngest Lariosaurus record from the Meride Limestone (Stockar, 2010) of the Kalkschieferzone (late Longobardian, Ladinian) of Perledo, Italy (Deecke, 1886), is not validated as a distinct species. Clear identifications of the latter, Lower Keuper species are missing in the Germanic Basin, but might be expected in the Lower Keuper bonebed material of southern Germany after further studies.



Figure 1 – *A*, Global distribution of *Lariosaurus* in Europe and China (composed after Curioni, 1841; Sanz, 1976; Tschanz, 1989; Rieppel, 2000; Rieppel et al., 2003). *B*, Newly identified *Lariosaurus* sites in Germany. *C*, Biostratigraphy of the Germanic Basin *Lariosaurus* remains from Lower and Upper Muschelkalk layers (Middle Triassic) within different carbonate facies types (composed and modified from Diedrich, 2009, 2012b, 2013d).

MATERIAL AND METHODS

A disarticulated incomplete skeleton from the Lower Muschelkalk was excavated in 1995 by the author at Borgholzhausen (Diedrich, 1996) in Germany at a house construction site, which is in the collection of the Museum Natur und Umwelt Osnabrück (= MNUO). Single skulls and postcranial bones from the Upper Muschelkalk from Bad Sulza (Central Germany) (Diedrich, 2013d) are housed in the Naturkundemuseum Erfurt (= NME), the Saline and Heimatmuseum Bad Sulza (= SHMBS), and the Natural History Museum of the Humboldt-University Berlin (= MB). Skull and postcranial remains from Bayreuth (Diedrich, 2012a) are stored in the Senckenbergmuseum Frankfurt (= SMF) and Urweltmuseum Oberfranken (= UM-O). All available Middle Triassic Germanic Basin material was compared directly with the original skeleton from the University Munich and the cast of L. buzzii from Perledo (Figs. 2B-D), which is kept in the University Tübingen (= UT). Finally, all herein described German Lariosaurus sites were studied for their stratigraphy, facies/ paleoenvironments and vertebrate content, whereas several new studies on skeletons and isolated vertebrate material from the Germanic Basin (cf. Diedrich, 2009a, 2010, 2011a-b, 2012ab, 2013a-d) allowed identifications of Lariosaurus single bones within various German Triassic vertebrate museum collections.

SYSTEMATIC PALEONTOLOGY

Sauropterygia Owen, 1860 Nothosauridae Baur, 1889 Lariosauridae Lydekker, 1889 *Lariosaurus* Curioni, 1847 *Lariosaurus buzzii* Tschanz, 1989

Figs. 2, 3

Material. Two skulls and some postcranial remains from Bindlach and Bad Sulza sites in Germany, that include both pectoral and fore limb, pelvic and hindlimb and axial skeletal remains (Fig. 3).

Stratigraphy. Atavus zone, *atavus* bonebeds (oolite bonebed) of the lowermost Upper Muschelkalk (middle Illyrian) in Germany (Diedrich, 2012a, 2013d). *Lariosaurus buzzii* is known by skeletons from the Illyrian/Fassanian or Anisian/Ladinian boundary of Monte San Giorgio, Switzerland (Tschanz, 1989, Fig. 2B-E).

Localities. Isolated remains are from Bindlach (Bayreuther localities) and Bad Sulza, Germany (Diedrich, 2012a-b). Skeletons are described from Monte San Giorgio, Switzerland and Italy (Tschanz, 1989; Rieppel, 2000).

Description. This species is well described by skeleton records (cf. Tschanz, 1989).

Lariosaurus teutonicus (Diedrich, 1996)

Holotype material. Cranial and postcranial elements of a single disarticulated individual skeleton (coll. MNUO).

Stratigraphy. Serpianites antecedens/Balatonites balatonicus cephalopod ceratite biozone, Basal Upper Terebratula bed, Terebratula substage, Osnabrück Formation, middle Lower Muschelkalk, middle Pelsonian, lower Anisian, Middle Triassic.

Locality. Northern corner of the private house grounding of Family Penner, Riesberg in Borgholzhausen, Middle Teutoburg Forst mountain chain, NW-Germany (coordinates Long. 3453,195 and Lat. 5774,350).

Description and differential diagnosis of the skeleton. From the skull only the incomplete lower jaw is represented by the dentary, which was found in close contact to the distal parts, the splenials/angulars/praearticulars indicating a not fully grown or a very old individual (Figs. 4-5). The left side is 66 mm long and has two large anterior teeth, then a small tooth, another large one (fang tooth formula 5/1), and, following, five more 2-4 mm high small teeth. The 17 mm wide symphysis is synostritic, with 12 mm of elongation. It seems, each side has five larger alveolar grooves for the 8-12 mm long fang teeth. All teeth have coarse enamel striations. The alveoli are asymmetricaly toothed, whereas on the left side three typically longer (about 3-5 times higher than the other teeth) fang teeth are present. The dental angle is 24°. The splenial overlaps the dentary about 35 mm.

Pectoral girdle - The coracoids are 56 mm long, proximally 31 mm and distally 26 mm wide and are distally partly incomplete (right one) and proximally wide-flattened. One coracoid seems to be pathological on its margin. The 1.5 mm foramina coracoidea are very small. The coracoid angle measures 35°. Distally, the symphysis has only 50% of the margin, which is also found in stratigraphically younger species. The scapula shape fits much more the outline shape of *L. buzzii* (cf. Tschanz, 1989) rather then of *Nothosaurus (N. marchicus)*. The left, complete 36 mm long scapula is 20 mm in maximum width, and has the typical distinct long processus. Other pectoral bone elements (clavicula, interclavicula) are missing.

Fore limb - The left limb is represented by one incomplete humerus (proximally and distally, estimated 62 mm long), the ulna, radius, the intermedium, distale IV, pisiform, and one phalanx, which allow a reconstruction of the extremity similar to the younger Lariosaurus species. The humerus is typically strongly curved in this genus (see Fig. 2C), whereas the noncompressed cross section in the specimen from Germany changes from proximally long-round to oval to distally high-oval. The tuberculus deltoideus is quite large, also seen in other species (L. buzzii, Fig. 2C; Tschanz, 1989), whereas the sulcus epicondylaris is deep. The 33 mm long, 14 mm proximal and 13 mm distal and wide, flattened ulna is complete. The inner anterior lateral margin has a convex hump, the outer lateral margin is slightly concave. The ulna outline shape of L. teutonicus fits that of L. buzzii (Fig. 2C; Tschanz, 1989) best. The length/width index is 4.25. The slimmer and also flattened complete radius is 32 mm long, proximally 9 mm and distally 8 mm wide. The outer margin is irregular. The length/width index is 7.75. Also, this is in general shape and size similar to the radius of L. buzzii (Fig. 2C; Tschanz, 1989). The left carpalia are represented by the nearly 10 mm wide intermedium, 4 mm wide and quadratic distale IV,



Figure 2 – Marine sauropterygian *Larisosaurus* skeletons from the Middle Triassic of Europe. *A*, Original of a young small individual of *Larisosaurus balsami* Curioni, 1847 from the late Longobardian (Ladinian/Carnian boundary) of Perledo, Italy (SMF without no.), ventral view. *B*, Cast of a large older individual of the neotype of *Lariosaurus balsami* Curioni, 1847 (original figured in Kuhn-Schnyder, 1987) from the late Longobardian (Ladinian/Carnian boundary) of Perledo, Italy (UT without no.), dorsal, *C*, Fore limb, *D*, Hindlimb of *L. balsam*i Curioni, 1847 cast, *E*, Skeleton reconstruction after *Lariosaurus balsami* Curioni, 1847, see skeleton of Fig. 2B.

the 6 mm wide oval pisiforme, and one 4 mm wide phalanx.

Axial skeleton – Four dorsal vertebrae are represented, but the neural arches are fused with the centra. The closed sutura neurocentralis indicate an older individual, but not very old compared to the non-fused jaw sutures. The prae- and postzygapophyses are wide and flattened. The distance between the praezygapophyses is 13 mm. The total height of the vertebra is only 21 mm, whereas the centrum is 10 mm and the dorsal arch 11 mm in height. The cross-section of the neural canal is kidney-formed, with the convex part up. The dorsal vertebrae are similar again to those of *L. buzzii* (Fig. 2C; Tschanz, 1989), whereas they are not isolated and deformed in the skeleton, including the neural arch.

The ribs of the German skeleton specimen are represented by



four cervical (different positions), nine dorsal, and one caudal rib. The four 10–35 mm long cervical ribs are from anterior to posterior and have small caput costae. Also, the dorsal ribs are from anterior to posterior 36–62 mm long. Their rib head is a figure eight in the shape of the articulatar surface. The 32 mm long flattened caudal rib is from an anterior position. Each of the gastral ribs is composed of five elements. The v-shaped middle gastralia have in the middle and anterior a small spine, whereas the ribs end on both sides in a spine. The angle and length (60 mm to 55 mm) of the middle gastralia increases from caudal to cranial. On this, 32 mm long middle part, the middle gastralia (both sides needle-like) are attached, as are the about 34 mm long end gastralia (one end needle-like, the distal one round). All ribs and gastralia fit descriptions and the skeleton of *L. buzzii* (Fig. 2C; Tschanz, 1989).

Pelvic girdle - Only the left pubis and ilium and fragments of the ischium are represented. The pubis is complete. The distal end of the pubis of *Lariosaurus* has a typically sigmoidal margin (see *L. buzzii*, Tschanz, 1989), but is damaged and clearly not separated into two forked parts as are found in similar aged *Nothosaurus* (*N. marchicus*). The length is 56 mm long, whereas the distal articulation area is 19 mm wide, and the other part measures 28 mm. The foramen obturatum is not open to the margin and only 3 x 4 mm in oval size. In *Nothosaurus* (see *L. buzzii*, Tschanz, 1989) in shape and less wide and without an elongated processus, as in *Nothosaurus*, and is complete and 17 mm in height with a length of 23 mm. The facies pubica measures 14 mm, the facies ischiadica 12 mm. The left ilium is fragmentary with the proximal part preserved as a negative impression.

In conclusion, the isolated bone material from an old adult, but not senile animal, in estimated total length of 100-120 cm of *L. teutonicus* fits well to similar and a few larger sized *L. buzzii* (cf. Tschanz, 1989) and is distinct from the similar aged *N. marchicus* in many osteological ways, which have been analyzed already (see Rieppel, 2000). The osteological and taxonomically relevant differences between *L. teutonicus* and *L. buzzii* are mainly found in the shorter dentary symphyses, and different outlined form of the coracoid (compare Figs. 3 and 5). The ulna of *L. teutonicus* is less wide (probably a result of non compaction in the black shale skeletons, in general a problem). The *L. teutonicus* skeleton from Germany is stratigraphically much younger (middle Pelsonian) than the about two million years younger *L. buzzii* (Illyrian/Fassanian or Anisian/Ladinian boundary, cf. Fig. 6).

DISCUSSION

Previous authors have not been able recognize *Lariosaurus* in the Germanic Basin (Rieppel, 2000), which was in fact widely distributed in the north-western Tethys and the Germanic Basin in the Illyrian, and later in the Fassanian and Longobardian. The claims of Monte San Giorgio endemism (Rieppel, 2000), especially of *L. balsami*, must be revised. Using the new chronostratigraphy of Kozur and Bachmann (2008) and newest stratigraphic results for the Bayreuth and Bad Sulza localities (Diedrich, 2012a-b), such as the updated Triassic time scale (cf. Ogg et al., 2014), and new palaeogeographical maps for the Middle Triassic of Europe, including extensive and changing carbonate intertidal flats (Diedrich, 2009, 2011b, 2012a, b), *Lariosaurus* species occur in different strata and show different distributions changing within the transgression/regression of the Middle Triassic.

Pelsonian – "*Paranothosaurus teutonicus*" (Diedrich, 1996) must be attributed consequently to the genus *Lariosaurus* and represents the oldest record of this genus (Fig. 6), because the Chinese finds are not yet dated well. The facies is a platy dolomite limestone, representing a lagoon deposit (cf. Lukas, 1991; Diedrich, 1996, 2012a, 2013d).

Illyrian – Overlooked cranial and postcranial material (Fig. 3.1-16) comes from Bad Sulza from stratigraphic levels of the *atavus* zone of the oolithic very shallow marine to lagoonal facies, and this material was recently identified as *Lariosaurus* (Diedrich, 2013d). A skull was misidentified by Rieppel and Wild (1996) as "*N. marchicus*" and is reassigned here to *Lariosaurus* cf. *balsami* Curioni, 1847 because the skull shape, nasal, orbit and parietal foramina shapes and measurements match more closely to those of *Lariosaurus* (cf. Tschanz, 1989) than to those of a small *Nothosaurus*. *N. marchicus* has furthermore only been positively identified from the Bithynian to the early Illyrian ("Lower Muschelkalk") in the Germanic Basin (e.g. Rieppel, 2000).

With the subsequent trangression, *Nothosaurus* appeared as a more highly evolved and few larger form (= *N. juvenilis*) at the end of the Middle Muschelkalk (Edinger, 1921), as has recently been described from a large quantity of cranial and postcranial

[←] Figure 3 – Lariosaurus buzzii Tschanz, 1989 remains mainly from the 1-16 Oolite Bed (atavus biozone, Lower Bad Sulza Fm) of the basal Upper Muschelkalk (Illyrian) of Bad Sulza, Central Germany. 1, Skull from the Oolite Bed (NME no. 78.234), a. dorsal. b. reconstruction. 2, Lower jaw from the Oolite Bed (NME no. 78/233), a. ventral, b. ventral reconstruction, c. lateral. 3, Scapula from the Oolite Bed (NME no. 78/349), lateral. 4, Humerus from the Oolite Bed (MB no. R.455), lateral. 5, Ischium from the Oolite Bed (NME no. R.126), dorsal. 6, Femur from the Oolite Bed (SHMBS no. 45), lateral. 8, Middle dorsal costa from the Oolite Bed (NME no. R.453), lateral. 9, Anterior cervical vertebral neural arch (NME no. 78/242), a. lateral, b. dorsal, c. cranial. 10, Posterior cervical/anterior dorsal vertebral neural arch centrum (NME no. 78/241), a. cranial, b. dorsal, c. lateral. 11, Middle dorsal vertebral neural arch (NME no. 78/476), a. dorsal, c. caudal. 12, Middle dorsal vertebral centrum (NME no. 78/240a), a. cranial, b. dorsal, c. lateral. 13, Middle Dorsal vertebral centrum from the Oolite Bed (MB no. 145.1), a. cranial, b. dorsal, c. lateral. 14, Middle Dorsal vertebral centrum (NME no. 78/78/240), a. cranial, b. dorsal, c. lateral. 15, Posterior dorsal vertebral centrum (NME no. 78/240b), a. cranial, b. dorsal, c. lateral. 16, Anterior caudal vertebral centrum (NME no. 93/12), a. cranial, b. dorsal, c. Lateral. 17-23. Oolite Bed and Terebratulid tempestite beds (atavus to maximum ?compressus biozone, Bindlach Fm) of the basal Upper Muschelkalk (Illyrian) of Bayreuth, South Germany. 17, Skull from Bindlach, Bindlacher Berg (SMF No. R4572, original in Meyer, 1847-55), a. dorsal, b. lateral, ventral. 18, Coracoid from Bindlach, Bindlacher Berg (UM-O without no.), ventral. 19, Humerus from Bindlach, Bindlacher Berg (SMF no. R784), ventral. 20, Ischium from Bindlach, Bindlacher Berg (UM-O no. BT-670; original to Meyer, 1847-55), dorsal. 21, Femur from Bindlach, Bindlacher Berg (UM-O no. BT-722), dorsal. 22, Middle dorsal vertebral centrum (SMF no. R951), a. cranial, b. dorsal, c. lateral. 23, Posterior dorsal vertebral centrum (SMF without no.), a. cranial, b. dorsal, c. lateral (composed from Diedrich, 2012a, 2013d).



Figure 4 – *Lariosaurus teutonicus* (Diedrich, 1996), holotype skeleton from the Lower Terebratula Bed (Terebratula substage) of the Lower Muschelkalk (Pelsonian) of Borgholzhausen, NW Germany (coll. MNUO).

material from Bad Sulza (Diedrich, 2013d). Also, the skull from Bindlach (Fig. 3.17) illustrated in Edinger (1921) as *Nothosaurus* sp., and identified by Rieppel and Wild (1996) again as "*N. marchicus*," was revised in Diedrich (2012a) to *Lariosaurus* cf. *buzzii*, which was also found in the atavus oolite bonebeds of Bayreuth/Bindlach (Germany, Fig. 6, Map A). In addition to the cranium from the Bayreuth sites, the postcranial remains include a humerus, coracoid, femur and ischium, and also possibly a dorsal vertebra centrum (Figs. 3.18-23), which all are quite characteristic of *Lariosaurus* compared to a skeleton cast in the BSP (from Perledo, Italy, Fig. 2B), the descriptions of Curioni (1847) and Ticli (1984) and the figures of Tschanz (1989). Those remains from Bayreuth and Bad Sulza thus support the presence of this genus, as would be expected, in the Germanic Basin.

Other new evidence comes from a skull of similar age from Bad Sulza (Fig. 3.17) that was misidentified as "*N. marchicus*" in Rieppel and Wild (1996), which is in fact a similar *Lariosaurus* skull (Diedrich, 2013d). Also, from Bad Sulza, several postcranial bones with similar proportions, from all parts of the body, again indicate the presence of *Lariosaurus buzzii* Tschanz, 1989 in the Germanic Basin during the late Illyrian (?Fassanian). Also, the *L. buzzii* remains from the Anisian/Ladinian boundary of the northwestern Tethys (Fig. 6 Map B) are from lagoons and strata that formed during the maximum high stand of the Anisian/ Ladinian boundary (intraplatform lagoons: Furrer, 1995).

Longobardian – *Lariosaurus balsamii* (early Longobardian, Ladinian) skeletons are from the Kalkschieferzone (Uppermost Ladinian), such as those recorded at Valceresio (Varese, northern Italy (Tintori and Renesto, 1990) to which the herein juvenile Frankfurt specimen (Fig. 2A) is assigned. Also from Perledo (It), similar material was described (Curioni, 1847). This species seems to be represented by different aged individual skeletons, and is recently limited in the northern Tethys, but must also be expected in the Germanic Basin at least in the Bavarian-Hessian depression lagoons in southern Germany (Fig. 6 Map C).

CONCLUSIONS

In the Middle Triassic of Europe Lariosaurus is present not only in the northern Tethys as suggested before, it also occurs in the Germanic Basin, and these are even the oldest records (middle Pelsonian, middle Lower Muschekalk, Anisian) with the herein described holotype skeleton of *L. teutonicus* Diedrich, 1996, which was found in dolomite platy limestones of a lagoon facies. This is validated herein taxonomically with a detailed description and figures of the material. From the Germanic Basin the next records are isolated skulls and postcranial bones of similar age from the atavus bonebeds (middle Illyrian, basal Upper Muschelkalk, Anisian/Ladinian boundary) and similar oolitic, very shallow marine to lagoonal facies of Bad Sulza and Bayreuth, Germany. This material is dated in more detail, whereas this can be attributed to the about two million year younger species L. buzzii Tschanz, 1989, which is therefore represented starting with the new transgression in both the Germanic Basin and the northern Tethys, whereas a clear facies related (lagoon) occurrence of this genus can be demonstrated. The stratigraphically youngest record species is present somehow in the Lower Keuper (Longobardian, Ladinian), whereas in the Germanic basin those seem to have been disappeared (regression) and remained only in the northern Tethys with *L. balsami* Curioni (1841).

ACKNOWLEDGMENTS

The Naturkundemuseum Osnabrück (A. Leipner) made the studied specimens available, and supported this work with the preparation of the L. teutonicus holotype specimen and with photos. I thank Dr. G.R. Riedel for the possibility to study the main collection in the Naturkundemuseum Erfurt, and Dr. J.M. Rabold for the study of the historical Meyer collection in the Urwelt-Museum Oberfranken Bayreuth. Dr. R. Brocke from the Senckenberg Natural History Museum, Frankfurt, provided access to the historical bone collection from the Bayreuth sites. Dr. O. Hampe supported the collections work in the Natural History Museum of the Humboldt-University Berlin. The sponsor of this research, as part of the "Middle Triassic Megatracksite and Reptile Project of the Germanic Basin," was the Palaeologic Private Research Institution (www.paleologic.eu). I thank D. Surmik for a review of the first draft. Comments by an anonymous reviewer and S. G. Lucas improved the manuscript.

REFERENCES

- Curioni, G. 1847. Cenni sopra un nuovo saurio fossile die monti di Perledo sul Lario e sul terreno che le racchiude. Giornale del' J.R. Instituto Lombardo di Science Lettre ed Arti NS, 16: 159–170.
- Deecke, W. 1886. Ueber *Lariosaurus* und einige andere Saurier der Lombardischen Trias. Zeitschrift der Deutschen Geologischen Gesellschaft, 38 (1): 170–197.
- Diedrich, C. 1996. Paranothosaurus teutonicus n. sp. Aus dem Unteren Muschelkalk (Terebratel-Zone) von Borgholzhausen (NW-Deutschland) und seine Bedeutung für die Phylogenie und Taphonomie der Nothosauriden des germanischen Beckens. Terra Nostra, 96 (6): 36.
- Diedrich, C. 2009. The vertebrates of the Anisian/Ladinian boundary (Middle Triassic) from Bissendorf (NW Germany) and their contribution to the anatomy, palaeoecology, and palaeobiogeography of the Germanic Basin reptiles. Palaeogeography, Palaeoclimatology, Palaeoecology, 273: 1–16.
- Diedrich, C. 2009. Palaeogeographic evolution of the marine Middle Triassic marine Germanic Basin changements – with emphasis on the carbonate tidal flat and shallow marine habitats of reptiles in Central Pangaea. Global and Planetary Change, 65: 27–55.
- Diedrich, C. 2010. Palaeoecology of *Placodus gigas* (Reptilia) and other placodontids - MiddleTriassic macroalgae feeders in the Germanic Basin of central Europe - and evidence for convergent developed sirenia. Palaeogeography, Palaeoclimatology, Palaeoecology, 285: 287–306.
- Diedrich, C. 2011a. Fossil Middle Triassic "sea cows" placodont reptiles as macroalgae feeders along the North-western Tethys



Figure 5 – Lariosaurus teutonicus (Diedrich, 1996) holotype skeletal elements (coll. MNUO) from the Lower Terebratula Bed (Terebratula Substage) of the Lower Muschelkalk (Pelsonian) of Borgholzhausen, NW Germany.

coasts. Natural Science, 3(1): 8–26.

- Diedrich, C. 2011b. Middle Triassic limulid crustacean reproduction intertidal flats of Europe and contribution to the variability of *Kouphichnium* Nopcsa 1923 tracks – early Mesozoic limulid mating migrations into the Germanic Basin and reptile food chain reactions. Biological Journal of the Linnaean Society London, 103: 76–105.
- Diedrich, C. 2012a. The Middle Triassic marine reptile biodiversity in the Germanic Basin in the centre of the Pangaean world. Central European Journal of Geosciences,

4 (1): 9–46.

- Diedrich, C. 2012b. Middle Triassic chirotherid trackways on earthquake influenced intertidal limulid reproduction flats of the European Germanic Basin coasts. Central European Journal of Geosciences, 4(3): 495–529.
- Diedrich, C. 2013a. The "Middle Triassic Sea Cow" reptile *Placodus gigas* (Reptilia) populations of Pangaea's shallow marine macroalgae habitats. New Mexico Museum of Natural History and Science, 61: 104–130.

Diedrich, C. 2013b. The earliest "subaquatic flying" reptile -



Figure 5 – Palaeobiogeography and biostratigraphy of the three very shallow marine/lagoonal-adapted *Lariosaurus* species. *L. teutonicus*, *L. buzzii*, *L. balsami* in Europe (maps and stratigraphy from Diedrich, 2009b, 2012a, 2013d).

Pistosaurus longaevus Meyer 1839 (Sauropterygia) - from the Middle Triassic of the Germanic Basin of Central Europe. New Mexico Museum of Natural History and Science, 61: 169–215.

- Diedrich, C. 2013c. Marine pachypleurosaur *Serpianisaurus germanicus* nov. spec. remains from the late Pelsonian (Middle Triassic) of the Germanic Basin in Europe. New Mexico Museum of Natural History and Science, 61: 159–168.
- Diedrich, C. 2013d. Shallow marine sauropterygian reptile biodiversity and change in the Bad Sulza Fm (Illyrian, Middle Triassic) of Central Germany - and a contribution to the evolution of *Nothosaurus* in the Germanic Basin. New Mexico Museum of Natural History and Science, 61: 132–158.
- Furrer, H. 1995. The Kalkschieferzone (Upper Meride Limestone Ladinian) near Meride (Canton Ticino, Southern Switzerland) and the evolution of a Middle Triassic intraplatform basin.

Eclogae Geologicae Helvetiae, 88(3): 827-852.

- Furrer, H. 2003. Der Monte San Giorgio im Südtessin vom Berg der Saurier zur Fossillagerstätte internationaler Bedeutung. Neujahrsblatt der Naturforschenden Gesellschaft Zürich, 2004: 1–64.
- Kuhn-Schnyder, E. 1987. Die Triasfauna der Tessiner Kalkalpen. XXVI. *Lariosaurus lavizzarii* n. sp. (Reptilia, Sauropterygia). Schweizerische Paläontologische Abhandlungen, 110: 335–348.
- Kozur, H.W. & Bachmann, G.H. 2008, Updated correlation of the Germanic Triassic with the Tethyan scale and assigned numeric ages. Berichte der Geologischen Bundesanstalt Wien, 76: 53–58.
- Lukas, V. 1991. Die Terebratel-Bänke (Unterer Muschelkalk, Trias) in Hessen - ein Abbild kurzzeitiger Faziesänderungen im westlichen Germanischen Becken. Geologisches Jahrbuch

Hessen, 119: 119-175.

- Mazin, J.-M. 1985. A specimen of *Lariosanrus balsami* Curioni 1847, from the Eastern Pyrenees (France). Palaeontographica A, 189: 159–169.
- Meyer, von H. 1847–1855. Zur Fauna der Vorwelt. 2 Abt. Die Saurier des Muschelkalkes mit Rücksicht auf die Saurier aus Buntem Sandstein und Keuper (8 + 167 pp.). Frankfurt a. Main.
- Motani, R., Da-Yong Jiang, D.-Y. Tintori, A., Sun, Y.-L. Hao, W.-C Boyd, A., Sanja, H.F., Schmitz, L., Shin, J.-Y. & Sun, Z.-Y. 2008. Horizons and Assemblages of Middle Triassic Marine Reptiles from Panxian, Guizhou, China. Journal of Vertebrate Paleontology, 28(3): 900–903.
- Ogg, J.G., Huang, C. & Hinnov, L. 2014. Triassic time scale status: a brief overview. Albertiana, 41: 3–30.
- Peyer, B. 1933–34. Die Triasfauna der Tessiner Kalkalpen. VII. Neubeschreibung der Saurier von Perledo. Abhandlungen der Schweizerischen Paläontologischen Gesellschaft, 53–54: 1–130.
- Rieppel, O. 1994. *Lariosaurus balsami* Curioni (Reptilia, Sauropterygia) aus den Gailtaler Alpen. Carinthia, 184: 345–356.
- Rieppel, O. 1998. The status of the sauropterygian reptile genera *Ceresiosaurus*, *Lariosaurus*, and *Silvestrosaurus* from the Middle Triassic of Europe. Fieldiana. Geology New Series, 38: 1-46.
- Rieppel, O. 2000. Handbuch der Paläoherpetologie, Teil 12A. Sauropterygia I. Placodontia, Pachypleurosauria, Nothosauroidea, Pistosauroidea. Friedrich Pfeil-Verlag, München. 134 pp.

- Rieppel, O., Jinling, L. & Jun, L. 2003. *Lariosaurus xingyiensis* (Reptilia: Sauropterygia) from the Triassic of China. Canadian Journal of Earth Sciences, 40(4): 621–634
- Rieppel, O. & Wild, R. 1996. A revision of the genus *Nothosaurus* (Reptilia: Sauropterygia) from the Germanic Triassic, with comments on the Status of *Conchiosaurus clavatus*. Fieldiana Geology NS, 34: 1-82.
- Sanz, J.L. 1976. *Lariosaurus balsami* (Sauropterygia, Reptilia) de Estada (Huesca). Estudios geológicos, 32: 547–567.
- Stockar, R. 2010. Facies, depositional environment, and palaeoecology of the Middle Triassic Cassina beds (Meride Limestone, Monte San Giorgio, Switzerland). Swiss Journal of Geosciences, 103(1): 101–119.
- Ticli, B. 1984. Due esemplari di *Lariosaurus balsami* Curioni presenti nei Museo Civici di Lecco Natura. Memoire della Civico di Storia Naturale di Milano, 75(1–4): 69–74.
- Tintori, A. & Renesto, S. 1990. A new *Lariosaurus* from the Kalkschieferzone (Uppermost Ladinian) of Valceresio (Varese, N. Italy). Bolletino della Società Paleontologica Italiana, 29, 309–319.
- Tschanz, K. 1989. *Lariosaurus buzzii* n. sp. from the Middle Triassic of Monte San Giorgio (Switzerland) with comments on the classification on nothosaurs. Palaeontographica A, 208: 153–179.



Research Article

A SUMMARY OF NEW CONODONT BIOSTRATIGRAPHY AND CORRELATION OF THE ANISIAN (MIDDLE TRIASSIC) STRATA IN BRITISH COLUMBIA, CANADA

Martyn L. Golding¹, Michael J. Orchard², and John-Paul Zonneveld³

¹Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, 2020-2207 Main Mall, Vancouver, British Columbia, Canada V6T 1Z4. Email: mgolding@eos.ubc.ca

²Natural Resources Canada - Geological Survey of Canada, 1500 - 605 Robson Street, Vancouver, British Columbia, Canada, V6B 5J3. Email: morchard@nrcan.gc.ca

³Ichnology Research Group, Department of Earth & Atmospheric Sciences, University of Alberta, Edmonton, Alberta, Canada, T6G 2E3. Email: zonnevel@ualberta.ca

Abstract – New conodont collections from both outcrop sections and subsurface drill core improve the correlation of Anisian (Middle Triassic) strata in British Columbia. New conodont species and morphotypes identified in these collections are used to define new informal conodont faunal assemblages. This preliminary biostratigraphical scheme allows correlation of the Anisian between surface and subsurface sections in British Columbia for the first time, as illustrated by four of the sections studied. Improved correlation has led to a revision of the interpretation of the depositional environment of Anisian rocks in British Columbia, from that of a continental margin to that of a restricted basin. The recognition of some of the new conodont taxa in collections from Nevada and the Canadian Arctic indicates that some of the faunal assemblages may prove useful for regional correlation and placement of the sub-stage boundaries of the Anisian.

INTRODUCTION

This paper introduces new conodont taxa from the Anisian of northeastern British Columbia, and new biostratigraphical correlations based on their distribution. This work both summarises and expands upon observations and conclusions made by Golding (2014).

Ammonoids and conodonts are the two most useful and widely used taxonomic groups for biostratigraphical subdivision of the marine Anisian. The base of the Anisian has not yet been defined by a Global Boundary Stratotype Section and Point (GSSP; Ogg et al., 2014). The only candidate section so far proposed is at Desli Caira in Romania, where the boundary has been proposed at the first appearance of the conodont Chiosella timorensis (Nogami) (Gradinaru et al., 2007; Orchard et al., 2007 a). However, the first appearance of this species is now known to occur with Spathian (upper Olenekian) ammonoids of the Neopopanoceras haugi Zone in Nevada and China (Ovtcharova et al., 2010; Goudemand et al., 2012), therefore its use as a definitive index species for the base of the Anisian is questionable, and would require the Neopopanoceras haugi Zone to be re-defined as Anisian in part. The complementary section at Guandao in China (Lehrmann et al., 2006) provides a superior conodont record (Orchard et al., 2007b), but lacks good ammonoid control

		Ammonoid and Conodont Zones (Orchard and Tozer, 1997)							
	U	chischa	aldae		excelsa				
		deleeni	constricta						
	м	minor	"n. sp. P"		bulgarica				
z		minor	shoshonensis		regalis				
ISIA		hayesi	"n. sp. N"						
AN		hagei							
	L	americanum							
		caurus	"n. sp. M"						
		mulleri	Gorman		l	1			
			timorer		nsis				

Figure 1 – Existing ammonoid and conodont biozonation of the Anisian in British Columbia based on Orchard and Tozer (1997). L = Lower, M = Middle, U = Upper.

(Ogg et al., 2014). The upper boundary of the Anisian is formally defined by the first appearance of the ammonoid *Eoprotrachyceras curionii* (Mojsisovics) at the base of the Ladinian, in the section at Bagolino in Italy (Brack et al., 2005).

In British Columbia, the Anisian ammonoid record is fairly well developed, with eight ammonoid zones and two subzones recognized (Fig. 1; Tozer, 1967, 1994; Silberling & Tozer, 1968; Bucher, 2002). However, the Anisian sequence in British Columbia appears rather incomplete by reference to that of Nevada, where 13 ammonoid zones and 31 subzones are recognized (Silberling & Wallace, 1969; Silberling & Nichols, 1982; Bucher, 1988, 1989, 1992 a, b, 1994; Bucher & Orchard, 1995). Lowermost Anisian ammonoids appear to be missing in British Columbia, and there are large gaps in the ammonoid record of the Middle and Upper Anisian (Monnet & Bucher, 2005). The reasons for this discrepancy between the two areas are unclear, but may be due to unconformities in the Anisian succession of British Columbia, or the difference may simply be a reflection of the higher ammonoid diversity in Nevada (Monnet & Bucher, 2005).

The conodont record of the Anisian in British Columbia is generally poor in resolution compared with that of the ammonoids. Eleven partially overlapping conodont zones have been outlined for the Anisian (Orchard & Tozer, 1997), although several of these were based on undescribed taxa. In the upper part of the Middle Anisian and in the Upper Anisian the precision of the existing conodont zonation is comparable to that of the ammonoids, however, within the Lower Anisian and lower part of the Middle Anisian, the precision is not as good (Fig. 1).

The deficiency in the existing conodont biozonation is due, in part, to an underdeveloped taxonomy for Anisian conodonts in North America, particularly the ubiquitous Neogondolellins and Paragondolellins (sensu Orchard, 2005). In an attempt to improve the stratigraphic utility of the conodonts, new conodont samples were collected from both surface outcrop and subsurface drill core in British Columbia, and examined together with existing collections which are calibrated with ammonoid zones (Orchard & Tozer, 1997). This has resulted in the taxonomic revision of a number of conodont species groups, the refinement of the faunal succession that is tied to the existing ammonoid biozonation, and improved correlation between outcrop and subsurface Triassic.

GEOLOGICAL SETTING

The Anisian rocks of British Columbia belong to two major tectonic settings: some were deposited on the continental margin of North America, whilst others were deposited on pericratonic and exotic terranes that lay to the west of the continent during the Triassic. These terranes have subsequently accreted onto the margin of North America, and rocks of Anisian age are preserved throughout British Columbia (Orchard, 1991). Those that were deposited on the continental margin form a near-continuous outcrop belt running along the eastern margin of the Canadian Rocky Mountains (Fig. 2). To the east of this outcrop belt, Anisian rocks are present in the subsurface of British Columbia and the



Figure 2 – Map showing the distribution of Triassic outcrop in northeastern British Columbia and the location of the four sections mentioned in the text. Triassic outcrop is in purple.

neighboring province of Alberta.

The Anisian rocks of the subsurface and eastern outcrop belt were deposited in the Western Canada Sedimentary Basin (WCSB), which was made up of a complex series of sub-basins formed by a combination of transtension and extension (Davies, 1997). The basin initiated during the Mesoproterozoic and persisted as a continental margin basin until terrane accretion began in the Triassic (Ross et al., 1989; Beranek and Mortensen, 2011). During the Anisian, the WCSB was located on the western margin of North America at mid-latitudes (Smith et al., 1995), and experienced an arid climate influenced by trade winds and upwelling (Davies, 1997).

Anisian rocks of the outcrop belt belong to the Toad Formation. This unit was named by Kindle (1944) and it consists of up to 800 m of siltstone, shale, fine-grained sandstone and carbonate (Gibson, 1975). The formation is thought to range in age from the Smithian (lower Olenekian) to the Ladinian on the basis of ammonoid and conodont collections (Tozer, 1961, 1994; Orchard & Tozer, 1997; Zonneveld, 2010). In the subsurface, the Anisian rocks belong to the Montney and Doig formations, both defined by Armitage (1962). The Montney Formation consists of up to 450 m of siltstone and sandstone with very minor amounts of shale and carbonate (Armitage, 1962; Davies et al., 1997). The formation is thought to range in age from Changhsingian to Anisian (Zonneveld, 2010; Zubin-Stathopoulos et al., 2012). The Doig Formation unconformably
overlies the Montney Formation, and consists of up to 150 m of siltstone, fine sandstone and carbonate (Armitage, 1962; Evoy & Moslow, 1995). The age of this formation is uncertain, but has been estimated as Spathian to Ladinian (Hunt & Ratcliffe, 1959; Qi, 1995; Zonneveld, 2010). Both the Montney and Doig formations contain economically important reserves of natural gas (Adams, 2013).

A total of 42 conodont collections from 11 outcrop sections and 7 subsurface wells, all representing the ancestral North American continental margin, were examined for this study; the location of those discussed in this paper are shown in Fig. 2. A full description of all of the localities is available in Golding (2014).

NEW CONODONT SPECIES

Previously, the majority of conodont species from the Anisian of North America have been assigned to one of six named species: Neogondolella regalis Mosher, N. constricta (Mosher and Clark), N. bulgarica (Budurov and Stefanov), N. shoshonensis Nicora, N. mombergensis (Tatge), and Paragondolella excelsa Mosher (Mosher & Clark, 1965; Mosher, 1973; Nicora, 1976, 1977; Nicora & Kovács, 1984; Ritter, 1989; Orchard, 1994), although Nicora et al. (1980) and Orchard (2010) also differentiated several other less common taxa from Nevada. The current taxonomic revision has resulted in the recognition of a number of new morphotypes of some of the common species from British Columbia, which have been re-classified as species groups. Previous reports of N. mombergensis, the type species of Neogondolella, from North America (e.g. Ritter, 1989) are regarded as erroneous, and the existence of this species outside of Tethys is questioned (Bucher & Orchard, 1995; Orchard & Rieber, 1999). In addition to 18 new morphotypes of four species groups assigned to Neogondolella and Paragondolella, a further nine new species of Neogondolella have been differentiated; these new taxa will all be fully described in future publications. A selection of some of the most stratigraphically important new taxa are shown in Fig. 3.

The Canadian succession of these new conodont taxa has allowed the recognition of 11 new, informal faunal assemblages that are useful for local correlation. These assemblages are groups of species that will hopefully define biozones, once the variations in morphology, and geographic and stratigraphic ranges of the constituent species have been fully determined. Seven of these assemblages are introduced informally here to illustrate their use in correlation (Fig. 4). Currently, the new biostratigraphical scheme is a hybrid, with the *N*. ex gr. *shoshonensis* faunal assemblage equivalent in concept to an assemblage biozone, whilst the others are akin to interval biozones, defined by first appearances of taxa.

CORRELATION OF THE ANISIAN IN B.C.

In order to demonstrate the utility of the new conodont faunal assemblages in the correlation of the Anisian of northeastern British Columbia, four sections are chosen. Two of these are outcrop sections located on the Alaska Highway, at Mile Post 375 East and Mile Post 375 West (Fig. 2). The Toad Formation is exposed at these sections on the two limbs of an anticline, and they represent the type sections of the Lenotropites caurus, Hollandites minor, and Eogymnotoceras deleeni ammonoid Zones (Tozer, 1967, 1994); Bucher (2002) also differentiated the Paracrochordiceras americanum beds at the Mile Post 375 East section. At these sections, the age of the conodont faunal assemblages has been directly tied to the ammonoid biozonation (Fig. 4). The other two sections are drill core from hydrocarbon wells drilled in the vicinity of Fort St. John, named Talisman Altares 16-17-083-25W6 and Murphy Swan d-054-B/093-P-09 (Fig. 2). The drill core from these wells belongs to the Montney and Doig formations, and conodont collections from the wells can be assigned to the same faunal assemblages as in the outcrop sections. However, no ammonoids were collected from the drill core. The correlation between this drill core and the outcrop sections is shown in Fig. 4. The stratigraphy and conodont zonation of the Talisman Altares 16-17-083-25W6 well is discussed in detail in another manuscript (Golding et al., in press).

The geographical separation of these sections is quite large, nevertheless it is possible to draw firm correlations between them, particularly around the Lower-Middle Anisian boundary. This is placed between the *Paracrochodiceras americanum* and *Buddhaites hagei* ammonoid zones (Tozer, 1994) and, although no conodont fauna was recovered from the *Buddhaites hagei* Zone (the presence of which was reported from the Mile Post 375 East section by Bucher, 2002), the division between the *Neogondolella* ex gr. *regalis* morphotype D and the *Neogondolella* ex gr. *regalis* morphotype B faunal assemblages is inferred to approximate the boundary. The recognition of these faunal assemblage in the subsurface drill core allows the correlation of the Lower-Middle Anisian boundary between the outcrop and the subsurface (Fig. 4).

IMPLICATIONS OF THE NEW CORRELATION

The correlation of the Anisian using new conodont faunal assemblages allows some of the previously accepted ideas about the Anisian of British Columbia to be tested. The boundary between the Montney and Doig formations has previously been thought to be diachronous, with estimates for its age ranging from Spathian to Ladinian (Hunt & Ratcliffe, 1959; Qi, 1995; Zonneveld, 2010), however neither the degree nor pattern of diachroneity has been well understood. The new correlation described herein shows that the formational boundary is oldest in the center of the study area, just west of Fort St. John, and becomes younger both to the west and to the east (Golding, 2014). The base of the Doig Formation is Spathian in age in the Altares area of British Columbia, and ranges into the Middle Anisian in the Dawson area. This pattern of diachroneity is not what would have been expected if the sediment of the Montney and Doig formations was deposited on a west-facing continental margin. Instead, it implies that the depositional basin was restricted during the Anisian, with localized palaeo-highs present to the west as well as to the east of the depocentre (Golding, 2014). The new correlation also shows that the basal Doig Formation is condensed, with



Figure 3 – Some stratigraphically important conodonts from the Anisian of northeastern British Columbia. **1–3**, *Neogondolella* ex gr. *regalis* Mosher, morphotype A nov., GSC 131502, *Tetsaoceras hayesi* Zone, Toad Formation, Mile Post 375 West (GSC cur. no. 68294). **4–6**, *Neogondolella* ex gr. *shoshonensis* Nicora, GSC 131503, *Hollandites minor* Zone, Toad Formation, Mile Post 375 West (GSC cur. no. C-201926). **7–9**, *Paragondolella* ex gr. *liebermani* Kovács and Krystyn, morphotype A nov., GSC 131504, *Eogymnotoceras deleeni* Zone, Toad Formation, Mile Post 375 West (GSC cur. no. C-201926). **7–9**, *Paragondolella* ex gr. *liebermani* Kovács and Krystyn, morphotype A nov., GSC 131504, *Eogymnotoceras deleeni* Zone, Toad Formation, Mile Post 375 West (GSC cur. no. C-201925). **10–12**, *Neogondolella* ex gr. *regalis* morphotype B nov., GSC 131505, *Tetsaoceras hayesi* Zone, Doig Formation, Murphy Swan d-054-B/093-P-09 well (GSC cur. no. V-002957); **13–15**, *Neogondolella* ex gr. *regalis* morphotype C nov., GSC 131506, *Tetsaoceras hayesi* Zone, Toad Formation, Mile Post 375 East (GSC cur. no. 68204). **16–18**, *Neogondolella* n. sp. N sensu Orchard and Tozer, 1997, GSC 131507, *Tetsaoceras hayesi* Zone, Toad Formation, Mile Post 375 East (GSC cur. no. 68204). **19–21**, *Neogondolella* n. sp. M sensu Orchard and Tozer, 1997, GSC 131508, *Lenotropites caurus* Zone, Toad Formation, Mile Post 375 East (GSC cur. no. C-201922).



Figure 4 – Correlation of the sections at Mile Post 375 East, Mile Post 375 West, Talisman Altares 16-17-083-25 W6 and Murphy Swan d-054-B/093-P-09 using seven of the new conodont faunal assemblages, that are in turn calibrated to the existing ammonoid biozonation. Dashed line between *Paracrochordiceras americanum* and *Tetsaoceras hayesi* zones indicates the absence of the *Buddhaites hagei* Zone. N. = *Neogondolella*, P. = *Paragondolella*. In outcrop sections, the scale is in meters above the base of the section, and in subsurface drill core, the scale is in meters below kelly bushing.

the highest degree of condensation occurring where the base of the formation is youngest, and the least where it is oldest. This pattern of condensation again suggests the presence of localized paleo-highs within the basin (Golding, 2014).

The large number of new conodont species and morphotypes used to define the new faunal assemblage scheme currently precludes correlation beyond the local area. Anisian collections of conodonts from Nevada contain rich faunas of *Neogondolella* ex gr. *constricta*, and it is probable that some of the new species recognized in British Columbia can also be recognized amongst this diverse fauna. Morphotypes of *N*. ex gr. *shoshonenesis* have been identified in Nevada (Nicora, 1976) and may be useful for stratigraphic correlation between the two areas; this species group appears to be restricted to the Middle and Upper Anisian (Nicora, 1976; Golding, 2014). *Neogondolella* n. sp. C sensu Orchard, 2008 appears in the Lower Anisian in British Columbia and similar forms seem to occur at the same level in Nevada and in the upper Spathian of the Canadian Arctic (Orchard, 2008). Further investigation of Anisian faunas from Nevada and the Canadian Arctic is required to determine the degree to which these new faunal assemblages are suitable for regional correlation, or to what extent they may be useful for correlation with sub-stage boundaries used in Tethys.

← Figure 3 (continued). 22–24, Neogondolella ex gr. regalis morphotype D nov., GSC 131476, Paracrochordiceras americanum Zone, Doig Formation, Talisman Altares 16-17-083-25 W6 well (GSC cur. no. V-002978). 25–27, Neogondolella ex gr. regalis morphotype E nov., GSC 131509, Paracrochordiceras americanum Zone, Toad Formation, Mile Post 375 East (GSC cur. no. C-209952). Magnification is × 80.

CONCLUSIONS

The recognition of new conodont taxa from the Anisian of British Columbia has improved the resolution of the conodont biozonation for this time period. The new faunal assemblages allow more precise correlation between sections, including from the outcrop to the subsurface for the first time. The correlation of the subsurface sections has allowed the depositional environment of the Anisian rocks to be re-assessed; it is suggested that these rocks were not deposited on a continental margin, but within a semi-restricted basin. A number of the new conodont taxa identified within the Anisian of British Columbia are also present in Nevada and the Canadian Arctic, suggesting the potential for recognizing the conodont faunal assemblages throughout North America.

ACKNOWLEDGEMENTS

Charles Henderson (University of Calgary) and Chris McRoberts (SUNY Cortland) are thanked for their helpful reviews of the manuscript. Financial support for this study was provided in part by the Yukon Basins Project of the Geological Survey of Canada GEM Program, and by Geoscience B.C.

REFERENCES

- Adams, C. 2013. Summary of shale gas activity in northeast British Columbia 2012. B.C. Ministry of Energy and Mines: 1–27.
- Armitage, J.H. 1962. Triassic oil and gas occurrences in northeastern British Columbia, Canada. Journal of the Alberta Society of Petroleum Geologists, 10: 35–56.
- Beranek, L.P. & Mortensen, J.K. 2011. The timing and provenance record of the Late Permian Klondike orogeny in northwestern Canada and arc-continent collision along western North America. Tectonics, 30: 1–23.
- Brack, P., Rieber, H., Nicora, A. & Mundil, R., 2005. The Global boundary Stratotype Section and Point (GSSP) of the Ladinian Stage (Middle Triassic) at Bagolino (Southern Alps, Northern Italy) and its implications for the Triassic timescale. Episodes, 28: 233–244.
- Bucher, H. 1988. A new Middle Anisian (Middle Triassic) ammonoid zone from northwestern Nevada (USA). Eclogae Geologicae Helvetiae, 81: 723–762.
- Bucher, H. 1989. Lower Anisian ammonoids from the northern Humboldt Range (northwestern Nevada, USA) and their bearing upon the Lower-Middle Triassic boundary. Eclogae Geologicae Helvetiae, 82: 945–1002.
- Bucher, H. 1992 a. Ammonoids of the *Hyatti* Zone and the Anisian transgression in the Star Peak Group, northwestern Nevada, USA. Palaeontographica Abteilung A, 223: 137–166.

Bucher, H. 1992 b. Ammonoids of the *Shoshonensis* Zone (Middle Anisian, Middle Triassic) from northwestern Nevada (USA). Jahrbuch der Geologischen Bundesanstalt Wien, 135: 425-465.

- Bucher, H. 1994. New ammonoids from the Taylori Zone (Middle Anisian, Middle Triassic) from northwestern Nevada (USA). Mémoires de Géologie (Lausanne), 22: 1–8.
- Bucher, H. 2002. Early Anisian (Middle Triassic) ammonoid biostratigraphy of northeastern British Columbia. Eclogae Geologicae Helvetiae, 95: 277–287.
- Bucher, H. & Orchard, M.J. 1995. Intercalibrated ammonoid and conodont succession, Upper Anisian-Lower Ladinian of Nevada. Albertiana, 15: 66–71.
- Davies, G.R. 1997. The Triassic of the Western Canada Sedimentary Basin: tectonic and stratigraphic framework, paleogeography, paleoclimate and biota. *In*, Moslow, T.F. & Wittenberg, J. (eds.), Triassic of the Western Canada Sedimentary Basin, Bulletin of Canadian Petroleum Geology, 45: 434–460.
- Davies, G.R., Moslow, T.F. & Sherwin, M.D. 1997. The Lower Triassic Montney Formation, west-central Alberta. *In*, Moslow, T.F. & Wittenberg, J. (eds.), Triassic of the Western Canada Sedimentary Basin, Bulletin of Canadian Petroleum Geology, 45: 474–505.
- Evoy, R.W. & Moslow, T.F. 1995. Lithofacies associations and depositional environments in the Middle Triassic Doig Formation, Buick Creek Field, northeastern British Columbia. Bulletin of Canadian Petroleum Geology, 43: 461–475.
- Gibson, D.W. 1975. Triassic rocks of the Rocky Mountain Foothills and Front Ranges of northeastern British Columbia and west-central Alberta. Geological Survey of Canada Bulletin 247, 61 pp.
- Golding, M.L. 2014. Biostratigraphy and sedimentology of Triassic hydrocarbon-bearing rocks in northeastern British Columbia. [Ph.D. Dissertation]: University of British Columbia, 402 pp.
- Golding, M.L., Orchard, M.J., Zonneveld, J.-P., Henderson, C.M. & Dunn, L. in press. An exceptional record of the sedimentology and biostratigraphy of the Montney and Doig formations in British Columbia. Bulletin of Canadian Petroleum Geology.
- Goudemand, N., Orchard, M.J., Bucher, H. & Jenks, J. 2012. The elusive origin of *Chiosella timorensis* (Conodont Triassic). Geobios, 45: 199–207.
- Gradinaru, E., Orchard, M.J., Nicora, A., Gallet, Y., Besse,
 J., Krystyn, L., Sobolev, E.S., Atudorei, V. & Ivanovna, D.
 2007. The Global Boundary Stratotype Section and Point (GSSP) for the base of the Anisian Stage: Desli Caira Hill,
 North Dobrogea, Romania. Albertiana, 36: 54–71.
- Hunt, A.D. & Ratcliffe, J.D. 1959. Triassic stratigraphy, Peace River area, Alberta and British Columbia, Canada. Bulletin of the American Association of Petroleum Geologists, 43: 563–589.
- Kindle, E.D. 1944. Geological reconnaissance along Fort Nelson, Liard and Beaver Rivers, northeastern British Columbia and southeastern Yukon. Geological Survey of Canada Paper 44-16, 19 pp.
- Lehrmann, D.J., Ramezani, J., Bowring, S.A., Martin, M.W.,

Montgomery, P., Enos, P., Payne, J.L., Orchard, M.J., Hongmei, W. & Jiayong, W. 2006. Timing of recovery from the end-Permian extinction: geochronologic and biostratigraphic constraints from south China. Geology, 34: 1053–1056.

Monnet, C. & Bucher, H. 2005. Anisian (Middle Triassic) ammonoids from North America: quantitative biochronology and biodiversity. Stratigraphy, 2: 281–296.

Mosher, C. & Clark, D.L. 1965. Middle Triassic conodonts from the Prida Formation of Northwestern Nevada. Journal of Paleontology, 39: 551–565.

Nicora, A. 1976. Conodont-fauna, stratigraphic position and relations to the Tethyan succession of the *Shoshonensis* Zone (Pelsonian) of Nevada. Rivista Italiana di Paleontologia e Stratigrafia, 82: 627–648.

Nicora, A. 1977. Lower Anisian platform-conodonts from the Tethys and Nevada: taxonomic and stratigraphic revision. Palaeontographica Abteilung A, 157: 88–107.

Nicora, A. & Kovács, S. 1984. Conodont fauna from the *Rotelliformis, Meeki* and *Occidentalis* zones (Middle Triassic) of Humboldt Range, Nevada, Western North America. Rivista Italiana di Paleontologia e Stratigrafia, 90: 135–164.

Nicora, A., Kozur, H. & Mietto, P. 1980. Gondolella pridaensis sp. n. A new conodont species from the Middle Triassic. Rivista Italiana di Paleontologia, 86: 761–768.

Ogg, J.G., Huang, C. & Hinnov, L. 2014. Triassic timeset acale status: a brief overview. Albertiana, 41: 3–30.

Orchard, M.J. 1991. Conodonts, time and terranes: an overview of the biostratigraphic record in the western Canadian Cordillera. *In*, Orchard, M.J. & McCracken, A.D. (eds.), Ordovician to Triassic conodont paleontology of the Canadian Cordillera. Geological Survey of Canada Bulletin 417: 1–26.

Orchard, M.J. 1994. Conodont biochronology around the Early-Middle Triassic boundary: new data from North America, Oman and Timor. Mémoires de Géologie (Lausanne), 22: 105–114.

Orchard, M.J. 2005. Multielement conodont apparatuses of Triassic Gondolelloidea. *In*, Purnell, M.A. & Donoghue, P.C.J. (eds.), Conodont biology and phylogeny: interpreting the fossil record. Special Papers in Palaeontology, 73: 73–101.

Orchard, M.J. 2008. Lower Triassic conodonts from the Canadian Arctic, their intercalibration with ammonoidbased stages and a comparison with other North American Olenekian faunas. Polar Research, 27: 393–412.

Orchard, M.J. 2010. Triassic conodonts and their role in stage boundary definition. Journal of the Geological Society of London, 334: 139–161.

Orchard, M.J., Gradinaru. E. & Nicora, A. 2007 a. Conodonts from the Olenekian-Anisian boundary interval at Desli Caira, Dobrogea, Romania. New Mexico Museum of Natural History and Science Bulletin 41: 341–346.

Orchard, M.J., Lehrmann, D., Wei Jiayong, Wang Hongmei, & Taylor, H. 2007 b. Conodonts from the Olenekian-Anisian boundary beds, Guandao, Guizhou Province, China. New Mexico Museum of Natural History and Science Bulletin 41: 347–354.

Orchard M.J. & Rieber, H. 1999. Multielement *Neogondolella* (Conodonta, upper Permian – middle Triassic). Bollettino della Società Paleontologica Italiana, 37: 475–488.

Orchard, M.J. & Tozer, E.T. 1997. Triassic conodont biochronology, its calibration with the ammonoid standard, and a biostratigraphic summary of the Western Canada Sedimentary Basin. *In*, Moslow, T.F. & Wittenberg, J. (eds.), Triassic of the Western Canada Sedimentary Basin, Bulletin of Canadian Petroleum Geology, 45: 675–692.

Ovtcharova, M., Bucher, H., Goudemand, N., Schaltegger, U., Brayard, A. & Galfetti, T. 2010. New U/Pb ages from Nanpanjiang Basin (South China): implications for the age and definition of the Early-Middle Triassic boundary. Geophysical Research Abstracts, 12: EGU2010-12505-3

Qi, F. 1995. Seismic stratigraphy and sedimentary facies of the Middle Triassic strata, Western Canada Sedimentary Basin, northeastern British Columbia. [Ph.D. Dissertation]: University of Alberta, 219 pp.

Ritter, S.M. 1989. Morphometric patterns in Middle Triassic *Neogondolella mombergensis* (Conodonta), Fossil Hill, Nevada. Journal of Paleontology, 63: 233–245.

Ross, G.M., McMechan, M.E. & Hein, F.J. 1989. Proterozoic history: the birth of the miogeocline. *In*, Ricketts, B.D. (ed.), Western Canada Sedimentary Basin – a Case History, Canadian Society of Petroleum Geologists, Special Publication, 30: 79–104.

Silberling, N.J. & Nichols, K.M. 1982. Middle Triassic molluscan fossils of biostratigraphic significance from the Humboldt Range, northwestern Nevada. U.S. Geological Survey Professional Paper 1207, 77 pp.

Silberling, N.J. & Tozer, E.T. 1968. Biostratigraphic classification of the marine Triassic in North America. U.S. Geological Survey Special Paper 110, 63 pp.

Silberling, N.J. & Wallace, R.E. 1969. Stratigraphy of the Star Peak Group (Triassic) and overlying Mesozoic rocks, Humboldt Range, Nevada. U.S. Geological Survey Professional Paper 592, 50 pp.

Smith, A.G., Smith, D.G. & Funnell, B.M. 1995. Atlas of Mesozoic and Cenozoic Coastlines. Cambridge University Press, 99 pp.

Tozer, E.T. 1961. The sequence of marine Triassic faunas in western Canada. Geological Survey of Canada Paper 61-6, 20 pp.

Tozer, E.T. 1967. A standard for Triassic time. Geological Survey of Canada Bulletin 156, 103 pp.

Tozer, E.T. 1994. Canadian Triassic Ammonoid Faunas. Geological Survey of Canada Bulletin 467, 663 pp.

Zonneveld, J.-P. 2010. The Triassic of northeastern British Columbia: sedimentary characteristics and stratigraphic architecture of conventional and unconventional reservoir successions. Geological Association of Canada-Mineralogical Association of Canada Joint Annual Meeting (GeoCanada 2010), Williston Lake Field Trip Guidebook, 158 pp.

Zubin-Stathopoulos, K.D., Beauchamp, B., Davydov, V.I. & Henderson, C.M. 2012. Variability of PennsylvanianPermian Carbonate Associations and implications for NW Pangea Paleogeography, east-central British Columbia, Canada. *In*, Gasiewicz, A. & Slowakiewicz, M. (eds.), Paleozoic Climate Cycles: Their Evolutionary and Sedimentological Impact, Geological Society, London, Special Publications, 376: 47–71.



Research Article

TIME CALIBRATED EARLY TRIASSIC $\delta^{13}C_{_{carb}}$, $\delta^{18}O_{_{apatite}}$ and SST curves from south china: An update

Nicolas Goudemand

Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006 Zurich, Switzerland, Email: nicolas.goudemand@pim.uzh.ch.

Abstract – Since Sun et al. (2012) published the first $\delta^{18}O_{apatite}$ data of conodonts from sections in the Nanpanjiang Basin, South China and used it to reconstruct Late Permian to Middle Triassic eastern Tethyan, equatorial seawater temperatures, errata were published and new radiometric ages became available, which prompted me to reanalyze this unique dataset. In the following I explore how the new data and corresponding modifications impact the shape of the Early Triassic $\delta^{13}C$ and $\delta^{18}O$ curves from South China, their implications for the evolution of sea surface temperatures, and their relationship with the biotic recovery in the aftermath of the Permo-Triassic crisis.

INTRODUCTION

In 2012, Sun et al. published the first $\delta^{18}O_{apatite}$ data of conodonts from sections in the Nanpanjiang Basin, South China and they used this data to reconstruct Late Permian to Middle Triassic eastern Tethyan, equatorial seawater temperatures. Since then, new information and mistakes have come to light. Goudemand et al. (2013) wrote a technical comment and proposed corrections to Sun et al.'s (2012) figures 2 and 3. As admitted by Sun et al. in their reply (2013), some data points were erroneous (their table S4 does not match their figs. 2 and 3). Goudemand et al.'s (2013) figure 1 was drawn after figure 4 in Galfetti et al. (2007), however, which referenced an incorrect age of ca. 247.9 Ma for the Early-Middle Triassic boundary instead of the new age of 247.2 Ma proposed by Lehrmann et al. (2007). This age (247.2 Ma) remains valid with the new (but still informal) boundary definition by Goudemand et al. (2012), which places the boundary between the 'carinata beds' and the 'tethydis beds', clearly above the FAD of Chiosella timorensis. Furthermore, Burgess et al. (2014) proposed a new radiometric age of 251.9 Ma for the base of the Triassic as well as new absolute time constraints. In the following I explore how these findings impact the shape of the Early Triassic $\delta^{\rm 13}C$ and $\delta^{\rm 18}O$ curves from South China and their implications for the relationship between seawater temperatures and biotic recovery in the aftermath of the Permo-Triassic crisis.

Stratigraphic sequence in Zuodeng

Sun et al.'s (2012) figure 2 shows that most $\delta 18O_{apatite}$ and $\delta^{13}C_{carb}$ data for the Early Triassic of the Nanpanjiang basin is from Zuodeng section, and it spans from the base of the Dienerian to the late Spathian. Hence, I used the data from

Zuodeng section as a basis for the time calibration. Once this data was calibrated, the biochronological correlations could be used to fit the data from other sections.

Figure 1 displays the raw data from Zuodeng II section from Sun et al. (2012, tab. S4). The y-axis represents here the stratigraphical position of the samples. The light and dark green bands indicate the extent of each biozone as given by Sun et al., taking into account that ZDC50 and ZDC51 have been reassigned to the *Novispathodus pingdingshanensis* Zone (Sun et al., 2013).

ABSOLUTE AGE CONSTRAINTS

In order to plot the corresponding biozones on an absolute timescale, the following absolute ages were taken into consideration:

(1) An age of 252.0 Ma for the middle of bed 23 in Meishan (Burgess et al., 2014). This age constrains the position of the *Neogondolella yini* Zone (C2 in Fig. 2), which starts from bed 24 in Meishan (Joachimski et al., 2012).

(2) An age of 251.9 Ma for the Permo-Triassic Boundary (Burgess et al., 2014).

(3) & (4) Two ages at 251.6 Ma and 251.5 Ma respectively from beds 33 and 34 at Meishan (Burgess et al., 2014), which are within the *Neogondolella planata* Zone (Zhang et al., 2007). These two ages constrain respectively the lower and upper ends of a minimum $\delta^{13}C_{carb}$ plateau (see Burgess et al., 2014; data from Cao et al., 2010).

(5) An age of 251.2Ma between the *Kashmirites densistriatus* beds and the *Flemingites rursiradiatus* beds (Galfetti et al., 2007, Brayard and Bucher, 2008). This age is late early Smithian but not earliest Smithian sensu the latest definition of the Dienerian/



Figure 1 – Stratigraphical evolution of $\delta^{13}C_{carb}$ and $\delta^{18}O_{apatite}$ data in the Zuodeng II section. Data from Sun et al. (2012). Sun et al. (2012) report that the measurement of NBS 120c was 22.6±0.16‰ VSMOW. C9-C14, conodont biozones after Sun et al. (2012). Assignment of samples to corresponding biozones was done after Sun et al. (2012) with corrections from Sun et al. (2013); C9, *Neospathodus dieneri* Zone (Dienerian); C10, *Novispathodus waageni* Zone (early and middle Smithian); C11, *Parachirognathus* Zone (mid-late Smithian); C12: *Novispathodus pingdingshanensis* Zone (early Spathian?); C13, *Icriospathodus collinsoni* Zone (late early Spathian); C14, *Triassospathodus homeri* Zone (middle and late Spathian).

Smithian boundary (Induan-Olenekian boundary, IOB) based on the latest ammonoid results (Brühwiler et al., 2010) that are also corroborated by conodonts and compatible with the latest consensus of the IOBWG about a conodont-based definition of the IOB (Goudemand, this issue). This age does however correlate quite well with the $\delta^{13}C_{carb}$ maximum near the base of the Smithian. This shows that the base of the Smithian should not be considered to fall exactly on that $\delta^{13}C_{carb}$ maximum, and this is confirmed at the West Pingdingshan section (Chaohu, Anhui province, South China), a GSSP candidate for the IOB (Tong et al., 2007, fig. 2). (6) An age of 250.5 Ma within the *Tirolitesl Columbites* beds (Galfetti et al., 2007). This age correlates roughly with the end of an early Spathian, positive $\delta^{13}C_{carb}$ plateau. Given the uncertainty associated with this age, Goudemand et al. (2013) had used the lower bound of this age (about 250.15 Ma) for calibrating the $\delta^{13}C_{carb}$ curve (their fig.1).

(7) An age of 248.1 Ma within the late Spathian *Haugi* Zone (Ovtcharova et al., 2006). The corresponding ash layer is found below the maximum $\delta^{13}C_{carb}$ plateau that straddles the Early-Middle Triassic boundary, approximately before the Chinese composite $\delta^{13}C_{carb}$ curve outreaches +2‰ (see for instance



Figure 2 – Temporal evolution of $\delta^{13}C_{carb}$ and $\delta^{18}O_{apattle}$ data from Meishan. Brown circles: $\delta^{13}C_{carb}$ data from Cao et al. (2010); red circles: $\delta^{18}O_{apattle}$ data from Joachimski et al. (2012), $\delta^{13}C_{carb}$ data from Xie et al. (2007), Shangsi, Wantuo, Jinya, Jiarong, Bianyang, Guandao and Zuodeng (all data from Sun et al., 2012) sections. The $\delta^{18}O_{apattle}$ datasets are standardized to NBS 120c=22.6±0.16‰ VSMOW. C1-C18, conodont biozones modified after Sun et al. (2012). C16 and C17 are added because they are used Sun et al. (2012, tab. S4). Assignment of samples to corresponding biozones was done after Sun et al. (2012) with a few corrections (see main text). C1, *Neogondolella changxingensis* Zone; C2, *Neogondolella yini* Zone; C3, *Neogondolella meishanensis* Zone; C4, *Hindeodus changxingensis* Zone; C5, *Hindeodus parvus* Zone; C6, *Isarcicella staeschei* Zone; C11; *Parachirognathus* Zone; C12, *Novispathodus pingdingshanensis* Zone; C13, *Icriospathodus collinsoni* Zone; C14, *Triassospathodus homeri* Zone; C15, *Chiosella timorensis* Zone; C16, *Neogondolella regalis* Zone; C17, *Nicoraella germanica* Zone; C18, *Nicoraella kockeli* Zone. The arrows with circled numbers 1 to 10 correspond to the absolute time constraints (see main text).

Galfetti et al., 2007).

(8) An age of 247.4 Ma for the FO of *Chiosella timorensis* in Guandao, South China (age by Lehrman et al., 2007, samples PGD1 and PGD2; revision of conodont ranges in Guandao by Goudemand et al., 2012).

(9) An age of 247.2 Ma for the Early-Middle Triassic Boundary (EMTB, Lehrmann et al., 2007). Note that, since *Chiosella timorensis* has been found in unambiguous Spathian strata, the *C. timorensis* Zone (sensu Sun et al., 2012, that is, based on the FAD of *C. timorensis*) actually straddles this boundary (Goudemand et al., 2012). Except for the position of the FAD of *C. timorensis*, the new EMTB definition (Goudemand et al., 2012) is fully compatible with the previous one (e.g. Lehrmann et al., 2007) and correlates roughly with the lowest part of the maximum $\delta^{13}C_{cath}$ plateau.

(10) An age of 246.8 Ma within the *Platycuccoceras* beds (Ovtcharova et al., 2006; sample CHIN29), and associated with the conodont *Nicoraella kockeli* (Lehrmann et al., 2006; sample GDGB-110; see also Galfetti et al., 2007, fig. 4), and hence in the *N. kockeli* Zone (C18, Fig. 2).

Unless constrained by absolute ages, the relative durations of biozones were set to reflect their relative stratigraphical thicknesses in the main section of the corresponding time interval (C1-C8 at Meishan, Joachimski et al., 2012; Zhang et al., 2007; Xie et al., 2007; Cao et al., 2010; C9-C13 at Zuodeng II section, Sun et al., 2012; C15-C18 at Guandao II (Upper Guandao) section, Lehrmann et al., 2007; Orchard et al., 2007). This is a somewhat arbitrary choice since it is unknown whether these thicknesses are a linear function of time. This would be the case if sedimentation rates were constant and no hiatuses were present but this is not necessarily true here (Joachimski et al., 2012; Sun et al., 2012, and references therein). Note that setting a sub-series of unconstrained biozones to be of equal duration would be an equally justified and equally arbitrary option.

The new PTB ages published by Burgess et al. (2014) lead to shorten the duration of the Dienerian considerably. This suggests that the early Smithian and possibly the early Spathian ages by Ovtcharova et al. (2006) should be reassessed too, using the same procedure as Burgess et al. (2014).

CORRELATION OF ADDITIONAL DATA

The data from Shangsi, Jinya, Jiarong, Wantuo, Bianyang Quarry, and Bianyang Hill sections were added using the biochronological correlations and, when available, the chemostratigraphical information. Additionally, if within a given biozone a series of $\delta^{13}C_{carb}$ points were associated with the $\delta^{18}O_{apatite}$ data, we adjusted the position of the sample series as a whole in order to 'fit' the Meishan-ZuodengII-GuandaoII composite $\delta^{13}C_{carb}$ curve. It is possible to do so because it has been shown that the trends of the $\delta^{13}C_{carb}$ evolution are the same within the basin (and globally). Yet, caution is warranted as the absolute $\delta^{13}C$ values and the amplitude of excursions do largely differ even within the same basin from one facies to another (see Meyer et al., 2011).

On the other hand, one could hardly justify adjusting the position of the series (as a whole or otherwise) to 'fit' the $\delta^{18}O_{apatite}$ 'curve'. In particular, isolated data points for which the associated $\delta^{13}C_{carb}$ data did not help constraining their position on the time axis were arranged so as to reflect the uncertainty on their temporal position: unless further information was provided, single points were placed at the center of the corresponding biozone, the topmost sample of a set of points was placed at the top of the biozone and the bottommost sample at the base of that biozone.

The early Smithian samples from Zuodeng I section (ZDC29, ZDC30) could not be positioned reliably on the time axis. They may or may not confirm a gradual warming within that time interval (compare with Sun et al., 2012). Similarly, it is hard to correlate sample ZDC24 (Sun et al. 2012, reported it from the *Neogondolella carinata* Zone but Sun et al. did not define a *Ng. carinata* Zone; we assigned it to the *Ng. planata* Zone): the associated $\delta^{13}C_{carb}$ data suggest that it is coeval to the minimum $\delta^{13}C_{carb}$ plateau in the middle of the *Ng. planata* Zone.

Sun et al.'s (2012) $\delta^{13}C_{carb}$ data from Jinya is consistent with Galfetti et al.'s (2007) data from Jinya if the early Smithian sample (LRC01) is considered younger than absolute age 5.

Sun et al. (2012) mentioned (their table S4; Sun, personal communication, 2013) that the samples at the Smithian-Spathian boundary from Bianyang Quarry section were ramiform elements, mostly robust ones. If P1 elements were present, they belonged to Parachirognathus spp (an ellisonid conodont). Ellisonids are not known from the Spathian (see for instance Orchard, 2007 for the latest review about Early Triassic conodonts). In my abundant collections from Guangxi, South China, ellisonids are clearly the dominant group in the middle-late Smithian Owenites koeneni beds (Galfetti et al., 2007). Yet, none of them survived the Smithian-Spathian boundary crisis. In other words, none is known in the Novispathodus pingdingshanensis Zone or in younger strata. Hence, it is doubtful that BYC19-1 (which contains Parachirognathus spp.) and BYC16-2 should be displayed in the Nv. pingdingshanensis Zone as in Sun et al.'s (2012) figure 2. Instead, I consider samples BYC19-1, BYC16-2, BYC14.3, BYB5 and BYB4 as belonging to the Parachirognathus Zone. It may sound trivial to discuss the location of these four points, but displaying them as straddling the Smithian-Spathian boundary gives the (misleading) impression that the evolution of $\delta^{18}O_{_{apatite}}$ at Bianyang section parallels that at Zuodeng II section. The evidence for such a parallel (offset) evolution of $\delta^{18}O_{anatise}$ in both sections is not available yet.

Figure 2 summarizes the results concerning the time calibration of Sun et al.'s (2012) $\delta^{13}C_{carb}$ and $\delta^{18}O_{anatise}$ data.

SST CURVE

 $\delta^{18}O_{apatite}$ data may be interpreted in terms of seawater temperatures, and, more precisely, in terms of Sea Surface Temperatures (SST). Yet, $\delta^{18}O_{apatite}$ data may also reflect the different life habits (water depth preferences) of the various conodont genera that were sampled and the corresponding



Figure 3 – Temporal evolution of δ 13Ccarb and estimated sea surface temperatures at Meishan, Zuodeng, Wantuo, Bianyang Hill, and Guandao II sections. For clarity, only the δ ¹³C_{carb} data is displayed for Guandao II because its δ ¹⁸O_{apatite} data is offset from the main trend (see main text and Fig. 4). Data modified after Sun et al. (2012) as described in main text. Samples using hindeodid or ellisonid (*Platyvillosus, Pachycladina, Parachirognathus*) material are considered as recording the SST. Samples using neospathodid material are corrected by a +3°C offset in order to retrieve the SST. Similarly, samples using neospatholellid material are corrected by a +4.7°C offset. The color bands show the first-order trends.

biases should be corrected for in order to retrieve the SST. Let us first focus on a main composite Meishan-Zuodeng II-Wantuo/ Bianyang Hill curve. In this case, Wantuo and Bianyang Hill sections are preferred to Guandao II because, contrary to Guandao II whose data is clearly offset, the $\delta^{18}O_{apatite}$ data from Wantuo and Bianyang Hill is compatible with the data from Zuodeng II. Without comparative samples, one can assume that the data from Meishan is not offset compared to that from Zuodeng II.

The original Meishan $\delta^{18}O_{apatite}$ data is from Joachimski et al. (2012) and the data points were obtained by processing either *Hindeodus* or *'Clarkina'* (*Neogondolella*) material. For the lower (Permian) part, up to bed 25, both *'Clarkina'* and *Hindeodus* are interpreted as recording $\delta^{18}O_{apatite}$ data that correspond to the SST (Joachimski et al., 2012, p. 196). Above bed 25, and concurrently with the PTB sea level rise, *'Clarkina'* records lower temperatures than *Hindeodus*, which is interpreted as *'Clarkina'* migrating to deeper and colder waters while *Hindeodus* remains at the sea surface (ibid).

For the Zuodeng II section, most measurements were performed with material comprising neospathodid elements (Sun et al., 2012, tab. S4). Based on two comparative analyses of Neospathodus and Platyvillosus (Sun et al., 2012, tab. S1, samples ZDC44 and ZDC B), Sun et al. (2012, supp. text, p. 7) interpreted neospathodids as having lived at a water depth of ca. 70 meters and *Platyvillosus* at sea surface. The case of the middlelate Smithian ellisonids (Pachycladina, Parachirognathus) is more difficult: ellisonids favored shallow-water environments (e.g. Carr et al., 1984) and their dominance in middle-late Smithian samples may reflect the recorded coeval regression. Yet, without comparative analyses from the same samples, it is not clear by how much their $\delta^{18}O_{_{apatite}}$ signal should be offset relative to that of neospathodids. In the following we shall assume, as Sun et al. (2012), that ellisonids lived at sea surface and their $\delta^{18}O_{anarite}$ offsets by -0.7‰ compared to that of neospathodids.

The Wantuo and Bianyang Hill data was obtained using either neospathodid or unidentified ramiform material (Sun et al., 2012, tab. S4). Without further evidence, we shall assume most samples correspond to animals that lived like neospathodids at a water depth of ca. 70 meters. One sample from Wantuo (WTC05) was used to compare neospathodid and neogondolellid material and the corresponding 0.4‰ offset suggests that neogondolellids lived at a water depth of ca. 85 meters (Sun et al., 2012, tab. S1 and supp. text p. 7).

The Zuodeng I section was measured using either neospathodids (ZDC28-31), neogondolellids (ZDC24 and ZDC32) or hindeodids (ZDC21). The same offsets as above apply. Namely: samples using hindeodid or ellisonid (*Platyvillosus, Pachycladina, Parachirognathus*) material are considered as recording the SST; samples using neospathodid material are corrected by a +3°C offset in order to retrieve the SST; and similarly, samples using neogondolellid material are corrected by a +4.7°C offset.

The SST curve for the 'Meishan-Zuodeng-Wantuo/Bianyang Hill' composite is shown on Figure 3. The temperatures are computed using the equation of Pucéat et al. (2010), assuming constant salinity and an ice-free world ($\delta^{18}O_{seawater} = -1\%$) (Sun et al., 2012).

On Figure 4, the same procedure is applied to the data from Guandao II, Bianyang Quarry, Jinya and Jiarong sections. The evolution of the SST in Guandao II section is clearly offset (by about -8°C) from that in Wantuo, Bianyang Hill and Zuodeng II sections. Sun et al. (2012) proposed that it is due to higher evaporation in Guandao (salinity offset) since this section is close to the Great Bank of Guizhou. The few data points from Jinya fit with the temperatures in Zuodeng for the early Smithian and the early Spathian but they differ within the Parachirognathus Zone. The evolution of the SST in Jiarong seems to parallel that in Zuodeng but it is offset by 2-4°C. Since we are comparing sea surface temperatures, this offset cannot be explained by a difference in water depth (contra Sun et al., 2012). It could instead reflect regional variation but it is puzzling that the offset in Jiarong is opposite to that in Guandao although Jiarong and Guandao are neighboring sections. Sun et al. (2012, fig. 3) suggested that the δ^{18} O data from Jiarong, Bianyang and Jinya around the Smithian-Spathian boundary, was reflecting that of Tsoteng and was recording the temporal evolution of the SST: namely the occurrence of a presumed late Smithian thermal maximum.

CONCLUSION

Though based on the same assumptions as Sun et al. (2012) about the life habitats of the measured conodonts, our new calibration offers a significantly different picture for the Early Triassic evolution of sea surface temperatures in South China: the late Smithian *Parachirognathus* Zone may in fact correspond to a decrease of temperatures, while the early Spathian *Novispathodus pingdingshanensis* Zone would correspond to a rebound of temperatures, possibly with a peak in the earliest Spathian. The sea surface temperatures drop again during the *Icriospathodus collinsoni* Zone and then they rise until the middle and late Spathian where they reach levels similar to those observed in the middle Smithian and early Spathian.

ACKNOWLEDGEMENTS

Albertiana's chief editor Chris McRoberts and reviewer Margaret Fraiser are thanked for their comments, which helped to improve this manuscript.

REFERENCES

- Brayard A. & Bucher H. 2008. New Smithian (Early Triassic) ammonoid faunas from northwestern Guangxi (South China): taxonomy and biochronology. Fossils & Strata, 55 (= Lethaia, 41, S2): 179 pp.
- Brühwiler T., Ware D., Bucher H., Krystyn L. & Goudemand N. 2010. New Early Triassic ammonoid faunas from the Dienerian/Smithian boundary beds at the Induan/ Olenekian GSSP candidate at Mud (Spiti, Northern India). Journal of Asian Earth Sciences, 39(6): 724–739.



Figure 4 Temporal evolution of $\delta^{13}C_{carb}$ and estimated sea surface temperatures at Meishan, Shangsi, Zuodeng, Jinya, Jiarong, Bianyang Quarry, Wantuo, Bianyang Hill and Guandao II sections. Data modified after Sun et al. (2012), see main text and Fig. 3. The SST in Jiarong and Guandao are offset in the opposite way relative to the main composite SST curve.

Burgess, S.D., Bowring, S. & Shen, S.z. 2014. High-precision timeline for Earth's most severe extinction. Proceedings of the National Academy of Sciences, 111(9): 3316–3321.

Cao, C, Yang, Y.C., Shen, S.Z., Wang, W., Zheng, Q.F. & Summons, R.E. 2009. Biogeochemical evidence for euxinic oceans and ecological disturbance presaging the end-Permian mass extinction event. Earth and Planetary Science Letters, 281(3-4): 188–201.

Galfetti, T., Bucher, H., Ovtcharova, M., Schaltegger, U., Brayard, A., Brühwiler, T., Goudemand, N., Weissert, H., Hochuli, P.A., Cordey, F. & Guodun, K. 2007. Timing of the Early Triassic carbon cycle perturbations inferred from new U-Pb ages and ammonoid biochronozones. Earth and Planetary Science Letters, 258: 593–604.

Goudemand, N. 2014. Note on the conodonts from the Induan-Olenekian boundary. Albertiana, 42: 49–51.

Goudemand, N., Orchard. M.J., Bucher, H. & Jenks, J. 2012.The elusive origin of *Chiosella timorensis* (Conodonts, Triassic). Geobios, 45: 199–207.

Goudemand, N., Romano, C., Brayard, A., Hochuli, P.A. & Bucher, H. 2013. Comment on "Lethally hot temperatures during the Early Triassic greenhouse". Science, 339: 1033.

Joachimski, M.M., Lai, X., Shen, S., Jiang, H., Luo, G., Chen, B., Chen, J. & Sun, Y. 2012. Climate warming in the latest Permian and the Permian–Triassic mass extinction. Geology, 40: 195–198.

Lehrmann, D.J., Ramezani, J., Bowring, S.A., Martin, M.W., Montgomery, P., Enos, P., Payne, J.L., Orchard, M.J., Wang, H. & Wei, J. 2006. Timing of recovery from the end-Permian extinction: geochronologic and biostratigraphic constraints from South China. Geology, 34: 1053–1056.

Lehrmann, D.J., Ramezani, J., Bowring, S.A., Martin, M.W., Montgomery, P., Enos, P., Payne, J.L., Orchard, M.J., Wang, H. & Wei, J. 2007. Timing of recovery from the end-Permian extinction: geochronologic and biostratigraphic constraints from South China: reply. Geology, 35: e136–e137.

Meyer, K.M., Yu, M., Jost, A.B. & Payne, J.L. 2011. Carbon isotope evidence for a vigorous biological pump in the aftermath of end-Permian mass extinction. Earth and Planetary Science Letters, 302(3-4): 378–384.

Orchard, M.J. 2007. Conodont diversity and evolution through the latest Permian and Early Triassic upheavals. Palaeogeography, Palaeoclimatology, Palaeoecology, 252: 93–117. Orchard, M.J., Lehrmann, D.J., Wei, J., Wang, H. & Taylor, H.J. 2007. Conodonts from the Olenekian-Anisian boundary beds, Guandao, Guizhou Province, China. *In* Lucas, S.G. & Spielmann, J.A. (eds.), The Global Triassic. New Mexico Museum of Natural History and Science Bulletin, 41: 347–354.

Ovtcharova, M., Bucher, H., Schaltegger, U., Galfetti, T., Brayard, A. & Guex, J. 2006. New Early to Middle Triassic U–Pb ages from South China: calibration with ammonoid biochronozones and implications for the timing of the Triassic biotic recovery. Earth and Planetary Science Letters, 243: 463–475.

Pucéat, E., Joachimski, M.M., Bouilloux, A., Monna, F., Bonin, A, Motreuil, S., Morinière, P., Hénard, S., Mourin, J., Dera, G. & Quesne, D. 2010. Revised phosphatewater fractionation equation reassessing paleotemperatures derived from biogenic apatite. Earth and Planetary Science Letters, 298: 135–142.

Sun, Y., Joachimski, M.M., Wignall, P.B., Yan, C., Chen, Y., Jiang, H., Wang, L. & Lai, X. 2012. Lethally hot temperatures during the Early Triassic greenhouse. Science, 338: 366–370.

Sun, Y., Joachimski, M.M., Wignall, P.B., Yan, C., Chen, Y., Jiang, H., Wang, L. & Lai, X. 2013. Response to Comment on "Lethally hot temperatures during the Early Triassic greenhouse". Science 339: 1033-d.

Tong, J., Zuo, J. & Chen, Z.Q. 2007. Early Triassic carbon isotope excursions from South China: proxies for devastation and restoration of marine ecosystems following the end-Permian mass extinction. Geological Journal, 42: 371–389.

Xie, S., Pancost, R.D., Huang, X., Wignall, P., Yu, J., Tang, X., Chen, L., Huang, X. & Lai, X. 2007. Changes in the global carbon cycle as two episodes during the Permian-Triassic crisis. Geology, 35: 1083–1086.

Zhang, K., Tong, J., Shi, G.R., Lai, X., Yu, J., He, W., Peng, Y. & Jin, Y. 2007. Early Triassic conodont–palynological biostratigraphy of the Meishan D Section in Changxing, Zhejiang Province, South China. Palaeogeography, Palaeoclimatology, Palaeoecology, 252: 4–23.

Research Note

NOTE ON THE CONODONTS FROM THE INDUAN-OLENEKIAN BOUNDARY

Nicolas Goudemand

Paleontological Institute and Museum, University of Zurich, Karl-Schmid-Strasse 4, CH-8006 Zurich, Switzerland, Email: nicolas.goudemand@pim.uzh.ch

INTRODUCTION

A brief summary of results of studies into the Induan-Olenekian Boundary (IOB) conodont material from Waili (Guangxi Province, South China) and Mud (Spiti, Himashal Pradesh, Northern India) is presented (Goudemand et al., in prep. a, b). The latest consensus about the definition of the IOB was to use the FAD of Neospathodus waageni sensu lato as an index for the base of the Olenekian (IOB Working Group, as suggested by Tong et al., 2004). Up to now and until the foreseen proposal of the Nammal Gorge section, Pakistan as a GSSP candidate, only two sections have been formally proposed: the Mud section M04 in Northern India (Krystyn et al., 2007a, b), and the West Pindingshan section at Chaohu in Anhui Province, South China (Tong et al., 2003, 2004). Conodont-based correlations between these two sections remain hitherto unclear. Furthermore, new ammonoid data recently questioned the stratigraphic position of the IOB in Mud (Brühwiler et al., 2010). Our new conodont data demonstrates that the northern Indian and south Chinese conodont sequences agree better with one another and with the ammonoid data than was previously thought.

NEW DATA

In the Mud section, the presumed FO of *Neospathodus* waageni was reported in subbed 13A3 (Krystyn et al., 2007a, b). In order to re-assess in details the Mud section, intense bedby-bed sampling of ammonoids and conodonts was performed below this proposed GSSP. It led first to a revision of the ammonoid biozonation and to the recognition that the most relevant ammonoid turnover occurred between beds 9 and 10 (Brühwiler et al., 2010), about 1m below bed 13 where the boundary was proposed in 2007 (Krystyn et al., 2007a, b). The conodont material led to surprising outcomes too, since 'typical' Smithian forms occurred also below bed 13 (Goudemand et al., in prep. a). In particular, Ns. waageni sensu lato was also found in bed 10 (together with the ammonoid *Flemingites bhargavai*), the middle part of which had not been sampled previously. In bed 10 of Mud section, the elements pertaining to Ns. waageni sensu lato (see Fig. 1.A) are similar to the specimens of "Ns. waageni eowaageni" that were described from beds 24-16, 24-20, 24-21 and 24-22 at Chaohu (Zhao et al., 2008a, fig. 3/1-4). Neospathodus waageni is a highly variable species that deserves further species differentiation (Goudemand et al., in prep. a, b). These early forms of "Ns. waageni eowaageni" were hitherto included in Ns. waageni sensu lato and discussed as such during earlier deliberations of the IOBWG. The fact that "Ns. waageni eowaageni" had not been recorded from Mud until now had actually led some people to question the completeness of the conodont record in Mud. Our results show that they do occur in Mud too and that they could be used to define the IOB. However, these early forms, which I will refer here to Novispathodus waageni new subspecies A, should be distinguished from true Ns. waageni eowaageni, a subspecies of 'Neospathodus' waageni whose holotype is from higher up in the Chaohu section (bed 27 at Chaohu, Zhao et al., 2007, pl.1, fig.5; Zhao et al., 2008a, fig. 3/5; not pl. 1, fig. 11 from Zhao et al., 2008b, which was published later). Though the two forms are superficially similar, Ns. waageni eowaageni is about twice as large and its upper profile is more arcuate. I have many specimens of both forms in my collections (e.g., Waili and Mud) and Novispathodus waageni new subspecies A cannot be considered a juvenile form of Ns. waageni eowaageni.

Note also that I assign them to the genus *Novispathodus*, whose type species is *Novispathodus abruptus*, an early Spathian species. The P₁ element of *Novispathodus* is segminate and very similar to the P₁ of all neospathodids, but the other elements of the apparatus are different from those of *Neospathodus*, especially the P₂ element, which is characteristically high-bladed. In collections from both Waili and Mud, the P₁s of *Novispathodus waageni* new subspecies A are found together with similarly denticulated, high-bladed P₂s, which in my opinion belong also to *Novispathodus waageni* new subspecies A (see Fig. 1.B). Based on further unpublished reconstructions of its apparatus, *Ns. waageni* should be similarly re- assigned to the genus *Novispathodus* (e.g., Orchard & Zonneveld, 2009). This may apply also to all species associated with *Nv. waageni*, such as '*Neospathodus' pakistanensis* and '*Neospathodus' posterolongatus*.

At Chaohu, further elements that Zhao et al. (2008a) assigned to '*Neospathodus*' posterolongatus were also illustrated from the first 'Olenekian' beds containing *Novispathodus* waageni new subspecies A (Zhao et al., 2008a, beds 24-20 and 24-22, fig. 3/7, 8). Novispathodus posterolongatus was originally



Figure 1 – *A*–*D*, *Novispathodus waageni* new subspecies A. *A*, P₁ element, Mud, bed 10. *B*, P₂ element, Mud, bed 10. *C*, *D*, P₁ elements, Waili cave section, bed W147 (=bed W235 of Galfetti et al., 2008). *E*, *Novispathodus* n. sp. B. P₁ element, Mud, bed 10. All SEM images, all × 80. All specimens are deposited at the Paleontological Institute and Museum, University of Zurich, Switzerland.

considered a subspecies of *Nv. waageni*. It is also considered as a good secondary proxy for the IOB (IOB Working Group). *Novispathodus posterolongatus* is similar to *Nv. pakistanensis* but differs in the elongated basal cavity of its P₁ element. The holotype of *Nv. posterolongatus* comes from bed 25-25 at Chaohu (Zhao et al., 2007), where it co-occurs with *Nv. waageni waageni*. In Chaohu, Waili and Mud, the FOs of *Nv. posterolongatus* and *Nv. waageni* sensu stricto are sub-contemporaneous. In my opinion, the elements illustrated by Zhao et al. from beds 24-20 and 24-22 are not true *Nv. posterolongatus* and are better differentiated as a new taxon (Goudemand et al., in prep. a, b) that I refer here to *Novispathodus* n. sp. B. In all cases, the very same elements are also found to co-occur with *Novispathodus waageni* new subspecies A in collections from Mud and Waili (see Figs. 1.C–E) and they could also be used as an index for the IOB.

Based on these results, bed 10 in Mud correlates with the upper part of bed 24 in Chaohu (West Pingdingshan section). In accordance with the Chaohu section and the ammonoid data from Mud (Brühwiler et al., 2010), the IOB in Mud should be placed between beds 9 and 10.

The biostratigraphic sequences are very similar in Mud, Waili and Chaohu. In Mud, we find successively: Novispathodus waageni new subspecies A and Novispathodus n. sp. B (bed 10), Borinella nepalensis (from bed 11), Eurygnathodus costatus and Eu. hamadai (beds 12, 13), N. waageni eowaageni (from bed 14). In the Waili cave section in Guangxi, the same sequence is recognized: Novispathodus waageni new subspecies A is found (see Figs. 1C-D) about 1 meter below the massive 'Flemingites rursiradiatus' beds (Brayard and Bucher, 2008), Borinella. cf. B. nepalensis was found 30 cm above that, Eu. costatus and Eu. hamadai are restricted to the bed just below the Flemingites beds and the lowermost part of the Flemingites beds, and Nv. waageni eowaageni is found in the middle of the Flemingites beds. In Chaohu, and as far as one can tell from the published material, a similar sequence occurs, except that no element of Borinella is known. Finally, note that at Nammal section (Salt Range, Pakistan), the type locality of Nv. waageni waageni, P. and P2 elements of Novispathodus waageni new subspecies A are also found in the earliest 'Smithian' strata (Goudemand et al., unpublished), much lower than the FO of Nv. waageni sensu stricto.

CONCLUSIONS

Based on our new data from Mud and Waili, we differentiate two new taxa (*Novispathodus waageni* new subspecies A and *Novispathodus* n. sp. B) that were previously thought to belong to *Nv. waageni eowaageni* and *Nv. posterolongatus* respectively. In Chaohu these two taxa occur in the earliest Smithian strata (sensu IOBWG). In Mud and Waili (and Nammal), they can also be used to define the base of the IOB. Such definition would be compatible with the ammonoid data from Mud. The conodont biostratigraphic sequences in Mud, Waili, and Chaohu are very similar.

REFERENCES

- Brayard, A. & Bucher, H. 2008. New Smithian (Early Triassic) ammonoid faunas from northwestern Guangxi (South China): taxonomy and biochronology. Fossils & Strata, 55 (= Lethaia 41 S2): 179 pp.
- Brühwiler, T., Ware, D., Bucher, H., Krystyn, L. & Goudemand, N. 2010. New Early Triassic ammonoid faunas from the Dienerian/Smithian boundary beds at the Induan/Olenekian GSSP candidate at Mud (Spiti, Northern India). Journal of Asian Earth Sciences, 39(6): 724–739.
- Galfetti, T., Bucher, H., Martini, R., Hochuli, P.A., Weissert, H., Crasquin-Soleau, S., Brayard, A., Goudemand, N., Brühwiler, T. & Kuang, G. 2008. Evolution of Early Triassic outer platform paleoenvironments in the Nanpangjiang Basin (South China) and their significance for the biotic recovery. Sedimentary Geology, 204: 36–60.
- Goudemand, N., Orchard, M., Krystyn, L., Brühwiler, T., Ware, D. & Bucher, H. in prep. a. New Early Triassic conodont faunas from the Dienerian/Smithian boundary beds at Mud (Himashal Pradesh, India).
- Goudemand, N., Galfetti, T., Brayard, A. & Bucher, H. in prep. b. New Early Triassic conodont faunas from the Dienerian/Smithian boundary beds at Waili (Guangxi, China).
- Krystyn, L., Bhargava, O.N. & Richoz S. 2007a. A candidate GSSP for the base of the Olenekian Stage: Mud at Pin

Valley; district Lahul & Spiti, Himachal Pradesh (Western Himalaya), India. Albertiana, 35: 5–29.

- Krystyn, L., Richoz, S. & Bhargava, O.N. 2007b. The Induan-Olenekian Boundary (IOB) in Mud – an update of the candidate GSSP section M04. Albertiana, 36: 33–45.
- Orchard, M.J. & Zonnenfeld, J.P. 2009. The Lower Triassic Sulphur Mountain Formation in the Wapiti Lake area: lithostratigraphy, conodont biostratigraphy and a new biozonation for the Lower Olenekian (Smithian). Canadian Journal of Earth Science, 46: 757–790.
- Tong, J., Zakharov, Y.D., Orchard, M.J., Yin, H. & Hansen, H.J. 2003. A Candidate of the Induan-Olenekian boundary stratotype in the Tethyan Region. Science in China (series D), 46(11): 1182–1200.
- Tong, J., Zakharov, Y.D., Orchard, M.J., Yin, H. & Hansen, H.J. 2004. Proposal of Chaohu Section as the GSSP Candidate of the Induan-Olenekian Boundary. Albertiana, 29: 13–28.

- Zhao, L., Orchard, M.J., Jinnan, T., Zhiming, S., Jinxun, Z., Suxin, Z. & Ailing, Y. 2007. Lower Triassic conodont sequence in Chaohu, Anhui Province, China and its global correlation. Palaeogeography, Palaeoclimatology, Palaeoecology, 252: 24–38.
- Zhao, L., Tong, J., Sun, Z. & Orchard, M.J. 2008a. A detailed Lower Triassic conodont biostratigraphy and its implications for the GSSP candidate of the Induan– Olenekian boundary in Chaohu, Anhui Province. Progress in Natural Science, 18: 79–90.
- Zhao, L., Tong, J., Zhang, S. & Sun, Z. 2008b. An update of conodonts in the Induan-Olenekian boundary strata at West Pingdingshan Section, Chaohu, Anhui Province. Journal of China University of Geosciences, 19(3): 207–216.



STS Task Group Report

NEW AND CORRECTED EVIDENCES ON EARLY TRIASSIC BIOSTRATIGRAPHY IN SOUTH PRIMORYE, SIBERIA, KAZAKHSTAN AND VIETNAM (REPORT ON THE IOBWG ACTIVITY IN 2014)

Yuri D. Zakharov

Chair, Induan-Olenekian Task Group.

Far Eastern Geological Institute, Russian Academy of Sciences (Far Eastern Branch), Stoletiya, Prospect 159, Vladivostok 690022, Russia,

MAIN RESULTS

1. Global correlation of the Permian-Triassic boundary sequences (*Otoceras* beds) in Siberia has been made on the basis of first carbon-isotope and known palaeontological data (Zakharov et al., 2014a).

2. The integrated approach in palaeontological investigations of the Olenekian (Lower Triassic) of Northern Vietnam was used for the first time: ammonoids (25 species), gastropods (eight species), bivalves (five species), conodonts (43 species), ostracodes (six species, including one new one), and radiolarians (12 species, including five new ones) (Shigeta et al., 2014). As a result, some new subdivisions are recognized in the middle Smithian (*Flemingites rursiradiatus, Urdyceras tulongensis*, and *Owenites koeneni* Beds), upper Smithian (*Xenoceltites variocostatus* Beds), and lower Spathian (*Tirolites* cf. *cassianus* and *T*. sp. nov. Beds). Conodonts *Novispathodus* ex gr. *waageni* were found in the middle Smithian in Vietnam.

3. The corrected data on distribution of invertebrates (brachiopods, ammonoids) in the Induan and Olenekian of South Primorye and the Olenekian of Kazakhstan (Mangyshlak) have been obtained (Zakharov and Popov, 2014).

4. First description of well-preserved foraminifera (*Ammodiscus* sp.) from the Lower Triassic (Smithian) of South Primorye was made for the Russian Far East (Bondarenko et al., 2014).

5. Ammonoids of the genus *Churkites*, typical representative of the early Olenekian assemblage (*Anasibirites nevolini* Zone), have been discovered in the upper part of the Tri Kamnya Cape section at the western coast of Ussuri Gulf in South Primorye (Fig. 1) for the first time. *Churkites* cf. syaskoi was met in association with ammonoids *Inyoites*, *Clypeoceras*, *Owenites*, *Juvenites*, *Prionites*, *Glyptophiceras*, *Brayardites*, *Mianwaliites*, *Pseudoflemingites*, *Aspenites*, *Rohillites*, *Pseudoaspedites*, *Anaxenaspis*, *Anasibirites*, and *Arctoceras*, most of which are known from the basic section for the *Anasibirites nevolini* Zone (SMID, Artyom). The rarity of ammonoids of the genus *Anasibirites* within the investigated interval and the lack of platform conodonts, including *Schythogondolella milleri* (Müller), within isolated material seem

to be an evidence for the property of fossil-bearing sediments to the upper part of the *Anasibirites nevolini* Zone (upper part of the Smithian Substage of the Olenekian). The lower part of the *Anasibirites nevolini* Zone in South Primorye, correlated with the *S. milleri* conodont Zone, more likely corresponds to the *Wasatchites clistractus* Zone of the Salt Range in Pakistan (Zakharov et al., 2014b).

6. New species of flemingitid ammonoids from the Smithian of South Primorye have been described (Smyshlyaeva and Zakharov, 2013).

Field-work results

Tirolites-bearing sediments located just above the Smithian were discovered in the Kamenushka River basin in South Primorye (new Kamenushka-1 and Kamenushka-2 sections).

Meetings

1. Cephalopod Symposium (Zurich, Sept. 2014). There was business-meeting on the Triassic subcommission in Zurich. Only a single question has been discussed: Induan-Olenekian boundary (IOB). Yuri Zakharov informed about unpublished carbon isotope results, obtained on this topic in Primorye region. Hugo Bucher promised to prepare a presentation on the candidate of global stratotype section of the IOB (Nammal, Salt Range, Pakistan) in nearest feature, when conodonts from the Nammal section will be described.

2. 4th International Palaeontological Congress (Mendoza, Sept. 28-Oct. 3, 2014). The poster "Global correlation of *Otoceras* beds in light of the first carbon isotope data on Permian-Triassic boundary in Siberia" (Y.D. Zakharov, A. S. Biakov, S. Richoz, and M. Horachek) was presented.

REFERENCES

Bondarenko, L.G., Zakharov Y.D. & Barinov, N.N. 2014. The first find of well-preserved foraminifera in the lower Triassic of Russian Far East. Albertiana, 41: 34–38.

Shigeta, Y., Komatsu, T., Mackawa, T. & Tran, H.D. 2014.



Figure 1 - Image of the Induan/Olenkian boundary Tri Kamnya Cape section at the western coast of Ussuri Gulf in South Primorye

Olenekian (Early Triassic) stratigraphy and fossil assemblage in northestern Vietnam. National Museum of Nature and Science, Tokyo. 309 pp.

- Smyshlyaeva, O.P. & Zakharov, Y.D. 2013. New members of the family Flemingitidae (Ammonoidea) from the Lower Triassic of South Primorye. Paleontological Journal, 47(3): 247–255.
- Zakharov Y.D., Biakov A.S. & Horacek, M. 2014a. Global correlation of basal layers in the Permian-Triassic boundary in Northeast Asia. Russian Journal of Pacific Geology, 8(1): 1–17.
- Zakharov, Y.D., Bondarenko, L.G. & Popov, A.M. 2014b. Lower Triassic stratigraphic units of Southern Primorye. Paper 1: first records of ammonoids of the genus *Churkites* on the coast of the Ussuri Gulf. Russian Journal of Pacific Geology, 8(6): 391–403.
- Zakharov, Y.D. & Popov, A.M. 2014. Recovery of brachiopod and ammonoid faunas following the end-Permian crisis: additional evidence from the Lower Triassic of the Russian Far East and Kazakhstan. Journal of Earth Science, 27(1): 1–44.

Obituary

CARMINA VIRGILI (1927–2014)

Carmina Virgili died in Barcelona on the 21st of November, 2014 where she was born 87 years ago.

During her academic career she lectured in Stratigraphy at Barcelona (1949-1963), Oviedo (1963-1968) and Madrid (1968-1996).

She was a pioneer on the modern studies of the Permian and Triassic in Spain, starting with her Ph.D. Thesis (1956), in which she made the first correct interpretation of the Triassic successions of the Catalan Coastal Ranges and remains the reference work for the Triassic of Northeastern and Central Spain.

In 1968, she created a research team at the Complutense University of Madrid dedicated to the Permian and Triassic stratigraphy and paleogeography, still active. Under her direction, fundamental new data were discovered: the presence of Permian sediments in the Iberian Ranges and Central System (Central Spain), the correct description and dating of Early Permian, Late Permian and Triassic sediments in Central and Eastern Spain, the interpretation of their mutual relationship, boundaries and subdivision into Formations and Members, changing radically the previously accepted interpretations, among may others. Her strong leadership ensured the activity of this group up to now, starting a network of correspondent researchers all over Europe that was at the origin of many collaborative projects.

She introduced in Spain the palynological techniques in collaboration with the Laboratories of the Universities of Strasbourg (France) and Utrecht (Holland) with great success, as all the main continental and shallow marine formations of the Permian and the Triassic were dated and the inter-basinal correlations were refined up to European standards. She encouraged many of the members of the research team to have long periods of study at French and English Universities, a very unusual practice in Spain in the 70's and 80's as part of her concept of international collaborative research effort.

She acted as the Spanish representative in several IUGS-UNESCO Research Programs and in the IUGS-UNESCO Sub-commissions on Permian and Triassic Stratigraphy.

She published more than one hundred scientific articles and books. At the end of her scientific career she became interested in the History of Geology and wrote a book on Lyell and several articles on the relationship of the english



geologist and Spain (See selected publication list below).

She was the first woman to be Professor (1963) and Dean of a Faculty (1977) after the Spanish Civil War. She was Junior Minister for Universities and Research (1982-1985, Dean of the College d'Espagne, Cité Universitaire, Paris (1987-1996) and Senator for Barcelona, Labour Party (1996-2000). She also was a strong militant of women's rights in the society fighting against enormous odds in a male-dominated academic world in the 50' and the 60'.

She will be dearly missed by her former students, disciples, friends and colleagues in different institutions.

Alfredo Arche Madrid, Spain

SELECTED PUBLICATIONS OF CARMINA VIRIGILI

- Virgili, C. 1952. Hallazgo de nuevos *Ceratites* en el Triásico Mallorquin. Memorias de la Comisión del Instituto Geológico Provincial de Barcelona, 9: 19–38.
- Virgili, C. 1953. Le Trías de la región d'Olesa (Catalogne, Espagne). Compte-rendu sommaire et bulletin de la Société Géologique de France, 3(6): 208–210.
- Virgili, C. & Julivert, M. 1954. El Triásico del la Sierra de Prades. Estudios Geológicos, 10: 216–242.
- Virgili, C. 1955. El tramo rojo intermedio del Muschelkalk de las Catalánides. Memorias de la Comisión del Instituto Geológico Provincial de Barcelona, 13: 37–78.
- Virgili, C. 1958. El Triásico de los Catalánides. Boletin del Instituto Geológico y Minero de España, 69, 856 pp.
- Virgili, C. & Rosell. J. 1959. Fauna Toarciense y Bajo- ciense en la Sierra de Prades (Tarragona). Notas y Comunicaciones del Instituto Geológico y Minero de España, 54: 131–137.
- Virgili, C. 1963. Le Trias du Nordest de l'Espagne. Boletin del Instituto Geológico y Minero de España, 15: 469–481.
- Virgili, C., Mary, G., Suarez Vega, L.C. & Brell, J.M., 1968. Depositos marinos del cuaternario antiguo(?) sobre la cuarcita armoricana del Cabo Torres (Gijon). Breviora Geologica Astiirica, 12(2): 4–8.
- Virgili, C., Suarez-Vega, L.C. & Rincon, R. 1971. La cobertera mesozoica de Asturias. I Congreso Hispano-Luso Americano de Geología. Económica, Madrid, 1: 461–468.
- Dubar, G., Mouterde, R., Suárez-Vega, L.C. & Virgili, C. 1971. El Jurásico de Asturias (norte de España). Cuadernos Geología Ibérica, 2: 561–580.
- Virgili, C., Santiago, H., Ramos, A. & Sopena, A. 1973. La sedimentation permienne au centre de l'Espagne. Compterendu sommaire et bulletin de la Société Géologique de France, 7(15): 109–112.
- Virgili, C. 1974. Vue d'ensemble sur les affleurements du Trias évaporitique espagnol de la bordure de la Meseta au littoral mediterraneen. Compte-rendu sommaire et bulletin de la Société Géologique de France, 7(16): 27.
- Virgili, C., Hernando, S., Ramos, A. & Sopeña, A. 1976. Le Permien en Espagne. *In*, Falke, H. (ed.), The continental Permian in Central, Western and Southern Europe, Brill, Leiden, pp. 91–109.
- Virgili, C., Sopeña, A., Ramos, A. & Hernando, S. 1977. Problemas de la cronoestratigrafia del Trias en Espana. Cuadernos Geología Ibérica, 77(4): 57–87.
- Virgili, C. & Esteban, M. 1977. Bibliografia del Trias de los Catalanides. Cuadernos Geología Ibérica, 77(4): 691–699.
- Virgili, C., Via, L., de Villalta, J.F., Paris, R., Esteban, M., Marzo, M., Colombo, F. & Anadon, P. 1977. Guia de la excursion al Triasico de la Sierra de Prades, Cuadernos Geología Ibérica, 77(4): 527–557.
- Virgili, C., 1977. Le Trias du Nord de l'Espagne. Bulletin due Bureau de Recherches Géologiques et Minieres 13, 4(3): 205–213.
- Mouterde, R., Suarez Vega, L.C. & Virgili, C. 1977. Sobre el limite Triasico-Jurasico en Asturias (Noroeste de Espana).

Cuadernos Geología Ibérica, 77(4): 325-328.

- Sopeña, A., Virgili, C., Doubinger, J. & Adloff, M.C. 1977. Excursion al Triasico y Permico de la Cordillera Iberica y bordes del Sistema Central; dia 4; El Permico y el Triasico del borde sureste del Sistema Central. Field trip for the Triassic and Permian of the Iberian Mountains and the borders, 77(4): 623–647.
- Virgili, C. 1980. Cartografía del Triásico y Pérmico del borde oriental del Sistema Central y Rama Castellana de la Cordillera Ibérica, Cuadernos Geología Ibérica, 6: 1–95.
- Virgili, C., Sopeña, A., Ramos, A., Hernando, S. & Arche, A. 1980. El Permico en España, Boletín Instituto Geológico y Minero, 12(2): 255–262.
- Virgili, C. 1983. El Permico en Espana: Introduccion. *In*, Diaz, C.M. (ed.), Carbonífero y Pérmico de España. Instituto Geológico y Minero de España. Madrid.: 383– 388.
- Virgili, C., Sopeña, A., Ramos, A., Arche, A. & Hernando, S. 1983. El relleno posthercinico y el comienzo de la sedimentacion mesozoica. *In*, Fontbote, J.M. (ed.), Geologia de España (Libro Julilar de J. M. Rios), Instituto Geológico y Minero de España, Madrid, 2: 25–36.
- Virgili, C., Sopeña, A., Arche, A., Ramos, A. & Hernando, S. 1983. Some observations on the Triassic of the Iberian Peninsula. Schriftenreihe der Erdwissenschaftlichen Kommissionen, 5: 287–294.
- Virgili, C., Doubinger, J., Arche, A., Sopena, A., Ramos, A. & Broutin, J. 1984. The Autunian of central Spain. Compte Rendu - Congres International de Stratigraphie et de Geologie du Carbonifere, 9(2): 593–598.
- Sopeña, A., Virgili, C., Arche, A., Ramos, A. & Hernando, S. 1985. El Triásico. *In*, Comba, J.A. (ed.), Geología de España. Instituto Geológico Minero y Minero de España, Madrid, 2: 47–62,
- Virgili, C. 1987. Problematica del Trias y Permico Superior del bloque Iberico. Cuadernos Geología Ibérica, 11: 39–52.
- Sopeña, A., López-Gómez, J., Arche, A., Pérez-Arlucea, M., Ramos, A., Virgili, C. & Hernando, S. 1988. Permian and Triassic Rift basins of the Iberian Peninsula. *In*, Manspeizer, W. (ed.), Triassic-Jurassic Rifting, Developments in Geotectonics, 22: 757–786, Elsevier, Amsterdam.
- Virgili, C. 2003. El fin de los mitos geológicos: Lyell. Nivola, Castellano. 308 pp.
- Virgili, C., Cassinis, G. & Broutin, J. 2006. Permian to Triassic sequences from selected continental areas of southwestern Europe. Geological Society Special Publications, 265: 231-259.
- Virgili, C. 2008. The Permian-Triassic transition: Historical review of the most important ecological crisis, with special emphasis on the Iberian Peninsula and Western-Central Europe. Journal Iberian Geology, 34: 123–158.

NEW TRIASSIC LITERATURE

Geoffrey Warrington

Honorary Visiting Fellow, Department of Geology, The University of Leicester, LE1 7RH, UK; Email: gwarrington@btinternet.com

- Abdala, F., Jasinoski, S.C. & Fernandez, V. 2013. Ontogeny of the Early Triassic cynodont *Thrinaxodon liorhinus* (Therapsida): dental morphology and replacement. Journal of Vertebrate Paleontology, 33(6): 1408–1431.
- Abdala, F., Marsicano, C.A., Smith, R.M.H. & Swart, R. 2013. Strengthening western Gondwanan correlations: a Brazilian dicynodont (Synapsida, Anomodontia) in the Middle Triassic of Namibia. Gondwana Research, 23(3): 1151–1162.
- Abrajevitch, A., Hori, R.S. & Kodama, K. 2013. Rock magnetic record of the Triassic-Jurassic transition in pelagic bedded chert of the Inuyama section, Japan. Geology, 41(7): 803–806.
- Abrantes Júnior, F.R. & Nogueira, A.C.R. 2013. Palaeoenvironmental reconstitution of Motuca and Sambaiba formations, Permian-Triassic of the Parnaiba Basin in southwest Maranhão State, Brazil. Geologia USP, Série Cientifica, 13(3): 65–82.
- Abu Hamad, A.M.B., Jasper, A. & Uhl, D. 2013. Charcoal remains from the Mukheiris Formation of Jordan – the first evidence of palaeowildfire from the Anisian (Middle Triassic) of Gondwana. Jordan Journal of Earth and Environmental Sciences, 5: 17–22.
- Adams, C.J., Korsch, R.J. & Griffin, W.L. 2013. Provenance comparisons between the Nambucca Block, Eastern Australia and the Torlesse Composite Terrane, New Zealand: Connections and implications from detrital zircon age patterns. Australian Journal of Earth Sciences, 60(2): 241–253.
- Adams, C.J., Mortimer, N., Campbell, H.J. & Griffin, W.L. 2013. Detrital zircon geochronology and sandstone provenance of basement Waipapa Terrane (Triassic-Cretaceous) and Cretaceous cover rocks (Northland Allochthon and Houhora Complex) in northern North Island, New Zealand. Geological Magazine, 150(1): 89–109.
- Al Gahtani, F. 2013. The influence of diagentic alterations on porosity in the Triassic Narrabeen Group, southern Sydney Basin, Australia. Geological Quarterly, 57: 613–628.
- Al-Husseini, M.I. & Koehrer, B. 2013. Chrono- and sequencestratigraphy of the Mid-Permian to Early Triassic Khuff sequences of the Arabian Plate. GeoArabia, 18(3): 103–130.
- Alaug, A.S. & El-Nakhal, H.A. 2013. Nomenclature review of the rock units in the stratigraphic lexicon of Yemen. Iranian

Journal of Earth Sciences, 5(2): 82–99.

- Aleali, M., Rahimpour-Bonab, H., Moussavi-Harami, R. & Jahani, D. 2013. Environmental and sequence stratigraphic implications of anhydrite textures: a case from the Lower Triassic of the Central Persian Gulf. Journal of Asian Earth Sciences, 75: 110–125.
- Algeo, T.J., Fraiser, M.L., Wignall, P.B. & Winguth, A.M.E. 2013. Permian-Triassic paleoceanography. Global and Planetary Change, 105: 1–6.
- Algeo, T.J., Henderson, C.M., Jinnan Tong, Qinglai Feng, Hongfu Yin & Tyson, R.V. 2013. Plankton and productivity during the Permian-Triassic boundary crisis: an analysis of organic carbon fluxes. Global and Planetary Change, 105: 52–67.
- Álvarez, J., Mpodozis, C., Blanco-Quintero, I., Garcia-Casco, A., Arriagada, C. & Morata, D. 2013. U-Pb ages and metamorphic evolution of the La Pampa gneisses: implications for the evolution of the Chilenia Terrane and Permo-Triassic tectonics of north Central Chile. Journal of South American Earth Sciences, 47: 100–115.
- Ambrose, K. 2013. Breedon Hill. Mercian Geologist 18(2): 139–143.
- Anderson, K. & Woods, A.D. 2013. Taphonomy of Early Triassic fish fossils of the Vega-Phroso Siltstone Member of the Sulphur Mountain Formation near Wapiti Lake, British Columbia, Canada. Journal of Palaeogeography, 2(4): 321–343.
- Angiolini, L., Zanchi, A., Zanchetta, S., Nicora, A. & Vezzoli, G. 2013. The Cimmerian geopuzzle: new data from South Pamir. Terra Nova, 25(5): 352–360.
- Apaldetti, C., Pol., D. & Yates, A. 2013. The postcranial anatomy of *Coloradisaurus brevis* (Dinosauria: Sauropodomorpha) from the Late Triassic of Argentina and its phylogenetic implications. Palaeontology, 56(2): 277–301.
- Aretz, M. & Hesselbo, S. 2013. Global Stratotype Sections and Points (GSSP) and Stratotypes. Ciências da Terra, 18: 35–40.
- Aristov, D.S. 2013. New Grylloblattida (Insecta) from the Upper Permian and Lower Triassic of European Russia and Kazakhstan. Paleontological Journal, 47(1): 66–76.
- Armitage, P.J., Worden, R.H., Faulkner, D.R., Aplin, A.C., Butcher, A.R. & Espie, A.A. 2013. Mercia Mudstone Formation caprock to carbon capture and storage sites:

petrology and petrophysical characteristics. Journal of the Geological Society, London, 170(1): 119–132.

- Arp, G. 2013. Geologie von Göttingen und Umgebung (Exkursion D am 4. April 2013) [Geology of Göttingen and surrounding area]. Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereins, 95: 71–90.
- Arratia, G. 2013. Morphology, taxonomy and phylogeny of Triassic pholidophorid fishes (Actinopterygii, Teleostei). Journal of Vertebrate Paleontology, 33, Supplement 1: 1–138.
- Ash, S.R. & Hasiotis, S.T. 2013. New occurrences of the controversial Late Triassic plant fossil *Sanmiguelia* Brown and associated ichnofossils in the Chinle Formation of Arizona and Utah, USA. Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen, 268(1): 65–82.
- Asvesta, A. & Dimitriadis, S. 2013. Magma-sediment interaction during the emplacement of syn-sedimentary silicic and mafic intrusions and lavas into and onto Triassic strata (Circum-Rhodope Belt, northern Greece). Geologica Carpathica, 64(3): 191–194.
- Axsmith, B.J., Fraser, N.C. & Corso, T. 2013. A Triassic seed with an angiosperm-like wind dispersal mechanism. Palaeontology, 56(5): 1173–1177.
- Bacon, K.L., Belcher, C.M., Haworth, M. & McElwain, J.C. 2013. Increased atmospheric SO₂ detected from changes in leaf physiognomy across the Triassic-Jurassic boundary interval of East Greenland. PLoS ONE, 8(4): e60614. doi:10.1371/ journal.pone.0060614.
- Baczko, M.B. von & Ezcurra, M.D. 2013. Ornithosuchidae: a group of Triassic archosaurs with a unique ankle joint. Geological Society, London, Special Publications, 379: 187–202.
- Bai-Qiu Wang, Mei-Fu Zhou, Wei Terry Chen, Jian-Feng Gao & Dan-Ping Yan. 2013. Petrogenesis and tectonic implications of the Triassic volcanic rocks in the northern Yidun Terrane, Eastern Tibet. Lithos, 175–176: 285–301.
- Bai-Qiu Wang, Wei Wang & Mei-Fu Zhou. 2013. Provenance and tectonic setting of the Triassic Yidun Group, the Yidun Terrane, Tibet. Geoscience Frontiers, 4(6): 765–777.
- Bai-Qiu Wang, Wei Wang, Wei Terry Chen, Jian-Feng Gao, Xin-Fu Zhao, Dan-Ping Yan & Mei-Fu Zhou. 2013. Constraints of detrital zircon U-Pb ages and Hf isotopes on the provenance of the Triassic Yidun Group and tectonic evolution of the Yidun Terrane, eastern Tibet. Sedimentary Geology, 289: 74–98.
- Bai Yu-bin, Luo Jing-lan, Liu Xin-ju, Jin Wen-qi & Wang Xiaojun. 2013. Geochemical characteristics of crude oil and oil source correlation in Yanchang Formation (Upper Triassic) in Wubao area, Ordos Basin. Acta Sedimentologica Sinica, 31(2): 374–383.
- Bai Yun-shan, Duan Qi-fa, Niu Zhi-jun, Tu Bing & Zeng Bo-fu. 2013. Geological characteristics and tectonic significance of volcanic rocks from Kenan Formation of the Upper Triassic in Zhahe area, southern Qinghai Province. Journal of Geology and Mineral Resources of South China, 2013(2): 100–104.
- Bandel, K. & Abu Hamad, A.M.B. 2013. Permian and Triassic strata of Jordan. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in

Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 31–41.

- Banham, S.G. & Mountney, N.P. 2013. Controls on fluvial sedimentary architecture and sediment-fill state in salt-walled mini-basins: Triassic Moenkopi Formation, Salt Anticline region, SE Utah, USA. Basin Research, 25(6): 709–737.
- Banham, S.G. & Mountney, N.P. 2013. Evolution of fluvial systems in salt-walled mini-basins: a review and new insights. Sedimentary Geology, 296: 142–166.
- Banks, D., Withers, J.G., Cashmore, G. & Dimelow, C. 2013. An overview of the results of 61 *in situ* thermal response tests in the UK. Quarterly Journal of Engineering Geology and Hydrogeology, 46(3): 281–291.
- Barrionuevo, M.F., Arnosio, J.M. & Llambías, E.J. 2013. Nuevos datos geocronológicos en subsuelo y afloramientos del Grupo Choiyoi en el oeste de La Pampa: Implicancias estratigráficas. Revista de la Asociación Geológica Argentina, 70(1): 31–39.
- Barth, G., Kozur, H.W., Franz, M. & Weems, R.E. 2013. *Paraconularia bachmanni* n. sp. from the Germanic Upper Triassic, the first fresh-water occurrence of a conularid. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 42–47.
- Baud, A. & Richoz, S. 2013. Permian-Triassic transition and the Saiq/Mahil boundary in the Oman Mountains: proposed correction for lithostratigraphic nomenclature. GeoArabia, 18(3): 87–98.
- Bazkurt, E., Winchester, J.A. & Satir, M. 2013. The Çele mafic complex: evidence for Triassic collision between the Sakarya and İstanbul Zones, NW Turkey. Tectonophysics, 595–596: 198–214.
- Beardmore, S.R., Orr, P.J. & Furrer, H. 2013. Large scale environmental gradients as a control on the skeletal taphonomy of Ichthyopterygia. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 24.
- Bedi, Y., Vasilev, E., Dabovski, C., Ergen, A., Okuyucu, C., Doğan, A., Kağan-Tekin, U., Ivanova, D., Boncheva, I., Lokova, I., Sachanski, V., Kuşcu, I., Tuncay, E., Demiray, G. D., Soycan, H. & Göncüoğlu, M.C. 2013. New age data from the tectonostratigraphic units of the Istranca "Massif" in NW Turkey: a correlation with SE Bulgaria. Geologica Carpathica, 64(4): 255–277.
- Beltrán-Triviño, A., Winkler, W. & Von Quadt, A. 2013. Tracing Alpine sediment sources through laser ablation U–Pb dating and Hf-isotopes of detrital zircons. Sedimentology, 60(1): 197–224.
- Bennett, S.C. 2013. The phylogenetic position of the Pterosauria within the Archosauromorpha re-examined. Historical Biology, 25(5–6): 545–563.
- Benton, M.J., Qiyue Zhang, Shixue Hu, Zhong-Qiang Chen, Wen Wen, Jun Liu, Jinyuan Huang, Changyong Zhu, Tao Xie, Jinnan Tong & Choo, B. 2013. Exceptional vertebrate biotas from the Triassic of China, and the expansion of marine ecosystems after the Permo-Triassic mass extinction. Earth-Science Reviews, 125: 199–243.

- Benton, M.J., Ruta, M., Dunhill, A.M. & Sakamoto, M. 2013. The first half of tetrapod evolution, sampling proxies, and fossil record quality. Palaeogeography, Palaeoclimatology, Palaeoecology, 372: 18–41.
- Bergene, J.A., Taylor, E.L. & Taylor, T.N. 2013. *Dordrechtites arcanus* sp. nov., an anatomically preserved gymnospermous reproductive structure from the Middle Triassic of Antarctica. International Journal of Plant Sciences, 174(2): 250–265.
- Bernardi, M., Petti, F.M., Porchetti, S.d'O. & Avanzini, M. 2013. Large tridactyl footprints associated with a diverse ichnofauna from the Carnian of the southern Alps. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 48–54.
- Bialik, O.M., Korngreen, D. & Benjamini, C. 2013. Carnian (Triassic) aridization on the Levant margin: evidence from the M1 member, Mohilla Formation, Makhtesh Ramon, south Israel. Facies, 59(3): 559–581.
- Bin Wang, Guowei Zhang & Zhenyu Yang. 2013. New Mesozoic paleomagnetic results from the northeastern Sichuan basin and their implication. Tectonophysics, 608: 418–427.
- Bittencourt, J.S., Stock da Rosa, Á.A., Schultz, C.L. & Langer, M.C. 2013. Dinosaur remains from the 'Botucaraí Hill' (Caturrita Formation), Late Triassic of south Brazil, and their stratigraphic context. Historical Biology, 25(1): 81–93.
- Blackburn, T.J., Olsen, P.E., Bowring, S.A., McLean, N.M., Kent, D.V., Puffer, J., McHone, G., Rasbury, E.T. & Et-Touhami, M. 2013. Zircon U-Pb geochronology links the end-Triassic extinction with the Central Atlantic Magmatic Province. Science, 340(6135): 941–945.
- Blanchard, S., Rossignol, C., Bourquin, S., Dabard, M.-P., Hallot, E., Nalpas, T., Poujol, M., Battail, B., Jalil, N.-E., Steyer, J.-S., Vacant, R., Véran, M., Bercovici, A., Diez, J.B., Paquette, J.-L., Khenthavong, B. & Vongphamany, S. 2013. Late Triassic volcanic activity in South-East Asia: New stratigraphical, geochronological and paleontological evidence from the Luang Prabang Basin (Laos). Journal of Asian Earth Sciences, 70-71: 8–26.
- Blaźejowski, B., Duffin, C.J., Gieszcz, P., Malkowski, K., Binkowski, M., Walczak, M., Mcdonald, S.A. & Withers, P.J. 2013. *Saurichthys* (Pisces, Actinopterygii) teeth from the Lower Triassic of Spitsbergen, with comments on their stable isotope composition (δ¹³C and δ¹⁸O) and X-ray microtomography. Polish Polar Research, 34(1): 23–38.
- Bo Chen, Joachimski, M.M., Shu-zhong Shen, Lambert, L.L., Xu-long Lai, Xing-dong Wang, Jun Chen & Dong-xun Yuan. 2013. Permian ice volume and palaeoclimate history; oxygen isotope proxies revisited. Gondwana Research, 24(1): 77–89.
- Boekhout, F., Sempere, T., Spikings, R. & Schaltegger, U. 2013. Late Paleozoic to Jurassic chronostratigraphy of coastal southern Peru: temporal evolution of sedimentation along an active margin. Journal of South American Earth Sciences, 47: 179–200.
- Bogan, S., Taverne, L. & Agnolin, F. 2013. First Triassic and oldest record of a South American amiiform fish: *Caturus* sp. from the Los Menucos Group (lower Upper Triassic),

Rio Negro province, Argentina. Geologica Belgica, 16(3): 191–195.

- Bomfleur, B., Decombeix, A.-L., Escapa, I.H., Schwendemann, A.B. & Axsmith, B. 2013. Whole-plant concept and environment reconstruction of a *Telemachus* conifer (Voltziales) from the Triassic of Antarctica. International Journal of Plant Sciences, 174(3): 425–444.
- Bomfleur, B., Escapa, I.H., Serbet, R., Taylor, E.L. & Taylor, T.N. 2013. A reappraisal of *Neocalamites* and *Schizoneura* (fossil Equisetales) based on material from the Triassic of East Antarctica. Alcheringa, 37(3): 349–365.
- Bonaparte, J.F. 2013. Evolution of the Brasilodontidae (Cynodontia-Eucynodontia). Historical Biology, 25(5–6): 643–653.
- Bondarenko, L.G., Buryi, G.I., Zakharov, Y.D., Bushkareva, X.Y. & Popov, A.M. 2013. Late Smithian (Early Triassic) conodonts from Artyom, South Primorye, Russian Far East. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 53–66.
- Boonchaisuk, S., Siripunvaraporn, W. & Ogawa, Y. 2013. Evidence for middle Triassic to Miocene dual subduction zones beneath the Shan-Thai terrane, western Thailand from magnetotelluric data. Gondwana Research, 23(4): 1607–1616.
- Boro, H., Bertotti, G. & Hardebol, N.J. 2013. Distributed fracturing affecting isolated carbonate platforms, the Latemar Platform Natural Laboratory (Dolomites, North Italy). Marine and Petroleum Geology, 40: 69–84.
- Bortolotti, V., Chiari, M., Marroni, M., Pandolfi, L., Principi, G. & Saccani, E. 2013. Geodynamic evolution of ophiolites from Albania and Greece (Dinaric-Hellenic belt): one, two, or more oceanic basins? International Journal of Earth Sciences, 102(3): 783–811.
- Boulila, S., Galbrun, B. & Gardin, S. 2013. Cyclostratigraphy and recent developments in the astronomical calibration of the Geological Time Scale. Ciências da Terra, 18: 29–34.
- Bozkaya, Ö. & Yalçin, H. 2013. Geochemical monitoring of clays for diagenetic evolution of the Paleozoic-Lower Mesozoic sequence in the northern Arabian plate: Hazro and Amanos regions, Southeastern Turkey. Journal of African Earth Sciences, 86: 10–24.
- Brack, P. & Kustatscher, E. 2013. *Voltzia recubariensis* from the uppermost Angolo Limestone of the Bagolino succession (Southern Alps of Eastern Lombardy, Italy). GeoAlp, 10: 61–70.
- Bragin, N.Yu. & Bragina, L.G. 2013. Morphological characteristics of Boreal radiolarians from the Triassic and Late Cretaceous: comparative analysis. Paleontological Journal, 47(4): 351–362.
- Brandes, C., Schmidt, C., Tanner, D.C. & Winsemann, J. 2013. Paleostress pattern and salt tectonics within a developing foreland basin (north-western Subhercynian Basin, northern Germany). International Journal of Earth Sciences, 102(8): 2239–2254.
- Brandner, R. & Krystyn, L. 2013. Bericht 2012 über

Profilaufnahmen und biostratigraphische Probenbearbeitungen in der Mitteltrias der Nördlichen Kalkalpen (Karwendelgebirge) auf Blatt 2223 Innsbruck und auf Blatt 2217 Hinteriß. Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1–4): 417–420.

- Brayard, A., Bylund, K.G., Jenks, J.F., Stephen, D.A., Olivier, N., Escarguel, G., Fara, E. & Vennin, E. 2013. Smithian ammonoid faunas from Utah: implications for Early Triassic biostratigraphy, correlation and basinal paleogeography. Swiss Journal of Palaeontology, 132(2): 141–219.
- Brayard, A. & Escarguel, G. 2013. Untangling phylogenetic, geometric and ornamental imprints on Early Triassic ammonoid biogeography: a similarity-distance decay study. Lethaia, 46(1): 19–33.
- Brookfield, M.E., Algeo, T.J., Hannigan, R., Williams, J. & Bhat, G.M. 2013. Shaken *and* stirred: seismites and tsunamites at the Permian-Triassic boundary, Guryul Ravine, Kashmir, India. Palaios, 28(8): 568–582.
- Brosse, M., Brayard, A., Fara, E. & Niege, P. 2013. Ammonoid recovery after the Permian-Triassic mass extinction: a reexploration of morphological and phylogenetic diversity patterns. Journal of the Geological Society, London, 170(2): 225–236.
- Brosse, M., Goudemand, N., Frisk, A.M., Baud, A., Bagherpour, B. & Bucher, H. 2013. New conodonts from the Griesbachian microbialite in South China: implications for an improved definition of the base of the Triassic. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 26–27.
- Brusatte, S.L., Butler, R.J., Niedźwiedzki, G., Sulej, T., Bronowicz, R. & Satkünas, J. 2013. First record of Mesozoic terrestrial vertebrates from Lithuania: phytosaurs (Diapsida: Archosauriformes) of probable Late Triassic age, with a review of phytosaur biogeography. Geological Magazine, 150(1): 110–122.
- Buchs, D.M., Bagheri, S., Martin, L., Hermann, J. & Arculus, R. 2013. Paleozoic to Triassic ocean opening and closure preserved in Central Iran: Constraints from the geochemistry of meta-igneous rocks of the Anarak area. Lithos, 172–173: 267–287.
- Buono, G., Hadfield, L., Mitchell, C. & Twitchett, R.J. 2013. New information about a Triassic brachiopod association from Oman. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 64.
- Burchfiel, B.C. & Chen Zhiliang (eds). 2013. Tectonics of the southeastern Tibetan Plateau and its adjacent foreland. Geological Society of America Memoirs, 210: vii+231pp.
- Butler, R.J. 2013. '*Francosuchus*' *trauthi* is not *Paleorhinus*: implications for Late Triassic vertebrate biostratigraphy. Journal of Vertebrate Paleontology, 33(4): 858–864.
- Cadée, G.C. 2013. Can Recent traces help in recognising probable jellyfish remains from the Muschelkalk of Winterswijk? a comment. Netherlands Journal of Geosciences, 192(4): 301–302.
- Caifu Xiang, Xiongqi Pang & Danišík, M. 2013. Post-Triassic thermal history of the Tazhong Uplift Zone in the Tarim Basin, Northwest China: evidence from apatite fission-track

thermochronology. Geoscience Frontiers, 4(6): 743–754.

- Callegaro, S., Marzoli, A., Bertrand, H., Chiaradia, M., Reisberg, L., Meyzen, C., Bellieni, G., Weems, R.E. & Merle, R. 2013. Upper and lower crust recycling in the source of the CAMP basaltic dykes from southeastern North America. Earth and Planetary Science Letters, 376: 186–199.
- Canfield, D.E. 2013, Sulfur isotopes in coal constrain the evolution of the Phanerozoic sulphur cycle. Proceedings of the National Academy of Sciences, 110(21): 8443–8446.
- Čar, J. 2013. Ladinian skonca beds of the Idrija ore deposit. Geologija, 56(2): 151–174.
- Carroll, A.R., Dumitru, T.A., Graham, S.A. & Hendrix, M.S. 2013. An 800 million-year detrital zircon record of continental amalgamation: Tarim basin, NW China. International Geology Review, 55(7): 818–829.
- Carter, E.S. & Orchard, M.J. 2013. Intercalibration of conodont and radiolarian faunas from the Carnian-Norian boundary interval in Haida Gwaii, British Columbia, Canada. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 67–92.
- Cather, S.M., Zeigler, K.E., Mack, G.H. & Kelley, S.A. 2013. Toward standardization of Phanerozoic stratigraphic nomenclature in New Mexico. Rocky Mountain Geology, 48(2): 101–124.
- Catlos, E.J., Huber, K. & Shin, T.A. 2013. Geochemistry and geochronology of meta-igneous rocks from the Tokat Massif, north-central Turkey: implications for Tethyan reconstructions. International Journal of Earth Sciences, 102(8): 2175–2198.
- Cavin, L., Furrer, H. & Obrist, C. 2013. New coelacanth material from the Middle Triassic of eastern Switzerland, and comments on the taxic diversity of actinistans. Swiss Journal of Geology, 106(2): 161–177.
- Celarc, B., Goričan, S. & Kolar-Jurkovšek, T. 2013. Middle Triassic carbonate-platform break-up and formation of smallscale half-grabens (Julian and Kamnik-Savinja Alps, Slovenia). Facies, 59(3): 583–610.
- Cerda, I.A., Desojo, J.B., Scheyer, T.M. & Schultz, C.L. 2013. Osteoderm microstructure of "rauisuchian" archosaurs from South America. Geobios, 46(4): 273–283.
- Chaloner, W.G. 2013. Three palynological puzzles. International Journal of Plant Sciences, 174(3): 602–607.
- Charvet, J. 2013. Late Paleozoic-Mesozoic tectonic evolution of SW Japan: a review reappraisal of the accretionary orogeny and revalidation of the collisional model. Journal of Asian Earth Sciences, 72: 88–101.
- Chatalov, A. 2013. A Triassic homoclinal ramp from the Western Tethyan realm, Western Balkanides, Bulgaria: integrated insight with special emphasis on the Anisian outer to inner ramp facies transition. Palaeogeography, Palaeoclimatology, Palaeoecology, 386: 34–58.
- Chatterjee, S., Tewari, R. & Agnihotri, D. 2013. A *Dicroidium* flora from the Triassic of Allan Hills, South Victoria Land, Transantarctic Mountains, Antarctica. Alcheringa, 37(2): 209–221.

- Chatzaras, V., Dörr, W., Finger, F., Xypolias, P. & Zulauf, G. 2013. U-Pb single zircon ages and geochemistry of metagranitoid rocks in the Cycladic Blueschists (Evia Island): implications for the Triassic tectonic setting of Greece. Tectonophysics, 595-596: 125–139.
- Chemberski, H.I. & Botoucharov, N.D. 2013. Triassic lithostratigraphic correlation in the Moesian Platform (Bulgaria–Romania). Stratigraphy and Geological Correlation, 21(6): 609–627.
- Chen Guo-chao, Pei Xian-zhi, Li Rui-bao, Li Zuo-chen, Pei Lei, Liu Zhan-qing, Chen You-xin & Liu Cheng-jun, 2013. Late Triassic EKOB magma mixing: a granodiorite pluton and (not translated) case study. Geology in China, 2013/4: 1044–1065.
- Chen Xiaohong, Sander, M.P., Cheng Long & Wang Xiaofeng. 2013. A new Triassic primitive ichthyosaur from Yunnan, South China. Acta Geologica Sinica, 87(3): 672–677.
- Chen Xiao-hong, Wei Kai, Zhang Bao-ming & Cheng Long. 2013. Guanling group Yongning Town Lower Triassic carbon isotope stratigraphy and marine chemical oxidation events. Geology in China, 2013/5: 1466–1473.
- Cheng Ji, Da-Yong Jiang, Motani, R., Wei-Cheng Hao, Zuo-Yu Sun & Tao Cai. 2013. A new juvenile specimen of *Guanlingsaurus* (Ichthyosauria, Shastasauridae) from the Upper Triassic of southwestern China. Journal of Vertebrate Paleontology, 33(2): 340–348.
- Chenshou Chun, Liu Jun, Liu Yi Sha, Liu Dong Xi, Pujun Wei, Liu Fei & Han Jian. 2013. Reservoir characteristics of 2nd member of Jialingjiang Formation in Fuchengzhai structure of east Sichuan. Advances in Geosciences, 3(4): 214–220.
- Chiari, M., Baumgartner, P.O., Bernoulli, D., Bortolotti, V., Marcucci, M. & Photiades, A. 2013. Late Triassic, Early and Middle Jurassic Radiolaria from ferromanganese-chert 'nodules' (Angelokastron, Argolis, Greece): evidence for prolonged radiolarite sedimentation in the Maliac-Vardar Ocean. Facies, 59(2): 391–424.
- Choulet, F., Yan Chen, Cogné, J.-P., Rabillard, A., Bo Wang, Wei Lin, Faure, M. & Cluzel, D. 2013. First Triassic palaeomagnetic constraints from Junggar (NW China) and their implications for the Mesozoic tectonics in Central Asia. Journal of Asian Earth Sciences, 78: 371–394.
- Chrząstek, A. 2013. Trace fossils from the Lower Muschelkalk of Raciborowice Górne (North Sudetic Synclinorium, SW Poland) and their palaeoenvironmental interpretation. Acta Geologica Polonica, 63(3): 315–353.
- Chu Mei-juan, Li Shi-xiang, Liu Xian-yang, Deng Xiu-qin & Guo Zheng-quan. 2013. Accumultion mechanisms and modes of Yanchang Formation Chang 8 interval hydrocarbons in Ordos Basin. Acta Sedimentologica Sinica, 31(4): 683–692.
- Chunbo Yan, Wang L., Haishui Jing, Wignall, P.B., Yadong Sun, Yanlong Chen & Xulong Lai. 2013. Uppermost Permian to Lower Triassic conodonts at Bianyang section, Guihzou Province, South China. Palaios, 28(8): 509–522.
- Clapham, M.E., Fraiser, M.L., Marenco, P.J. & Shu-zhong Shen. 2013. Taxonomic composition and environmental distribution of post-extinction rhynchonelliform brachiopod faunas: Constraints on short-term survival and the role of anoxia in the end-Permian mass extinction. Palaeogeography,

Palaeoclimatology, Palaeoecology, 374: 284-292.

- Clarke, J. 2013. Feathers before flight. Science, 340(6133): 690-692.
- Clarkson, M.O., Richoz, S., Wood, R.A., Maurer, F., Krystyn, L., McGurty, D.J. & Astratti, D. 2013. A new high-resolution δ^{13} C record for the Early Triassic: insights from the Arabian Platform. Gondwana Research, 24(1): 233–242.
- Clémence, M.-E. & Hart, M.B. 2013. Proliferation of Oberhauserellidae during the recovery following the Late Triassic extinction: paleoecological implications. Journal of Paleontology, 87(6): 1004–1015.
- Cobbold, P.R., Zanella, A., Rodrigues, N. & Løseth, H. 2013. Bedding-parallel fibrous veins (beef and cone-incone): worldwide occurrence and possible significance in terms of fluid overpressure, hydrocarbon generation and mineralization. Marine and Petroleum Geology, 43: 1–20.
- Coram, R.A. & Radley, J.D. 2013. A chirothere footprint from the Otter Sandstone Formation (Middle Triassic, late Anisian) of Devon, UK. Proceedings of the Geologists' Association, 124(3): 520–524.
- Corfe, I., Smith, A., Hakkinen, T., Lammi, L., Nieminen, P., Kallonen, A., Hamalainen, K., Gill, P. & Jernvall, J. 2013. The developmental basis of 200 million year old mammal teeth. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 31.
- Crane, P. 2013. Ginkgo The Tree that Time Forgot. Newhaven: Yale University Press: 404 pp.
- Császár, G., Szinger, B. & Piros, O. 2013. From continental platform towards rifting of the Tisza Unit in the Late Triassic to Early Cretaceous. Geologica Carpathica, 64(4): 279–290.
- Cuthbertson, R.S., Russell, A.P. & Anderson, J.S. 2013. Reinterpretation of the cranial morphology of *Utatsusaurus hataii* (Ichthyopterygia) (Osawa Formation, Lower Triassic, Miyagi, Japan) and its systematic implications. Journal of Vertebrate Paleontology, 33(4): 817–830.
- Cuthbertson, R.S., Russell, A.P. & Anderson, J.S. 2013. Cranial morphology and relationships of a new grippidian (Ichthyopterygia) from the Vega-Phroso Siltstone Member (Lower Triassic) of British Columbia, Canada. Journal of Vertebrate Paleontology, 33(4): 831–847.
- Dalla Vecchia, F.M. 2013. Triassic pterosaurs. Geological Society, London, Special Publications, 379: 119–155.
- Dalla Vecchia, F.M. & Selden, P.A. 2013. A Triassic spider from Italy. Acta Palaeontologica Polonica, 58(2): 325–330.
- Dan Yong, Lin Liangbiao, Zhong Yijiang & Fan Yu. 2013. The conglomerate composition of the Fourth Member of Xujiahe Formation, Upper Triassic, in the front of Micang-Daba mountains, Sichuan, China: implications for provenance analysis. Geological Review, 59(1): 15–23.
- De Franca, M.A.G., Langer, M.C. & Ferigolo, J. 2013. The skull anatomy of *Decuriasuchus quartacolonia* (Pseudosuchia: Suchia: Loricata) from the Middle Triassic of Brazil. Geological Society, London, Special Publications, 379: 469–501.
- De Jersey, N.J. & McKeller, J.L. 2013. The palynology of the Triassic-Jurassic transition in southeastern Queensland, Australia, and correlation with New Zealand. Palynology, 37(1): 77–114.

- De-Bin Yang, Wen-Liang Xu, Yi-Gang Xu, Fu-Ping Pei & Feng Wang. 2013. Provenance of sediments from Mesozoic basins in western Shandong: implications for the evolution of the eastern North China Block. Journal of Asian Earth Sciences, 76: 12–29.
- De-Feng He, Wei-Guang Zhu, Hong Zhong, Tao Ren, Zhong-Jie Bai & Hong-Peng Fan. 2013. Zircon U-Pb geochronology and elemental and Sr-Nd-Hf isotopic geochemistry of the Daocheng granitic pluton from the Yidun arc, SW China. Journal of Asian Earth Sciences, 67-68: 1–17.
- Decarlis, A., Dallagiovanna, G., Lualdi, A., Maino, M. & Seno, S. 2013. Stratigraphical evolution of the Ligurian Alps between Variscan heritages and the Alpine Tethyan opening: a review. Earth-Science Reviews, 125: 43–68.
- Dehcheshmehi, S.M. & Adabi, M.H. 2013. Depositional facies and geochemistry of the Kangan Formation in the South Fars Field, Persian Gulf (Iran). Carbonates and Evaporites, 28(3): 297–307.
- Deng Xiu-qin, Luo An-xiang, Zhang Zhong-yi & Liu Xin. 2013. Geochronological comparison on Indosinian tectonic events between Qinling Orogeny and Ordos Basin. Acta Sedimentologica Sinica, 31(6): 939–953.
- Desojo, J.B., Heckert, A.B., Martz, J.W., Parker, W.G., Schoch, R.R., Small, B.J. & Sulej, T. 2013. Aetosauria: A clade of armoured pseudosuchians from the Upper Triassic continental beds. Geological Society, London, Special Publications, 379: 203–239.
- Di-Cheng Zhu, Zhi-Dan Zhao, Yaoling Niu, Yildirim Dilek, Zeng-Qian Hou & Xuan-Xue Mo. 2013. The origin and pre-Cenozoic evolution of the Tibetan Plateau. Gondwana Research, 23(4): 1429–1454.
- Dias-da-Silva, S. & Dias, E.V. 2013. A comprehensive survey of Triassic stereospondyls from southern Brazil with comments on their overall significance. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 93–102.
- Diedrich, C.G. 2013. Review of the Middle Triassic "sea cow" *Placodus gigas* (Reptilia) in Pangea's shallow marine macroalgae meadows of Europe. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 103–131.
- Diedrich, C.G. 2013. Shallow marine sauropterygian reptile biodiversity and change in the Bad Sulza Formation (Illyrian, Middle Triassic) of central Germany, and a contribution to the evolution of *Nothosaurus* in the Germanic Basin. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 132–158.
- Diedrich, C.G. 2013. The marine pachypleurosaur *Serpianosaurus germanicus* nov. spec. skeleton and isolated bone remains from the Pelsonian (Middle Triassic) of the European Germanic Basin carbonate intertidals and its paleobiology and taphonomy. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy

and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 159–168.

- Diedrich, C.G. 2013. The oldest "subaquatic flying" reptile in the world – *Pistosaurus longaevus* Mayer, 1839 (Sauropterygia) from the Middle Triassic of Europe. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 169–215.
- Diedrich, C.G. & Gradinaru, E. 2013. Distribution of basal Middle Triassic fossil reptile placodonts in the Germanic Basin and northern Tethys. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 216–227.
- Dobretsov, N.L., Polyansky, O.P., Reverdatto, V.V. & Babichev, A.V. 2013. Dynamics of the Arctic and adjacent petroleum basins: a record of plume and rifting activity. Russian Geology and Geophysics, 54(8): 888–902.
- Dobretsov, N.L., Vernikovsky, V.A., Karyakin, Yu.V., Korago, E.A. & Simonov, V.A. 2013. Mesozoic-Cenozoic volcanism and geodynamic events in the Central and Eastern Arctic. Russian Geology and Geophysics, 54(8): 874–887.
- Dong Li-yang, Li Jun, Ning Kuo-bu, Ren Fei, Wang Jian, Wang Peng, Zhou Ye-xin & Zu Guang-fu. 2013. The alluvial fan-delta depositional systems in the Lower Triassic Qingtianbao Formation, Junmachang, Eryuan, western Yunnan. Sedimentary Geology and Tethyan Geology, 33(4): 6–15.
- Dong Li-yang, Wang Jian, Wang Peng, Yin Fu-guang & Zhou Ye-xin. 2013. The storm deposits from the Middle Triassic Beiya Formation in the Beiya region, western Yunnan and their palaeogeographic implications. Sedimentary Geology and Tethyan Geology, 33(4): 16–21.
- Dongdong Liu, Jolivet, M., Wei Yang, Zuja Zhang, Feng Cheng, Bei Zhu & Zhaojie Guo. 2013. Latest Paleozoic-Early Mesozoic basin-range interactions in South Tian Shan (northwest China) and their tectonic significance: Constraints from detrital zircon. Tectonophysics, 599: 197–213.
- Donskaya, T.V., Gladkochub, D.P., Mazukabzov, A.M. & Ivanov, A.V. 2013. Late Paleozoic – Mesozoic subduction-related magmatism at the southern margin of the Siberian continent and the 150 million-year history of the Mongol-Okhotsk Ocean. Journal of Asian Earth Sciences, 62: 79–97.
- Du Yuansheng, Huang Hu, Yang Jianghai, Wang Hongwei, Tao Ping, Huang Zhiqiang, Hu Lisha & Xie Chunxia. 2013. The basin translation from Late Paleozoic to Triassic of the Youjiang Basin and its tectonic signification. Geological Review, 59(1): 1–11.
- Duan Zhi-ming, Li Guang-ming, Zhang Hui & Duan Yau-yao. 2013. Tibet Bangong – Nujiang suture zone north edge of dragons ore district Late Triassic–Jurassic accretionary complex structure and its constraints on the geological background. Geological Bulletin of China, 32(5): 742–750.
- Duffy, O.B., Gawthorpe, R.L., Docherty, M. & Brocklehurst, S.H. 2013. Mobile evaporite controls on structural style and evolution of rift basins: Danish Central Graben, North Sea.

Basin Research, 25(3): 310–330.

- Dumitrica, P. 2013. Siamese twins and twin-like skeletons in Mesozoic Polycystine Radiolaria. Revue de Micropaléontologie, 56(2): 51–61.
- Dumitrica, P., Tekin, U.K. & Bedi, Y. 2013. Taxonomic study of the tetrahedral, pentagonal and hexagonal spongy spumellarian Radiolaria from the Middle Carnian (Late Triassic) of the Köseyahya nappe (Elbistan, SE Turkey) and other Triassic localities. Paläontologische Zeitschrift, 87(3): 311–343.
- Dumitrica, P., Tekin, U.K. & Bedi, Y. 2013. Taxonomic study of spongy spumellarian Radiolaria with three and four coplanar spines or arms from the Middle Carnian (Late Triassic) of the Köseyahya nappe (Elbistan, SE Turkey) and other Triassic localities. Paläontologische Zeitschrift, 87(3): 345–395.
- Dunhill, A.M., Benton, M.J., Newell, A.J. & Twitchett, R.J. 2013. Completeness of the fossil record and the validity of sampling proxies: a case study from the Triassic of England and Wales. Journal of the Geological Society, London, 170(2): 291–300.
- Durand, M. & Bourquin, S. 2013. Criteria for the identification of ventifacts in the geological record: a review and new insights. Comptes Rendus Geoscience, 345(3): 111-125.
- Dustira, A.M., Wignall, P.B., Joachimski, M., Blomeier, D., Hartkopf-Fröder, C. & Bond, D.P.G. 2013. Gradual onset of anoxia across the Permian-Triassic boundary in Svalbard, Norway. Palaeogeography, Palaeoclimatology, Palaeoecology, 374: 303–313.
- Egglseder, M. & Fügenschuh, B. 2013. Pre-Alpine fold interference patterns in the north-eastern Oetztal–Stubai– Complex (Tyrol, Austria). Austrian Journal of Earth Sciences, 106(2): 63–74.
- Elsner, M., Schöner, R., Gerdes, X. & Gaupp, R. 2013. Reconstruction of the early Mesozoic plate margin of Gondwana by U-Pb ages of detrital zircons from northern Victoria Land, Antarctica. Geological Society, London, Special Publications, 383: 211–232.
- Esrafili-Dizaji, B., Harchegani, F.K., Rahimpour-Bonab, H. & Kamali, M. 2013. Controls on reservoir quality in the Early Triassic Kangan Formation, Iran. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 219–245.
- Esrafili-Dizaji, B. & Rahimpour-Bonab, H. 2013. A review of Permo-Triassic reservoir rocks in the Zagros area, SW Iran: influence of the Qatar – Fars Arch. Journal of Petroleum Geology, 36(3): 257–279.
- Ezcurra, M.D., Butler, R.J. & Gower, D.J. 2013. 'Proterosuchia': the origin and early history of Archosauriformes. Geological Society, London, Special Publications, 379: 9–33.
- Fang Qiang, Zhang Haijun, Wang Xunlian & Wang Jingya. 2013. The discovery of Early Triassic conodont in the Huashixia area of Maduo County, Qinghai Province, NW China and its significance. Acta Micropalaeontologica Sinica, 30(4): 423–425.
- Faqira, M.I., Bakheit, A.F., Tang, D.Z., Tan, W. & Ahmed, A. 2013. A review of the Permo-Triassic gas play in the Arabian Gulf region. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic

Sequences of the Arabian Plate. EAGE Publications bv.: 163–198.

- Fei Li, Jiaxin Yan, Algeo, T. & Xia Wu. 2013. Paleoceanographic conditions following the end-Permian mass extinction recorded by giant ooids (Moyang, South China). Global and Planetary Change, 105: 102-120.
- Feldman, H.R. 2013. *Menathyris wilsoni* (Brachiopoda), a new genus and species from the Middle Triassic (Ladinian) of southern Israel. Annales Societatis Geologorum Poloniae, 83(2): 81–85.
- Fernandez, V., Abdala, F., Carlson, K.J., Cook, D.C., Rubidge, D.S., Yates, A. & Tafforeau, P. 2013. Synchrotron reveals Early Triassic odd couple: injured amphibian and aestivating therapsid share burrow. PLoS ONE, 8(6): e64978. doi:10.1371/journal.pone.0064978.
- Fijałkowska-Mader, A. 2013. Impact of the environmental stress on the Late Permian palynoflora from Poland. Biuletyn Państwowego Instytutu Geologicznego, 452: 23–32.
- Fijałkowska-Mader, A. 2013. Palynostratigraphy, palaeoecology and palaeoclimate of the Late Permian and Triassic of the Nida Basin. Biuletyn Państwowego Instytutu Geologicznego, 454: 15–70.
- Fio, K., Sremac, J., Vlahovic, I., Velíc, I. & Spangenberg, J.E. 2013. Permian deposits and the Permian-Triassic boundary in Croatia: palaeoclimatic implications based on palaeontological and geochemical data. Geological Society, London, Special Publications, 376: 539–548.
- Forchielli, A. & Pervesler, P. 2013. Phosphatic cuticle in thyalocephalans: A taphonomic case study of *Austriocaris* (Arthropoda, Thyalocephala) from the Fossil-Lagerstätte Polzberg (Reingraben shales, Carnian, Upper Triassic, Lower Austria). Austrian Journal of Earth Sciences, 106(1): 46–61.
- Forel, M.-B. 2013. The Permian-Triassic mass extinction: ostracods (Crustacea) and microbialites. Comptes Rendus Geoscience, 345(4): 203-211.
- Forel, M.-B., Crasquin, S., Hips, K., Kershaw, S., Collin, P.-Y. & Haas, J. 2013. Biodiversity evolution through the Permian-Triassic boundary event: ostracods from the Bükk Mountains, Hungary. Acta Palaeontologica Polonica, 58(1): 195–219.
- Forel, M.-B., Crasquin, S., Kershaw, S. & Collin, P.-Y. 2013. In the aftermath of the end-Permian extinction: the microbialite refuge? Terra Nova, 25(2): 137–143.
- Forke, H.C., Köhrer, B., Bendias, D., Walz, L. & Aigner, T. 2013. Biostratigraphy and biofacies of Khuff time-equivalent strata in the Al-Jabal Al-Akhdar area (Hajar Mountains), northern Oman. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 77–125.
- Foster, W.J. & Twitchett, R. 2013. Functional diversity across the Permian/Triassic boundary. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 70.
- Foth, C. & Rauhut, O.W.M. 2013. The Good, the Bad and the Ugly: the influence of skull reconstructions and intraspecific variability in studies of cranial morphometrics in theropods and basal saurischians. PLoS ONE, 8(8): e72007. doi:10.1371/journal.pone.0072007.
- Frankowiak, K., Mazur, M., Gothmann, A.M. & Stolarski, J.

2013. Diagenetic alteration of Triassic coral from the aragonite Konservat-Lagerstätte in Alakir Çay, Turkey: implications for geochemical measurements. Palaios, 28(6): 333–342.

- Franz, M., Henninger, M. & Barnasch, J. 2013. The strong diachronous Muschelkalk/Keuper facies shift in the Central European Basin: implications from the type-section of the Erfurt Formation (Lower Keuper, Triassic) and basin-wide correlations. International Journal of Earth Sciences, 102(3): 761–780.
- Fraser, N.C. & Furrer, H. 2013. A new species of *Macrocnemus* from the Middle Triassic of the eastern Swiss Alps. Swiss Journal of Geology, 106(2): 199–206.
- Fraser, N.C., Rieppel, O. & Li Chun. 2013. A long-snouted protorosaur from the Middle Triassic of southern China. Journal of Vertebrate Paleontology, 33(5): 1120–1126.
- Fröbisch, J. 2013. Vertebrate diversity across the end-Permian mass extinction – separating biological and geological signals. Palaeogeography, Palaeoclimatology, Palaeoecology, 372: 50–61.
- Fu Jin-hua, Deng Xiu-qin, Chu Mei-juan, Zhang Hai-feng & Li Shi-xiang. 2013. Features of deepwater lithofacies, Yanchang Formation in Ordos Basin and its petroleum significance. Acta Sedimentologica Sinica, 31(5): 928–938.
- Fu, X., Wang, J., Tan, F., Feng, X. & Zeng, S. 2013. Minerals and potentially hazardous trace elements in the late Triassic coals from the Qiangtang Basin, China. International Journal of Coal Geology, 116-117: 93–105.
- Fulai Liu, Fang Wang, Pinghua Liu & Chaohui Liu. 2013. Multiple metamorphic events revealed by zircons from the Diancang Shan – Ailao Shan metamorphic complex, southeastern Tibetan Plateau. Gondwana Research, 24(1): 429–450.
- Gabtni, H., Jallouli, C., Mikus, K.L. & Turki, M.M. 2013. Geodynamics of the southern Tethyan margin in Tunisia and Maghrebian domain: new constraints from integrated geophysical study. Arabian Journal of Geosciences, 6(1): 271–286.
- Gaetani, M., Nicora, A., Henderson, C., Cirilli, S., Gale, L., Rettori, R., Vuolo, I. & Atudorei, V. 2013. Refinements in the Upper Permian to Lower Jurassic stratigraphy of Karakorum, Pakistan. Facies, 59(4): 915–948.
- Galán-Abellán, B., Alonso-Azcárate, J., Newton, R.J., Bottrell, S.H., Barrenechea, J.F., Benito, M.I., De la Horra, R., López-Gómez, J. & Luque, F.J. 2013. Sources of Sr and S in aluminium-phosphate-sulphate minerals in Early-Middle Triassic sandstones (Iberian Ranges, Spain) and paleoenvironmental implications for the west Tethys. Journal of Sedimentary Research, 83(5): 406–426.
- Galán-Abellán, B., Barrenechea, J.F., Benito, M.I., De la Horra, R., Luque, F.J., Alonso-Azcárate, J., Arche, A., López-Gómez, J. & Lago, M. 2013. Palaeoenvironmental implications of aluminium phosphate-sulphate minerals in Early-Middle Triassic continental sediments, SE Iberian Range (Spain). Sedimentary Geology, 289: 169–181.
- Galán-Abellán, B., López-Gómez, J., Barrenechea, J.F., Marzo, M., De la Horra, R. & Arche, A. 2013. The beginning of the Buntsandstein cycle (Early-Middle Triassic) in the Catalan

Ranges, NE Spain: sedimentary and palaeogeographic implications. Sedimentary Geology, 296: 86–102.

- Gale, L., Kastelic, A. & Rožič, B. 2013. Taphonomic features of Late Triassic foraminifera from Mount Begunjščica, Karavanke Mountains, Slovenia. Palaios, 28(11): 771–792.
- Gale, L., Rettori, R., Martini, R. & Rožič, B. 2013. *Decapoalina* n. gen. (Miliolata, Milioporidae; Late Triassic), a new foraminiferal genus for "*Sigmoilina*" schaeferae" Zaninetti, Altiner, Dager & Ducret, 1982. Bollettino della Società Paleontologica Italiana, 52(2): 81–93.
- Gallois, R.W. 2013. The relationship of stratigraphically controlled lithological variations in the Mercia Mudstone Group (Triassic) in South-West England to weathering processes and landslide mechanisms. Geoscience in South-West England, 13(2): 172–182.
- Gallois, R.W. 2013. Possible tetrapod burrows in the Mid Triassic Otter Sandstone Formation at Sidmouth, Devon, UK. Geoscience in South-West England, 13(2): 208–215.
- Garcia-Lasanta, C., Oliva-Orcia, B., Román-Berdiel, T., Gil-Peña, I., Sánchez-Moya, Y. & Sopeña, A. 2013. Tectonically controlled magnetic fabrics in the Triassic Iberian Basin. Geogaceta, 53; 53–56.
- Gardener, J.D. & DeMar, D.G. Jr. 2013. Mesozoic and Palaeocene lissamphibian assemblages of North America: a comprehensive review. Palaeodiversity and Palaeoenvironments, 93(4): 459–515.
- Gastaldo, R.A., Pludow, B.A. & Neveling, J. 2013. Mud aggregates from the Katberg Formation, South Africa: additional evidence of Early Triassic degradational landscapes. Journal of Sedimentary Research, 83(7): 531–540.
- Gattolin, G., Breda, A. & Preto, N. 2013. Demise of Late Triassic carbonate platforms triggered the onset of a tide-dominated depositional system in the Dolomites. Sedimentary Geology, 297: 31–37.
- Gaździcki, A. & Uchman, A. 2013. Fatricum provenance of the Rhaetian deposits in the "Nad Moreną" section (Przysłop Miętusi area): a correction of the Hronicum stratigraphy in the Tatra Mts. Przegląd Geologiczny, 60(6): 333–340.
- Ge Zhong-wei, Wei Li-min, Fan Li & Zhang Zhuang. 2013. Sources and sedimentary facies of the 2nd Member of Xujiahe Formation, Upper Triassic in Dayi area. Journal of Geology and Mineral Resources of South China, 2013(1): 72–78.
- Genming Luo, Yongbiao Wang, Grice, K., Kershaw, S., Algeo, T.J., Xiaoyan Ruan, Hao Yang, Chengling Jia & Shucheng Xie. 2013. Microbial-algal community changes during the latest Permian ecological crisis: evidence from lipid biomarkers at Cili, South China. Global and Planetary Change, 105: 36–51.
- Gibson, S.Z. 2013. A new hump-backed ginglymodian fish (Neopterygii, Semionotiformes) from the Upper Triassic Chinle Formation of southeastern Utah. Journal of Vertebrate Paleontology, 33(5): 1037–1050.
- Goswami, S. & Singh, K.J. 2013. Floral diversity and geology of the Talcher Basin, Orissa, India during the Permian-Triassic interval. Geological Journal, 48(1): 39–56.
- Goudemand, N., Roghi, G., Rigo, M., Brosse, M., Meier, M., Tafforeau, P. & Urdy, S. 2013. New evidence for a lamprey-like motion of the grasping elements of ozarkodinid conodonts.

The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 36.

- Granier, B. 2013. Micropaleontology from the STRATI 2013 perspective. Ciências da Terra, 18: 89–94.
- Granier, B., Lethiers, A. & Sander, N.J. 2013. The XXIst Century Edition of the "New studies on Triassic Siphoneae verticillatae by Julius von Pia". Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1–4): 239–300.
- Granier, B. & Sander, N.J. 2013. The XXIst Century (the 100th Anniversary) edition of "New studies on Triassic Siphoneae verticillatae" by Julius von PIA. Carnets de Géologie, CG2013 Bo 1, 108pp.
- Grasby, S.E., Beauchamp, B., Embry, A. & Sanei, H. 2013. Recurrent Early Triassic ocean anoxia. Geology, 41(2): 175–178.
- Grasby, S.E., Sanei, H., Beauchamp, B. & Zhuoheng Chen. 2013. Mercury deposition through the Permo-Triassic biotic crisis. Chemical Geology, 351: 209–216.
- Gregori, D. & Benedini, L. 2013. The Cordon del Portillo Permian magmatism, Mendoza, Argentina, plutonic and volcanic sequences at the western margin of Gondwana. Journal of South American Earth Sciences, 42: 61–73.
- Gretz, M., Lathuilière, B., Martini, R. & Bartolini, A. 2013. The Hettangian corals of the Isle of Skye (Scotland): an opportunity to better understand the paleoenvironmental conditions during the aftermath of the Triassic–Jurassic boundary crisis. Palaeogeography, Palaeoclimatology, Palaeoecology, 376: 132–148.
- Griebeler, E.M. 2013. Body temperatures in dinosaurs: what can growth curves tell us? PLoS ONE, 8(10): e74317. doi:10.1371/journal.pone.0074317.
- Griebeler, E.M., Klein, N. & Sander, P.M. 2013. Aging, maturation and growth of sauropodomorph dinosaurs as deduced from growth curves using long bone histological data: an assessment of methodological constraints and solutions. PLoS ONE, 8(6): e67012. doi:10.1371/journal. pone.0067012.
- Gründel, J. & Nützel, A. 2013. Evolution and classification of Mesozoic mathildoid gastropods. Acta Palaeontologica Polonica, 58(4): 803–826.
- Guang-Hui Xu, Li-Jun Zhao, Ke-Qin Gao & Fei-Xiang Wu. 2013. A new stem-neopterygian fish from the Middle Triassic of China shows the earliest over-water gliding strategy of the vertebrates. Proceedings of the Royal Society, B.280(1750) 20122261; doi: 10.1098/rspb.2012.2261 (7 pp).
- Guangyou Zhu, Shuichang Zhang, Keyu Liu, Haijun Yang, Bin Zhang, Jin Su & Yaguang Zhang. 2013. A well-preserved 250 million-year-old oil accumulation in the Tarim Basin, western China: implications for hydrocarbon exploration in old and deep basins. Marine and Petroleum Geology, 43: 478–488.
- Guochen Dong, Xuanxue Mo, Zhidan Zhao, Dicheng Zhu, Goodman, R.C., Huilei Kong & Shuo Wang. 2013. Zircon U-Pb dating and the petrological and geochemical constraints on Lincang granite in western Yunnan, China: implications for the closure of the Paleo-Tethys ocean. Journal of Asian Earth Sciences, 62: 282–294.

Guoyi Hu, Cong Yu, Yunyan Ni, Shipeng Huang & Xingwang

Tan. 2013. Comparative study of stable carbon and hydrogen isotopes of alkane gases sourced from the Longtan and Xujiahe coal-bearing measures in the Sichuan Basin, China. International Journal of Coal Geology, 116-117: 293–301.

- Haas, J., Pelikán, P., Görög, A., Józsa, S. & Ozsvárt, P. 2013. Stratigraphy, facies and geodynamic settings of Jurassic formations in the Bükk Mountains, North Hungary: its relations with the other areas of the Neotethyan realm. Geological Magazine, 150(1): 18–49.
- Haijun Song, Wignall, P.B., Jinnan Tong & Hongfu Yin. 2013.Two pulses of extinction during the Permian–Triassic crisis.Nature Geoscience, 6(1): 52–56.
- Hancox, P.J., Angielczyk, K.D. & Rubidge, B.S. 2013. Angonisaurus and Shansiodon, dicynodonts (Therapsida, Anomodontia) from subzone C of the Cynognathus Assemblage Zone (Middle Triassic) of South Africa. Journal of Vertebrate Paleontology, 33(3): 655–676.
- Hara, H., Kon, Y., Usuki, T., Ching-Ying Lan, Kamata, Y., Hisada, K., Ueno, K., Charoentitirat, T. & Charusiri, P. 2013.
 U-Pb ages of detrital zircons within the Inthanon Zone of the Paleo-Tethyan subduction zone, northern Thailand: new constraints on accretionary age and arc activity. Journal of Asian Earth Sciences, 74: 50–61.
- Harris, D.B., Toro, J. & Prokopiev, A.V. 2013. Detrital zircon U-Pb geochronology of Mesozoic sandstones from the Lower Yana River, northern Russia. Lithosphere, 5(1): 98–108.
- Hauser, L.M. & Martill, D.M. 2013. Evidence for coelacanths in the Late Triassic (Rhaetian) of England. Proceedings of the Geologists' Association, 124(6): 982–987.
- Hautmann, M. & Hagdorn, H. 2013. Oysters and oysterlike bivalves from the Middle Triassic Muschelkalk of the Germanic Basin. Paläontologische Zeitschrift, 87(1): 19–32.
- Hautmann, M., Hofmann, R., Frisk, A.M., Brosse, M. & Bucher,
 H. 2013. Diversification and diversity partitioning in Early Triassic benthic ecosytems. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 39.
- Hautmann, M., Smith, A.B., McGowan, A.J. & Bucher, H. 2013. Bivalves from the Olenekian (Early Triassic) of south-western Utah: systematics and evolutionary significance. Journal of Systematic Palaeontology, 11(3): 263–293.
- Havlik, P., Aiglstorfer, M., Atfy, H.El. & Uhl, D. 2013. A peculiar bonebed from the Norian Stubensandstein (Löwenstein Formation, Late Triassic) of southern Germany and its palaeoenvironmental interpretation. Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen, 269(3): 321–337.
- Hawie, N., Gonni, C., Deschamps, R., Nader, F.H., Montadert,
 L., Granjean, D. & Baudin, F. 2013. Tectono-stratigraphic
 evolution of the northern Levant Basin (offshore Lebanon).
 Marine and Petroleum Geology, 48: 392–410.
- Heckert, A.B., Jenkins, H.S., Lucas, S.G. & Hunt, A.P. 2013.
 Mandibles of juvenile phytosaurs (Archosauria: Crurotarsi) from the Upper Triassic Chinle Group of Texas and New Mexico, USA. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of

Natural History and Science, Bulletin 61: 228–236.

- Heckert, A.B. & Miller-Camp, J.A. 2013. Tooth enamel microstructure of *Revueltosaurus* and *Krzyanowskisaurus* (Reptila: Archosauria) from the Upper Triassic Chinle Group, USA: implications for function, growth, and phylogeny. Palaeontologia Electronica, 16(1): 23 pp (Article 16.1.1A).
- Helbig, M., Keppie, J.D., Murphy, J.B. & Solari, L.A. 2013. Exotic rifted passive margin of a back-arc basin off western Pangea: geochemical evidence from the Early Mesozoic Ayú Complex, southern Mexico. International Geology Review, 55(7): 863–881.
- Henares, S., Viseras, C., Fernández, J. & Lizarazo, N. 2013. Point bar deposit characterization with preliminary subsurface data: a case from the Triassic of the Iberian Mesata. Geogaceta, 54: 7–10.
- Heydari, E., Arzani, N., Safaei, M. & Hassanzadeh, J. 2013. Ocean's response to a changing climate: clues from variations in carbonate mineralogy across the Permian-Triassic boundary of the Shareza section, Iran. Global and Planetary Change, 105: 79–90.
- Hiete, M., Röhling, H.-G., Heunisch, C. & Berner, U. 2013. Facies and climate changes across the Permian-Triassic boundary in the North German Basin: insights from a highresolution organic carbon isotope record. Geological Society, London, Special Publications, 376: 549–574.
- Hillebrandt, A.v., Krystyn, L., Kürschner, W.M., Bonis, N.R., Ruhl, M., Richoz, S., Schobben, M.A.N., Urlichs, M., Bown, P.R., Kment, K., McRoberts, C.A., Simms, M. & Tomásových, A. 2013. The Global Stratotype Sections and Point (GSSP) for the base of the Jurassic System at Kuhjoch (Karwendel Mountains, Northern Calcareous Alps, Tyrol, Austria). Episodes, 36(3): 162–198.
- Hminna, A., Voigt, S., Klein, H., Saber, H., Schneider, J.W. & Hmich, D. 2013. First occurrence of tetrapod footprints from the continental Triassic of the Sidi Said Maachou area (Western Meseta, Morocco). Journal of African Earth Sciences, 80: 1-7.
- Hofmann, R., Hautmann, M. & Bucher, H. 2013. A new paleoecological look at the Dinwoody Formation (Lower Triassic, Western USA): intrinsic versus extrinsic controls on ecosystem recovery after the end-Permian mass extinction. Journal of Paleontology, 87(5): 854–880.
- Hofmann, R., Hautmann, M., Schneebeli-Herman, E. & Bucher, H. 2013. Evidence for an intra-Early Triassic crisis of benthic ecosystems. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 39.
- Hofmann, R., Hautmann, M., Wasmer, M. & Bucher, H. 2013. Palaeoecology of the Spathian Virgin Formation (Utah, USA) and its implications for the Early Triassic recovery. Acta Palaeontologica Polonica 58(1): 149–173.
- Holliday, C.M. & Nesbitt, S.J. 2013. Morphology and diversity of the mandibular symphysis of archosauriforms. Geological Society, London, Special Publications, 379: 555–571.
- Holloway, W.L., Claeson, K.M. & O'keefe, F.R. 2013. A virtual phytosaur endocast and its implications for sensory system evolution in archosaurs. Journal of Vertebrate Paleontology, 33(4): 848–857.

- Hongfu Yin & Haijun Song. 2013. Mass extinction and Pangea integration during the Paleozoic-Mesozoic. SCIENCE CHINA Earth Sciences, 56(11): 1791–1803.
- Hong-Yan Li & Xiao-Long Huang. 2013. Constraints on the paleogeographic evolution of the North China Craton during the Late Triassic-Jurassic. Journal of Asian Earth Sciences, 70-71: 308–320.
- Horacek, M. & Brandner, R. 2013. Comment on: Plankton and productivity during the Permian-Triassic boundary crisis: an analysis of organic carbon fluxes (Algeo et al., 2013, Global and Planetary Change). Global and Planetary Change, 111: 327–330.
- Horn, B.L.D., Pereira, V.P. & Schultz, C.L. 2013. Calcretes of the Santa Maria Supersequence, Middle Triassic, Rio Grande do Sul: classification, genesis and paleoclimatic implications. Palaeogeography, Palaeoclimatology, Palaeoecology, 376: 39–47.
- Hornung, T. 2013. Berichte 2012 über geologische Aufnahmen im Bereich Zahmer Kaiser (Kaisergebirge) auf Blatt 3213 Kufstein. Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1-4): 423–430.
- Howell, E.R. & Blakey, R.C. 2013. Sedimentological constraints on the evolution of the Cordilleran arc: new insights from the Sonsela Member, Upper Triassic Chinle Formation, Petrified Forest National Park (Arizona, USA). Geological Society of America Bulletin, 125(7/8): 1349–1368.
- Hu Bin, Shen Dajun, Wei Wei & Liu Yi Sha. 2013. Study of diagenesis and diagenetic facies of 2nd member of Lower Triassic Jialingjiang Formation in east Sichuan. Advances in Geosciences, 3(5): 289–298.
- Hua-Hua Cao, Wen-Liang Xu, Fu-Ping Pei, Zhi-Wei Wang, Feng Wang & Zi-Jin Wang. 2013. Zircon U-Pb geochronology and petrogenesis of the Late Paleozoic-Early Mesozoic intrusive rocks in the eastern segment of the northern margin of the North China Block. Lithos, 170-171: 191–207.
- Huang Jianguo, Yang Ruidong, Yang Jin & Cui Chunlong. 2013. Geochemical characteristics and tectonic significance of Triassic granite from Taer region, the northern margin of west Kunlun. Acta Geologica Sinica, 87(2): 346–357.
- Huang Zhiying, Zhu Tong-xing, Zhang Qi-yue, Zhou Changyong & Xie Tao. 2013. Southeast Yunnan newly discovered group of Middle Triassic Guanling typical platform margin and slope slumping limestone reef. Geological Bulletin of China, 32(9): 1436–1439.
- Hughes, G.W. 2013. Late Permian to Late Jurassic "microproblematica" of Saudi Arabia: possible palaeobiological assignments and roles in the palaeoenvironmental reconstructions. GeoArabia, 18(1): 57–92.
- Hui Tian, Peng Cheng, Qin Zhou, Xianming Xiao & Wilkins, R.W.T. 2013. Abnormal elevations of $C_{34}2\alpha$ -methylhopane and $C_{34}2\alpha$ -methylbenzhopane in a Lower Triassic mudstone sample, NW Sichuan Basin. Organic Geochemistry, 63: 139–144.
- Hunt, A.P., Lucas, S.G. & Spielmann, J.A. 2013. Triassic vertebrate coprolite ichnofaunas. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New

Mexico Museum of Natural History and Science, Bulletin 61: 237–258.

- Husain, R., Khan, D.A., Sajer, A., Al-Ammar, N. & Al-Fares, A. 2013. Khuff Formation in Kuwait: an overview. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 303–326.
- Huttenlocker, A.K. & Botha-Brink, J. 2013. Body size and growth patterns in the therocephalian *Moschorhinus kitchingi* (Therapsida: Eutheriodontia) before and after the end-Permian extinction in South Africa. Paleobiology, 39(2): 253–277.
- Huyue Song, Jinnan Tong, Algeo, T.J., Horacek, M., Haiou Qiu, Haijun Song, Li Tian & Zhong-Qiang Chen. 2013. Large vertical $\delta^{13}C_{_{DIC}}$ gradients in Early Triassic seas of the South China craton: implications for oceanographic changes related to Siberian Trap volcanism. Global and Planetary Change, 105: 7–20.
- Hyeong Soo Kim & Jin-Han Ree. 2013. Permo-Triassic changes in bulk crustal shortening direction during deformation and metamorphism of the Taebaeksan Basin, South Korea using foliation intersection/inflection axes: implications for tectonic movement at the eastern margin of Eurasia during the Songrim (Indosinian) orogeny. Tectonophysics, 587: 133–145.
- Iannuzzi, R. & Shu-Zhong Shen. 2013. Event stratigraphy. Ciências da Terra, 18: 121–124.
- Ikeda, M. 2013. Cyclostratigraphy of the Triassic to Jurassic deepsea sequences in Japan: frequency modulation for stratigraphic correlation. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 259–267.
- Irmis, R.B., Nesbitt, S.J. & Sues, H.-D. 2013. Early Crocodylomorpha. Geological Society, London, Special Publications, 379: 275–302.
- Irmis, R.B., Whiteside, J.H. & Kammerer, C.F. 2013. Non-biotic controls of observed diversity in the paleontologic record: An example from the Permo-Triassic Karoo Basin of South Africa. Palaeogeography, Palaeoclimatology, Palaeoecology, 372: 62–77.
- Ivanov, A.V., Huayiu He, Leikun Yan, Ryabov, V.V., Shevko, A.Y., Palesskii, S.V. & Nikolaeva, I.V. 2013. Siberian Traps large igneous province: evidence for two flood basalt pulses around the Permo-Triassic boundary and in the Middle Triassic, and contemporaneous granitic magmatism. Earth-Science Reviews, 122: 58–76.
- Jaglarz, P. 2013. Multi-stage dolomitization in the Lower–Middle Triassic succession of the High-Tatric series (Western Tatra Mts., Poland). Przegląd Geologiczny, 60(5): 284–293.
- Janson, X., Lucia, F.J., Jennings, J.W., Bellian, J.A., AbuBshait, A.A., Al-Dukhayyil, R.K., Mueller, H.W. & Cantrell, D. 2013. Outcrop-based 3D geological and reservoir model of the uppermost Khuff Formation in central Saudi Arabia. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 269–302.
- Jaraula, C.M.B., Grice, K., Twitchett, R.J., Böttcher, M.E., LeMetayer, P., Dastidar, A.G. & Opazo, L.F. 2013. Elevated pCO₂ leading to Late Triassic extinction, persistent photic

- zone euxinia, and rising sea levels. Geology, 41(9): 955–958.
 Jenks, J., Guex, J., Hungerbühler, A., Taylor, D.G. & Bucher, H. 2013. Ammonoid biostratigraphy of the early Spathian *Columbites parisianus* zone (Early Triassic) at Bear Lake Hot Springs, Idaho. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 268–283.
- Ji Liming & Zhu Youhua. 2013. Sporo-pollen assemblages and paleoclimate of the Yanchang Formation in the Xifeng area, southwestern Ordos Basin, Gansu Province, NW China. Acta Micropalaeontologica Sinica, 30(4): 367–378.
- Jian-Bo Zhou & Wilde, S.A. 2013. The crustal accretion history and tectonic evolution of the NE China segment of the Central Asian Orogenic Belt. Gondwana Research, 23(4): 1365–1377.
- Jian-Wei Zi, Cawood, P.A., Wei-Ming Fan, Tohver, E., Yue-Jun Wang, McCuaig, T.C. & Tou-Ping Peng. 2013. Late Permian-Triassic magmatic evolution in the Jinshajiang orogenic belt, SW China and implications for orogenic processes following closure of the Paleo-Tethys. American Journal of Science, 313(2): 81–112.
- Jiang Hongxia & Wu Yasheng. 2013. Diagenesis of the microbialites in the Permian-Triassic boundary section at Laolongdong, Chongqing, South China. Journal of Palaeogeography, 2(2): 183–191.
- Jiang, L., Cai, C.F., Worden, R.H., Li, K.-K. & Xiang, L. 2013. Reflux dolomitization of the Upper Permian Changxing Formation and the Lower Triassic Feixianguan Formation, NE Sichuan Basin, China. Geofluids, 13(2): 232–245.
- Jianhua Li, Yueqiao Zhang, Shuwen Dong & Wei Shi. 2013. Structural and geochronological constraints on the Mesozoic tectonic evolution of the north Dabashan zone, south Qinling, central China. Journal of Asian Earth Sciences, 63: 99–114.
- Jianjun Cui, Yueqiao Zhang, Shuwen Dong, Bor-ming Jahn, Xianbing Xu & Licheng Ma. 2013. Zircon U-Pb geochronology of the Mesozoic metamorphic rocks and granitoids in the coastal tectonic zone of SE China: constraints on the timing of Late Mesozoic orogeny. Journal of Asian Earth Sciences, 62: 237–252.
- Jianren Mao, Haimin Ye, Kai Liu, Zilong Li, Takahashi, Y., Xilin Zhao & Weon-Seo Kee. 2013. The Indosinian collisionextension event between the South China Block and the Palaeo-Pacific plate: evidence from Indosinian alkaline granitic rocks in Dashuang, eastern Zhejiang, South China. Lithos, 172–173; 81–97.
- Jiawei Zhang, Zhilong Huang, Taiyi Luo, Zhikuan Qian & Ying Zhang. 2013. Origin of early Triassic rift-related alkaline basalts from southwest China: age, isotope, and trace-element constraints. International Geology Review, 55(9): 1162–1178.
- Jingen Dai, Chengshan Wang, Hourigan, J. & Santosh, M. 2013. Multi-stage tectono-magmatic events of the Eastern Kunlun Range, northern Tibet: insights from U-Pb geochronology and (U-Th)/He thermochronology. Tectonophysics, 599: 97–106.
- Jinyuan Huang, Feldmann, R.M., Schweitzer, C.E., Shixue Hu, Changyong Zhou, Benton, M.J., Qiyue Zhang, Wen Wen & Tao Xie. 2013. A new shrimp (Decapoda, Dendrobranchiata,

Penaeoidea) from the Middle Triassic of Yunnan, southwest China. Journal of Paleontology, 87(4): 603–611.

- Johannessen, K.C., Kohlmann, F., Ksienzyk, A.K., Dunkl, I. & Jacobs, J. 2013. Tectonic evolution of the SW Norwegian passive margin based on low-temperature thermochronology from the innermost Hardangerfjord area. Norwegian Journal of Geology, 93(3-4): 243–260.
- Jun Liu, Li-Jun Zhao, Chun Li & Tao He. 2013. Osteology of *Concavispina biseridens* (Reptilia, Thallatosauria) from the Xiaowa Formation (Carnian), Guanling, Guizhou, China. Journal of Paleontology, 87(2): 341–350.
- Jun Liu, Motani, R., Da-yong Jiang, Shi-xue Hu, Aitchison, J.C., Rieppel, O., Benton, M.J., Qi-yue Zhang & Changyong Zhou. 2013. The first specimens of the Middle Triassic *Phalarodon atavus* (Ichthyosauria: Mixosauridae) from South China, showing postcranial anatomy and peri-Tethyan distribution. Palaeontology, 56(4): 849–866.
- Jun Shen, Algeo, T.J., Qing Hu, Guozhen Xu, Lian Zhou & Qinglai Feng. 2013. Volcanism in South China during the Late Permian and its relationship to marine ecosystem and environmental changes. Global and Planetary Change, 105: 121–134.
- Jun Shen, Algeo, T.J., Qinglai Feng, Lian Zhou, Lanping Feng, Ning Zhang & Junhua Huang. 2013. Volcanically induced environmental change at the Permian-Triassic boundary (Xiakou, Hubei Province, South China): related to West Siberian coal-field methane releases? Journal of Asian Earth Sciences, 75: 95–109.
- Jun Shen, Yong Lei, Algeo, T.J., Qinglai Feng, Servais, T., Jianxin Yu & Lian Zhou. 2013. Volcanic effects on microplankton during the Permian-Triassic transition (Shangsi and Xinmin, South China). Palaios, 29(8): 552–567.
- Kai-Xing Wang, Tao Sun, Pei-Rong Chen, Hong-Fei Ling & Ting-Fu Xiang. 2013. The geochronological and geochemical constraints on the petrogenesis of the Early Mesozoic A-type granite and diabase in northwestern Fujian province. Lithos, 179: 364–381.
- Kaiho, K. & Koga, S. 2013. Impacts of a massive release of methane and hydrogen sulphide on oxygen and ozone during the late Permian mass extinction. Global and Planetary Change, 107: 91–101.
- Kammerer, C.F., Fröbisch, J. & Angielczyk, K.D. 2013. On the validity and phylogenetic position of *Eubrachiosaurus browni*, a kannemeyeriiform dicynodont (Anomodontia) from the Triassic North America. PLoS ONE, 8(5): e64203. doi:10.1371/journal.pone.0064203.
- Kaplanis, A., Koukouvelas, I., Xypolias, P. & Kokkolas, S. 2013. Kinematics and ophiolite obduction in the Gerania and Helicon Mountains, central Greece. Tectonophysics, 595-596: 215–234.
- Karádi, V., Kozur, H. & Görög, A. 2013. Stratigraphically important lower Norian conodonts from the Csővár borehole (CSV-1), Hungary – comparison with the conodont succession of the Norian GSSP candidate Pizzo Mondello (Sicily, Italy). *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History

and Science, Bulletin 61: 284-295.

- Karasev, E.V. 2013. Formal system of dispersed leaf cuticles of pteridosperms (Peltaspermaceae) from the Permian and Triassic of the Russian Platform. Paleontological Journal, 47(3): 335-349.
- Kawagoe, Y., Mori, N., Sano, S., Orihasi, Y., Tamamoto, K., Ishizaki, Y., Kouchi, Y. & Otoh, S. 2013. Evidence for Late Permian-Triassic volcanism in the Hida Gaien Belt, southwest Japan: new U-Pb ages from the Motodo, Ashidani, and Otani formations. Memoir of the Fukui Prefectural Dinosaur Museum, 12: 17–33.
- Keith Holmes, W.B. & Anderson, H.M. 2013. A synthesis of the rich Gondwana Triassic megafossil flora from Nymboida, Australia. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 284–295.
- Kerr, R.A. 2013. Mega-eruptions drove the mother of mass extinctions. Science, 342(6165): 1424.
- Kettanah, Y.A. 2013. Hydrocarbon fluid inclusions in the Argo salt, offshore Canadian Atlantic margin. Canadian Journal of Earth Sciences, 50(6): 607–635.
- Khan, A., Anis & Shahnawaz. 2013. Petrography and provenance of Early Triassic Pachmarhi Formation sandstones, Satpura Gondwana Basin, Madhya Pradesh, Central India. Open Journal of Geology, 3(2): 83–93.
- Kilian, S. 2013. Bericht 2012 über geologische und strukturgeologische Aufnahmen im Karwendelgebirge auf Blatt 2223 Innsbruck und auf Blatt 2217 Hinteriß. Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1–4): 411–417.
- Kimmig, J. 2013. Possible secondarily terrestrial lifestyle in the European phytosaur *Nicrosaurus kapffi* (Late Triassic, Norian): a preliminary study. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 296–312.
- Klein, E.L., Angélica, R.S., Harris, C., Jourdan, F. & Babinski, M. 2013. Mafic dykes intrusive into Pre-Cambrian rocks of the São Luís cratonic fragment and Gurupi Belt (Parnaíba Province), north-northeastern Brazil: geochemistry, Sr-Nd-Pb-O isotopes, ⁴⁰Ar/³⁹Ar geochronology, and relationships to CAMP magmatism. Lithos, 172-173: 222–242.
- Klein, H. & Lucas, S.G. 2013. The Late Triassic tetrapod ichnotaxon *Apatopus lineatus* (Bock 1952) and its distribution. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 313–324.
- Klein, H., Niedźwiedzki, G., Voigt, S., Lagnaoui, A., Hminna, A., Saber, H. & Schneider, J.W. 2013. The tetrapod ichnogenus *Protochirotherium* Fichter and Kunz 2004, a characteristic Early Triassic morphotype of central Pangea. Ichnos, 20(1): 24–30.
- Klojzy-Karczmarczyk, B., Kryza, A., Kurek, T. & Mazurek, J. 2013. Analysis of changes in water chemistry of the Triassic aquifer in the initial phase of Trzebionka mine flooding. Biuletyn Pánstowowego Instytutu Geologicznego, 456:

281-286.

- Knaust, D. 2013. The ichnogenus *Rhizocorallium*: classification, tracemakers, palaeoenvironments and evolution. Earth-Science Reviews, 126: 1–47.
- Knaust, D. 2013. Classification of bioturbation-related reservoir quality in the Khuff Formation (Middle East): towards a genetic approach. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 247–267.
- Knies, J., Grasby, S.E., Beauchamp, B. & Schubert, C.J. 2013. Water mass denitrification during the latest Permian extinction in the Sverdrup Basin, Arctic Canada. Geology, 41(2): 167–170.
- Kobayashi, F. 2013. Late Permian (Lopingian) foraminifers from the Tsukumi Limestone, southern Chichibu Terrane of eastern Kyushu, Japan. Journal of Foraminiferal Research, 43(2): 154–169.
- Kokkolas, S., Kamberis, E., Xypolias, P., Sotiropoulos, S. & Koukouvelas, I. 2013. Coexistence of thin- and thick-skinned tectonics in the Zakynthos area (western Greece): insights from seismic sections and regional seismicity. Tectonophysics, 597-598: 73–84.
- Kolar-Jurkovšek, T., Vuks, V.J., Aljinović, D., Hautmann, M., Kaim, M. & Jurkovšek, B. 2013. Olenekian (Early Triassic) fossil assemblage from eastern Julian Alps (Slovenia). Annales Societatis Geologorum Poloniae, 83(3): 213–227.
- Komatsu, T., Shigeta, Y., Dang Tran Huyen, Dinh Cong Tien, Maekawa, T. & Tanaka, G. 2013. *Crittendenia* (Bivalvia) from the Lower Triassic (Olenekian) Bac Thay Formation, an Chau Basin, northern Vietnam. Paleontological Research, 17(1): 1–11.
- Konietzko-Meier, D. & Schmitt, A. 2013. A histological study of a femur of *Plagiosuchus*, a Middle Triassic temnospondyl amphibian from southern Germany, using thin sections and micro-CT scanning. Netherlands Journal of Geosciences, 92(2/3): 97–108.
- Kong-Yang Zhu, Zheng-Xiang Li, Xi-Sheng Xu & Wilde, S.A. 2013. Late Triassic melting of a thickened crust in southeastern China: evidence for flat-slab subduction of the Paleo-Pacific plate. Journal of Asian Earth Sciences, 74: 265–279.
- Kongyou Wu, Paton, D. & Ming Zha. 2013. Unconformity structures controlling stratigraphic reservoirs in the northwest margin of Junggar basin, north-west China. Frontiers of Earth Science, 7(1): 55–64.
- Konietzko-Meier, D. & Klein, N. 2013. Unique growth pattern of *Metoposaurus diagnosticus krasiejowensis* (Amphibia, Temnospondyli) from the Upper Triassic of Krasiejów, Poland. Palaeogeography, Palaeoclimatology, Palaeoecology, 370: 145–157.
- Konietzko-Meier, D. & Sander, P.M. 2013. Long bone histology of *Metaposaurus diagnosticus* (Temnospondyli) from the Late Triassic of Krasiejów (Poland) and its paleobiological implications. Journal of Vertebrate Paleontology, 33(5): 1003–1018.
- Konstantinov, A.G., Sobolev, E.S. & Yadrenkin, A.V. 2013. Triassic stratigraphy of the eastern Laptev Sea coast and New Siberian Islands. Russian Geology and Geophysics, 54(8):

792–807.

- Kontorovich, V.A., Kontorovich, A.E., Gubin, I.A., Zoteev, A.M., Lapkovsky, V.V., Malyshev, N.A., Soloviev, M.V. & Fradkin, G.S. 2013. The Neoproterozoic-Phanerozoic section of the Anabar-Lena province: structural framework, geological model, and petroleum potential. Russian Geology and Geophysics, 54(8): 980–996.
- Korn, D., Hopkins, M.J. & Walton, S.A. 2013. Extinction space – a method for the quantification and classification of changes in morphospace across extinction boundaries. Evolution, 67(10): 2795–2810.
- Korn, D., Leda, L., Ghaderi, A., Hairapetian, V. & Schobben, M. 2013. The Lilliput Effect in the latest Permian ammonoids from Iran. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 41-42.
- Korngreen, D. & Benjamini, C. 2013. Stratigraphic evidence for shear in structural development of the Triassic Levant margin: new borehole data on the epicontinental to deep marine transition in Israel. Tectonophysics, 591: 3–15.
- Korngreen, D., Orlov-Labkovsky, O., Bialik, O. & Benjamini, C. 2013. The Permian-Triassic transition in the central coastal plain of Israel (North Arabian Plate Margin), David 1 Borehole. Palaios, 28(8): 491–508.
- Kotánski, Z. 2013. Anisian Dasycladales from Upper Silesia and adjacent regions. Carnets de Géologie, CG2013 Book 2, 205 pp.
- Kowal-Linka, M. & Stanikowski, W. 2013. Garnet and tourmaline as provenance indicators of terrigenous material in epicontinental carbonates (Middle Triassic, S Poland). Sedimentary Geology, 291: 27–47.
- Kozur, H.W., Franz, M. & Bachmann, G.H. 2013. Shipingia weemsi n. sp., a biostratigraphically important conchostracan species from the uppermost Carnian and lowermost Norian of central Europe. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 325–330.
- Krajewski, K.P. 2013. Organic matter-apatite-pyrite relationships in the Botneheia Formation (Middle Triassic) of eastern Svalbard: relevance to the formation of petroleum source rocks in the NW Barents Sea shelf. Marine and Petroleum Geology, 45: 69–105.
- Krapovickas, V., Mancuso, A.C., Marsicano, C.A., Domnanovich, N. & Schultz, C.L. 2013. Large tetrapod burrows from the Middle Triassic of Argentina: a behavioural adaptation to seasonal semi-arid climate? Lethaia, 46(2): 154–169.
- Kraus, S.H., Brandner, R., Heubeck, C., Kozur, H.W., Struck, U. & Korte, C. 2013. Carbon isotope signatures of latest Permian marine successions of the Southern Alps suggest a continental runoff pulse enriched in land plant material. Fossil Record, 16(1): 97–109.
- Kubo, T. & Kubo, M.O. 2013. Analysis of Triassic archosauriform trackways: difference in stride/foot ratio between dinosauromorphs and archosauriforms. Palaios, 28(4): 259–265.
- Kui-Dong Zhao, Shao-Yong Jiang, Wei-Feng Chen, Pei-Rong Chen & Hong-Fei Ling. 2013. Zircon U-Pb chronology and

elemental and Sr-Nd-Hf isotope geochemistry of two Triassic A-type granites in South China: implication for petrogenesis and Indosinian transtensional tectonism. Lithos, 160-161: 292–306.

- Kürschner, W., Batenburg, S.J. & Mander, L. 2013. Aberrant *Classopollis* pollen reveals evidence for unreduced (2n) pollen in the conifer family Cheirolepidiaceae during the Triassic– Jurassic transition. Proceedings of the Royal Society, B. 280(1768) 20131708; doi: 10.1098/rspb.2013.1708 (8pp).
- Kusov, A.V. & Stoupakova, A.V. 2013. Reservoir sequences of the Anabar-Khatanga saddle: formation conditions and reservoir properties. Moscow University Geology Bulletin, 68(3): 185–190.
- Kustatscher, E. & van Konijnenburg-van Cittert, J.H.A. 2013. Seed ferns from the European Triassic – an overview. *In*, Tanner, L.H., Speilmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 331–344.
- Labandeira, C.C., & Currano, E.D. 2013. The fossil record of plant-insect dynamics. Annual Review of Earth and Planetary Sciences, 41: 287–311.
- Laishi Zhao, Yonglin Chen, Zhong-qiang Chen & Ling Cao. 2013. Uppermost Permian to Lower Triassic conodont zonation from Three Gorges area, South China. Palaios, 28(8): 523–540.
- Laishi Zhao, Zhong-Qiang Chen, Algeo, T.J., Jianbo Chen, Yonglin Chen, Jinnan Tong, Shan Gao, Lian Zhou, Zhaochu Hu & Yongsheng Liu. 2013. Rare-earth element patterns in conodont albid crowns: evidence for massive inputs of volcanic ash during the latest Permian biocrisis? Global and Planetary Change, 105: 135–151.
- Laming, D.J.C. & Roche, D.P. 2013. Faulting in Permo-Triassic strata and buried channels revealed by excavation of the Lympstone – Powderham pipeline tunnel, Exe Estuary, Devon (Abstract). Geoscience in South-West England, 13(2): 244.
- Langer, M.C. & Ferigolo, J. 2013. The Late Triassic dinosauromorph *Sacisaurus agudoensis* (Caturrita Formation: Rio Grande do Sul, Brazil): Anatomy and affinities. Geological Society, London, Special Publications, 379: 353–392.
- Langer, M.C., Nesbitt, S.J., Bittencourt, J.S. & Irmis, R.B. 2013. Non-dinosaurian Dinosauromorpha. Geological Society, London, Special Publications, 379: 157–186.
- Leavitt, W.D., Halevy, I., Bradley, A.S. & Johnston, D.T. 2013. Influence of sulfate reduction rates on the Phanerozoic sulfur isotope record. Proceedings of the National Academy of Sciences, 110(28): 11244–11249.
- Legendre, L.J., Segaten, L. & Cubo, J. 2013. Evidence for high bone growth rate in *Euparkeria* obtained using a new paleohistological inference model for the humerus. Journal of Vertebrate Paleontology, 33(6): 1343–1350.
- Lei He, Yongbiao Wang, Woods, A., Guoshan Li, Hao Yang & Wei Liao. 2013. An oxygenation event occurred in deep shelf settings immediately after the end-Permian mass extinction in South China. Global and Planetary Change, 101: 72–81.
- Leitner, C., Neubauer, F., Genser, J., Borojević-Šoštarić, S. & Rantitsch, G. 2013. ⁴⁰Ar/³⁹Ar ages of crystallization and

recrystallization of rock-forming polyhalite in Alpine rocksalt deposits. Geological Society, London, Special Publications, 378: 207–224.

- Leitner, C., Neubauer, F., Marschallinger, R., Genser, J. & Bernroider, M. 2013. Origin of deformed halite hopper crystals, pseudomorphic anhydrite cubes and polyhalite in Alpine evaporites (Austria, Germany). International Journal of Earth Sciences, 102(3): 813–829.
- Lepper, H. & Röhling, H.-G. (Eds). 2013. Stratigraphy von Deutschland XI: Buntsandstein. Schiftenreihe der Deutschen Gesellschaft für Geowissenschaften, 69: 657 pp.
- Letnikova, E.F., Lobanov, S.S., Pokhilenko, N.P., Izokh, A.E. & Nikolenko, E.I. 2013. Sources of clastic material in the Carnian diamond-bearing horizon of the northeastern part of the Siberian Platform. Doklady Earth Sciences, 451(1): 702–705.
- Li, S.Z., Suo, Y.H., Santosh, M., Dai, L.M., Liu, X., Yu, S., Zhao, S.J. & Jin, C. 2013. Mesozoic to Cenozoic intracontinental deformation and dynamics of the North China Craton. Geological Journal, 48(5): 543–560.
- Li Wei & Wen Zhi-gang. 2013. Characteristics and assessment of the Chang-71 reservoirs from the Yanchang Formation in the Maling region, northern Shaanxi. Sedimentary Geology and Tethyan Geology, 33(3): 40–47.
- Li Yan, Wang Yun-peng, Zhao Chang-yi, Lu Jia-lan, Liao Lingling & Liu Jie. 2013. Triassic Xujiahe burial history of source rocks and thermal evolution of the Sichuan Basin. Geological Bulletin of China, 32(11): 1798–1805.
- Li Yong, Wang Ping & Sun Ming. 2013. Sedimentary features and reservoir characteristics of Chang 6 strata of Fanchuan district, Zhidan oilfield. Northwestern Geology, 46(3): 174–182.
- Lianbo Zeng, Hui Su, Xiaomei Tang, Yongmin Peng & Lei Gong. 2013. Fractured tight sandstone oil and gas reservoirs: a new play type in the Dongpu depression, Bohai Bay Basin, China. AAPG Bulletin, 97(3): 363–377.
- Liang Luo, Dong Jia, Jiafu Qi, Guoqi Wei & Fei Deng. 2013. Tectono-sedimentary evolution of the Late Triassic Xujiahe Formation in the Sichuan Basin. Acta Geologica Sinica, 87(6): 1554–1568.
- Liao Shi-yong, Yin Fu-guang, Sun Zhi-ming, Wang Dong-bing, Tang Yuan & Sun Jie. 2013. Western Yunnan Baoshan eastern edge of the Late Triassic Ho Chu Ci dacite porphyry discovery and its geological significance. Geological Bulletin of China, 32(7): 1006–1013.
- LiMei Tang, HanLin Chen, ChuanWan Dong, ShuFeng Yang, ZhongYue Shen, XiaoGan Cheng & LuLu Fu. 2013. Middle Triassic post-orogenic extension on Hainan Island: Chronology and geochemistry constraints of bimodal intrusive rocks. SCIENCE CHINA Earth Sciences, 56(5): 783–793.
- Lintnerová, O., Michalik, J., Uhlík, P., Soták, J. & Weissová, Z. 2013. Latest Triassic climate humidification and kaolinite formation (Western Carpathians, Tatric Unit of the Tatra Mts.). Geological Quarterly, 57: 701–728.
- Liparini, A., Oliveira, T.V., Pretto, F.A., Soares, M.B. & Schultz, C.L. 2013. The lower jaw and dentition of the traversodontid *Exaeretodon riograndensis* Abdala, Barberena & Dornelles,

from the Brazilian Triassic (Santa Maria 2 Sequence, Hyperodapedon Assemblage). Alcheringa, 37(3): 331-337.

- Liparini, A. & Schultz, C.L. 2013. A reconstruction of the thigh musculature of the extinct pseudosuchian *Prestosuchus chiniquensis* from the *Dinodontosaurus* Assemblage Zone (Middle Triassic Epoch), Santa Maria 1 Sequence, southern Brazil. Geological Society, London, Special Publications, 379: 441–468.
- Liu Jun. 2013. On the taxonomy of *Xinpusaurus* (Reptilia: Thallatosauria). Vertebrata Palasiatica, 51(1): 17-23.
- Liu Jun, Li Lu & Li Xing-wen. 2013. SHRIMP U-Pb zircon dating of the Triassic Ermaying and Tongchuan formations in Shanxi, China and its stratigraphic implications. Vertebrata Palasiatica, 51(2): 162–168.
- Liu Xingjun, Liu Yiqun, Zhou Dingwu, Li Hong, Chang Xinhu & Nan Yun. 2013. Deep fluid tracer in Ordos Basin: Characteristics and origin of high natural gamma sandstone in Triassic Yanchang Formation. Earth Science Frontiers, 20(5): 149–165.
- Lloyd, G.T. & Friedman, M. 2013. A survey of palaeontological sampling biases in fishes based on the Phanerozoic record of Great Britain. Palaeogeography, Palaeoclimatology, Palaeoecology, 372: 5–17.
- Lockley, M.G., Li Jianjun, Li Rihui, Matsukawa, M., Harris, J.D. & Xing Lida. 2013. A review of the tetrapod track record in China, with special reference to type ichnospecies: implications for ichnotaxonomy and paleobiology. Acta Geologica Sinica, 87(1): 1–20.
- Lockley, M.G. & Lucas, S.G. 2013. *Evazoum gatewayensis*, a new Late Triassic archosaurian ichnospecies from Colorado: implications for footprints in the ichnofamily Otozoidae. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 345–353.
- Löffler, N. & Prinz-Grimm, P. 2013. Quantitative analyse von Wirbeltierfragmenten aus einem Muschelkalk-Keuper-Bonebed bei Palzem (Mosel). Mainzer geowissenschaftliche Mitteilungen, 41: 37–50.
- Lombardo, C. 2013. A new basal actinopterygian fish from the Late Ladinian of Monte San Giorgio (Canton Ticino, Switzerland). Swiss Journal of Geology, 106(2): 219–230.
- Loope, G.R., Kump, L.R. & Arthur, M.A. 2013. Shallow water redox conditions from the Permian-Triassic boundary microbialite: the rare earth element and iodine geochemistry of carbonates from Turkey and South China. Chemical Geology, 351: 175–194.
- Lucas, S.G. 2013. Plant megafossil biostratigraphy and biochronology, Upper Triassic Chinle Group, western USA. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 354–365.
- Lucas, S.G. 2013. A new Triassic timescale. *In*, Tanner, L.H., Spielmann, J. A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin

61: 366–374.

- Lucas, S.G. & Bachmann, G.H. 2013. Dedication to Heinz W. Kozur. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 1–22.
- Lucas, S.G. & Spielmann, J.A. 2013. Magnetostratigraphy of the Upper Triassic Chinle Group in New Mexico: an appraisal of 40 years of analysis. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61:375–381.
- Lucas, S.G., Spielmann, J.A. & Hunt, A.P. 2013. A new doswelliid archosauromorph from the Upper Triassic of West Texas. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 382-388.
- Lucas, S.G., Spielmann, J.A. & Rinehart, L.F. 2013. Juvenile skull of the phytosur *Redondasaurus* from the Upper Triassic of New Mexico, and phytosaur ontogeny. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 389–400.
- Lutz, A.I. & Arce, F.E. 2013. Paleoflora of the Carrizal Formation (Middle-Upper Triassic), San Juan Province, Argentina. Revista Mexicana de Ciencias Geológicas, 30(2): 453–462.
- Maisch, M.W., Matzke, A.T. & Rathgeber, T. 2013. Reevaluation of the enigmatic archosaur *Dyoplax arenaceus* O. Fraas, 1867 from the Schilfsandstein (Stuttgart Formation, lower Carnian, Upper Triassic) of Stuttgart, Germany. Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen, 267(3): 353–362.
- Manafi, M., Arian, M., Reesi, S.H.T. & Solgi, A. 2013. Tethys subduction history in Caucasus region. Open Journal of Geology, 3(3): 222–232.
- Mander, L., Kürschner, W.M. & McElwain, J.C. 2013. Palynostratigraphy and vegetation history of the Triassic-Jurassic transition in East Greenland. Journal of the Geological Society, London, 170(1): 37–46.
- Mao Luo, Zhong-qiang Chen, Shixue Hu, Qiyue Zhang, Benton, M.J., Wen Wen & Jin-yuan Huang. 2013. Carbonate reticulated ridge structures from the lower Middle Triassic of the Luoping area, Yunnan, southwestern China: geobiological features and implications for exceptional preservation of the Luoping biota. Palaios, 28(8): 541–551.
- Marfil, R., La Iglesia, A. & Estupiñan, J. 2013. Origin and nature of the aluminium phosphate-sulphate minerals (APS) associated with uranium mineralization in Triassic red-beds (Iberian Range, Spain). Estudios Geológicas, 69(1): 31–34.
- Martindale, R.C., Krystyn, L., Bottjer, D.J., Corsetti, F.A., Senowbari-Daryan, B. & Martini, R. 2013. Depth transect of an Upper Triassic (Rhaetian) reef from Gosau, Austria: microfacies and community ecology. Palaeogeography, Palaeoclimatology, Palaeoecology, 376: 1–21.
- Martindale, R.C., Krystyn, L., Corsetti, F.A. & Bottjer, D.J.
2013. From fore reef to lagoon: evolution of the Upper Triassic Dachstein carbonate platform on the Tennengebirge (Salzburg, Austria). Palaios, 28(11): 755–770.

- Martinez, R.N., Apaldetti, C., Colombi, C.E., Praderio, A., Fernandez, E., Malnis, P.S., Correa, G.A., Abelin, D. & Alcober, O. 2013. A new sphenodontian (Lepidosauria: Rhynchocephalia) from the Late Triassic of Argentina and the early origin of herbivore opisthodontians. Proceedings of the Royal Society, B. 280(1772) 20132057; doi: 10.1098/ rspb.2013.2057 (7 pp).
- Martinez, R.N., Fernandez, E. & Alcober, O.A. 2013. A new non-mammaliaform eucynodont from the Carnian-Norian Ischigualasto Formation, northwestern Argentina. Revista Brasileira de Paleontologia, 16(1): 61–76.
- Marynowski, L. 2013. Mesozoic wildfires: distribution and importance for global events. Przegląd Geologiczny, 60(3): 148–154.
- Masoodi, M., Yassaghi, A., Sadat, M.A.A.N., Neubauer, F., Bernroider, M., Friedl, G., Genser, J. & Houshmandzadeh, A. 2013. Cimmerian evolution of the central Iranian basement: evidence from metamorphic units of the Kashmar-Kerman tectonic zone. Tectonophysics, 588: 189–208.
- Mastrantonio, B.M., Schultz, C.L., Desojo, J.B. & Garcia, J.B. 2013. The braincase of *Prestosuchus chiniquensis* (Archosauria: Suchia). Geological Society, London, Special Publications, 379: 425–440.
- Maxwell, E.E. & Kear, B.P. 2013. Triassic ichthyopterygian assemblages from the Svalbard archipelago: a reassessment of taxonomy and distribution. GFF, 135(1): 85–94.
- Maystrenko, Y.P., Bayer, U. & Scheck-Wenderoth, M. 2013. Salt as a 3D element in structural modelling – example fron the Central European Basin System. Tectonophysics, 591: 62–82.
- Mazzoli, S., Martín-Algarra, A., Reddy, S.M., López Sánchez-Vizcaíno, V., Fedele, L. & Noviello, A. 2013. The evolution of the footwall to the Ronda subcontinental mantle peridotites: insights from the Nieves Unit (western Betic Cordillera). Journal of the Geological Society, London, 170(3): 385–402.
- Medvedev, A.Ya. 2013. Platinum group elements in Permo-Triassic volcanics in West Siberia. Russian Geology and Geophysics, 54(7): 664–670.
- McGowan, A.J. & Kiessling, W. 2013. Using abundance data to assess the relative role of sampling biases and evolutionary radiations in Upper Muschelkalk ammonoids. Acta Palaeontologica Polonica, 58(3): 561–572.
- McHone, J.G. & Fyffe, L. 2013. Geology of the Island of Grand Manan, New Brunswick: Precambrian to Early Cambrian and Triassic formations. Geoscience Canada, 40(4): 361–362.
- MacDougall, M.J., Modesto, S.P. & Botha-Brink, J. 2013. The postcranial skeleton of the Early Triassic parareptile *Sauropareion anoplus*, with a discussion of possible life history. Acta Palaeontologica Polonica, 58(4): 737–749.
- Meinhold, G. & Kostopoulos, D.K. 2013. The Circum-Rhodope Belt, northern Greece: age, provenance, and tectonic setting. Tectonophysics, 595-596: 55-68.
- Meister, P., Mckenzie, J.A., Bernasconi, S.M. & Brack, P. 2013. Dolomite formation in the shallow seas of the Alpine Triassic. Sedimentology, 60(1): 270–291.

- 71
- Mercedes-Martin, R., Salas, R. & Arenas, C. 2013. Facies heterogeneity and depositional models of a Ladinian (Middle Triassic) microbial-dominated carbonate ramp system (Catalan Coastal Ranges, NE Spain). Marine and Petroleum Geology, 46: 107–128.
- Metcalfe, I. 2013. Gondwana dispersion and Asian accretion: tectonic and palaeogeographic evolution of eastern Tethys. Journal of Asian Earth Sciences, 66: 1–33.
- Metcalfe, I. 2013. Tectonic evolution of the Malay Peninsula. Journal of Asian Earth Sciences, 76: 195–213.
- Metcalfe, I., Nicoll, R.S., Willink, R., Ladjavadi, M. & Grice, K. 2013. Early Triassic (Induan–Olenekian) conodont biostratigraphy, global anoxia, carbon isotope excursions and environmental perturbations: new data from Western Australian Gondwana. Gondwana Research, 23(3): 1136– 1150.
- Meyer, K.M., Yu, M., Lehrmann, D., van de Schootbrugge, B. & Payne, J.L. 2013. Constraints on Early Triassic carbon cycle dynamics from paired organic and inorganic carbon isotope records. Earth and Planetary Science Letters, 361: 429–435.
- Michalik, J., Lintnerova, O., Wojcik-Tabol, P., Gazdzicki, A., Grabowski, J., Golej, M., Simo, V. & Zahradnikova, B. 2013. Paleoenvironments during the Rhaetian transgression and the colonization history of marine biota in the Fatric Unit (Western Carpathians). Geologica Carpathica, 64(1): 39–62.
- Mikhalevich, V.I. 2013. New insight into the systematics and evolution of the foraminifera. Micropaleontology, 59(6): 493–527.
- Miller, E.L., Soloviev, A.V., Prokopiev, A.V., Toro, J., Harris, D., Kuzmichev, A.B. & Gehrels, G.E. 2013. Triassic river systems and the paleo-Pacific margin of northwestern Pangea. Gondwana Research, 23(4): 1631–1645.
- Mirnejad, H., Lalonde, A.E., Obeid, M. & Hassanzadeh, J. 2013. Geochemistry and petrogenesis of Mashhad granitoids: An insight into the geodynamic history of the Paleo-Tethys in northeast of Iran. Lithos, 170–171: 105–116.
- Mohriak, W.U., Danforth, A., Post, P.J., Brown, D.E., Tari, G.C., Nemčok, M. & Sinha, S.T. (eds.). 2013. Conjugate Divergent Margins. Geological Society, London, Special Publications, 369: vi+568 pp.
- Moisan, P. & Voigt, S. 2013. Lycopsids from the Madygen Lagerstätte (Middle to Late Triassic, Kyrgyzstan, Central Asia). Review of Palaeobotany and Palynology, 192: 42–64.
- Moix, P., Vachard, D., Allibon, J., Martini, R., Wernli, R., Kozur,
 H.W. & Stampfli, G.M. 2013. Palaeotethyan, Neotethyan and Huğlu-Pindos Series in the Lycian Nappes (SW Turkey): geodynamical implications. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 401–444.
- Monnet, C., Bucher, H., Brayard, A. & Jenks, J.F. 2013. *Globacrochordiceras* gen. nov. (Acrochodiceratidae, late Early Triassic) and its significance for stress-induced evolutionary jumps in ammonoid lineages (cephalopods). Fossil Record, 16(2): 197–215.
- Monostori, M. & Tóth, E. 2013. Ladinian (Middle Triassic) silicified ostracod faunas from the Balaton Highland

(Hungary). Rivista Italiana di Paleontologia e Stratigrafia, 119(3): 303–23.

- Montefeltro, F.C., Bittencourt, J.S., Langer, M.C. & Schultz, C.L. 2013. Postcranial anatomy of the hyperodapedontine rhynchosaur *Teyumbaita sulcognathus* (Azevedo and Schultz, 1987) from the Late Triassic of southern Brazil. Journal of Vertebrate Paleontology, 33(1): 67–84.
- Moonsup Cho, Wonseok Cheong, Ernst, W.G., Keewook Yi & Jeongmin Kim. 2013. SHRIMP U-Pb ages of detrital zircons in metasedimentary rocks of the central Ogcheon fold-thrust belt, Korea: evidence for tectonic assembly of Paleozoic sedimentary protoliths. Journal of Asian Earth Sciences, 63: 24–249.
- Mørk, M.B.E. 2013. Diagenesis and quartz cement distribution of low-permeability Upper Triassic–Middle Jurassic reservoir sandstones, Longyearbyen CO₂ lab well site in Svalbard, Norway. AAPG Bulletin, 97(4): 577–596.
- Morton, A., Hounslow, M.W. & Frei, D. 2013. Heavymineral, mineral-chemical and zircon-age constraints on the provenance of Triassic sandstones from the Devon coast, southern Britain. Geologos, 19(1–2): 67–85.
- Nakae, S. 2013. Triassic and Middle Jurassic radiolarians from pelagic cherts in the Nanjö Mountains, southwest Japan Part 1. Imajö district. Bulletin of the Geological Survey of Japan, 64(3–4): 85–112.
- Nakae, S. 2013. Triassic and Middle Jurassic radiolarians from pelagic cherts in the Nanjö Mountains, southwest Japan – Part
 2. Kanmuri Yama district. Bulletin of the Geological Survey of Japan, 64(5–6): 151–190.
- Naydenov, K., Peytcheva, I., von Quadt, A., Sarov, S., Kolcheva, K. & Dimov, D. 2013. The Maritsa strike-slip shear zone between Kostenets and Krichin towns, south Bulgaria: structural, petrographic and isotope geochronology study. Tectonophysics, 595–596: 69–89.
- Naylor, D., Al-Rawi, M., Clayton, G., Fitzpatrick, M.J. & Green, P.F. 2013. Hydrocarbon potential in Jordan. Journal of Petroleum Geology, 36(3): 205–236.
- Neenan, J.M. & Scheyer, T.M. 2013. Systematics, origins and palaeoecology of placodont marine reptiles. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 51–52.
- Nesbitt, S.J., Barrett, P.M., Werning, S., Sidor, C. & Charig, A.J. 2013. The oldest dinosaur? A Middle Triassic dinosauriform from Tanzania. Biology Letters, 9(1), doi: 10.1098/ rsbl.2012.0949 (5 pp).
- Nesbitt, S.J., Brusatte, S.L., Desojo, J.B., Liparini, A., de Franca, M.A.G., Weinbaum, J.C. & Gower, D.J. 2013. Rauisuchia. Geological Society, London, Special Publications, 379: 241–274
- Nesbitt, S.J. & Butler, R.J. 2013. Redescription of the archosaur *Parringtonia gracilis* from the Middle Triassic Manda beds of Tanzania, and the antiquity of Erpetosuchidae. Geological Magazine, 150(2): 225–238.
- Nesbitt, S.J. & Butler, R.J. 2013. Redescription of the archosaur *Parringtonia gracilis* from the Middle Triassic Manda beds of Tanzania, and the antiquity of Erpetosuchidae CORRIGENDUM. Geological Magazine, 150(2): 239.

- Nesbitt, S.J., Butler, R.J. & Gower, D.J. 2013. A new archosauriform (Reptilia: Diapsida) from the Manda Beds (Middle Triassic) of southwestern Tanzania. PLoS ONE, 8(9): e72753. doi:10.1371/journal.pone.0072753.
- Nesbitt, S.J., Desojo, J.B. & Irmis, R.B. (eds.). 2013. Anatomy, Phylogeny and Palaeobiology of Early Archosaurs and their Kin. Geological Society, London, Special Publications, 379, vi+608 pp.
- Nesbitt, S.J., Desojo, J.B. & Irmis, R.B. 2013. Anatomy, phylogeny and palaeobiology of early archosaurs and their kin. Geological Society, London, Special Publications, 379: 1–7.
- Nestell, M.K. & Stevens, C.H. 2013. Mixed Tethyan and McCloud Belt rugose corals and fusulinids in an Upper Triassic conglomerate, central Oregon. Journal of Paleontology, 87(5): 909–921.
- Nguyen, B.T.T., Jones, S.J., Goulty, N.R., Middleton, A.J., Grant, J., Ferguson, A. & Bowen, L. 2013. The role of fluid pressure and diagenetic cements for porosity preservation in Triassic fluvial reservoirs of the Central Graben, North Sea. AAPG Bulletin, 97(8): 1273–1302.
- Nichols, K. & Lucas, S.G. 2013. Norman J. Silberling (1928-2011) In memoriam. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 23–30.
- Niedźwiedzki, G., Brusatte, S.L. & Butler, R.J. 2013. *Prorotodactylus* and *Rotodactylus* tracks: an ichnological record of dinosauromorphs from the Early-Middle Triassic of Poland. Geological Society, London, Special Publications, 379: 319–351.
- Niu Ya-zhuo, Jiang Bao-yu & Zhang Yan-wei. 2013. Late Triassic and Early Jurassic bivalve fossils from eastern South China and their biostratigraphic implications. Acta Palaeontologica Sinica, 52(3): 352–376.
- Novikov, I.S. 2013. Reconstructing the stages of orogeny around the Junggar basin from the lithostratigraphy of Late Paleozoic, Mesozoic, and Cenozoic sediments. Russian Geology and Geophysics, 54(2): 138–152.
- Nurnberg, S. & Aberhan, M. 2013. Habitat breadth and geographic range predict diversity dynamics in marine Mesozoic vertebrates. Paleobiology, 39(3): 360-372.
- Nutman, A.P., Buckman, S., Hidaka, H., Kamiichi, T., Belousova, E. & Aitchison, J. 2013. Middle Carboniferous – Early Triassic eclogite-blueschist blocks within a serpentinite mélange at Port Macquarie, eastern Australia: implications for the evolution of Gondwana's eastern margin. Gondwana Research, 24(3–4): 1038–1050.
- Nützel, A. 2013. Revision of genus *Sabrinella* (Gastropoda) and a new minute vetigastropod from the Late Triassic Cassian Formation (N Italy). Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen, 269(1): 63–72.
- Ocampo-Díaz, Y.Z.E. & Rubio-Ciseneros, I.I. 2013. Sedimentary recycling: compositional analysis associated with unconformities in northwestern Mexico, an Upper Triassic to Lower Cretaceous example. Boletín de la Sociedad Geológica Mexicana, 65: 553–572.

Okrostsvaridze, A.V. & Tormey, D.R. 2013. Phanerozoic

continental crust evolution of the Inner Caucasian Microplate: the Dzirula massif. Episodes, 36(1): 31–38.

- Oliver, G. & Prave, A. 2013. Palaeogeography of Late Triassic red-beds in Singapore and the Indosinian Orogeny. Journal of Asian Earth Sciences, 76: 214–224.
- Oosterhout, C.W.M. van & Pöppelreiter, M.C. 2013. Plate reconstructions and distribution of sedimentary facies during the Permo-Triassic opening of the Neo-Tethys ocean. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 23–42.
- Oosterink, H.W. & Winkelhorst, H. 2013. Probable remains of jellyfish (Cnidaria, Scyphozoa) from the lower Middle Triassic (Anisian) of Winterswijk, eastern Netherlands. Netherlands Journal of Geosciences, 92(1): 61–67.
- Orchard, M.J. 2013. Five new genera of conodonts from the Carnian-Norian boundary beds of Black Bear Ridge, northeast British Columbia, Canada. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 445–457.
- Osen, A.K., Winguth, A.M.E., Winguth, C. & Scotese, C.R. 2013. Sensitivity of Late Permian climate to bathymetric features and implications for the mass extinction. Global and Planetary Change, 105: 171–179.
- Ősi, A., Botfalvai, G., Prondvai, E., Hajdu, Z., Czirják, G., Szentesi, Z., Pozsgai, E., Götz, A.E., Makádi, L., Csengődi, D. & Sebe, K. 2013. First report of Triassic vertebrate assemblages from the Villány Hills (southern Hungary). Central European Geology, 56(4): 297–335.
- Otero, A. 2013. Functional anatomy in the transition Prosauropoda-Sauropoda. The Palaeontological Association Newsletter, 82: 72–74.
- Otero, A. & Pol, D. 2013. Postcranial anatomy and phylogenetic relationships of *Mussaurus patagonicus* (Dinosauria, Sauropodamorpha). Journal of Vertebrate Paleontology, 33(5): 1138–1168.
- Özbey, Z., Ustaömer, T. & Robertson, A.H.F. 2013. Mesozoic magmatic and sedimentary development of the Tavşanlı Zone (NW Turkey): implications for rifting, passive margin development and ocean crust emplacement. Geological Society, London, Special Publications, 372: 141–165.
- Özdamar, Ş., Billor, M.Z., Sunal, G., Esenli, F. & Roden, M.F. 2013. First U-Pb SHRIMP zircon and ⁴⁰Ar/³⁹Ar ages of metarhyolites from the Afyon – Bolkardag Zone, SW Turkey: implications for the rifting and closure of the Neo-Tethys. Gondwana Research, 24(1): 377–391
- Papanikolaou, D. 2013. Tectonostratigraphic models of the Alpine terranes and subduction history of the Hellenides. Tectonophysics, 595-596: 1–24.
- Parker, W.G. & Nesbitt, S.J. 2013. Cranial remains of *Poposaurus gracilis* (Pseudosuchia, Poposauroidea) from the Upper Triassic, the distribution of the taxon, and its implications for poposauroid evolution. Geological Society, London, Special Publications, 379: 503–523.
- Parrish, J.T. & Soreghan, G.S. 2013. Sedimentary geology and the future of paleoclimate studies. Sedimentary Record, 11(2): 4–10.

- Paul, J. 2013. Die Beckenfazies des Buntsandsteins: playaund fluss-ablagerungen, stromatolithen und klimazyklen (Exkursion C am 4. April 2013) [The basin facies of the Buntsandstein: playa and fluvial deposits, stromatolites and climatic cycles]. Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereins, 95: 49–70.
- Payne, J.L., Jost, A.B., Wang, S.C. & Skotheim, J.M. 2013. A shift in the long-term mode of foraminiferan size evolution caused by the end-Permian mass extinction. Evolution, 67(3): 816–827.
- Pearson, M.R., Benson, R.B.J., Upchurch, P., Fröbisch, J. & Kammerer, C.F. 2013. Reconstructing the diversity of early terrestrial herbivorous tetrapods. Palaeogeography, Palaeoclimatology, Palaeoecology, 372: 42–49.
- Peecook, B.R., Sidor, C.A., Nesbitt, S.J., Smith, R.M.H., Steyer, S.J. & Angielczyk, K.D. 2013. A new silesaurid from the upper Ntawere Formation of Zambia (Middle Triassic) demonstrates the rapid diversification of Silesauridae (Avemetatarsalia, Dinosauriformes). Journal of Vertebrate Paleontology, 33(5): 1127–1137.
- Pereira, R. & Alves, T.M. 2013. Crustal deformation and submarine canyon incision in a Meso-Cenozoic firstorder transfer zone (SW Iberia, North Atlantic Ocean). Tectonophysics, 601: 148–162.
- Perez-Gomez, I., Farqani, S., Claps, M. & Peters, H. 2013. Diagenetic evolution of a Permo-Triassic Khuff oolitic reservoir in north Oman. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 327–386.
- Perri, F., Critelli, S., Martín-Algarra, A., Martín-Martín, M., Perrone, V., Mongelli, G. & Zattin, M. 2013. Triassic redbeds in the Malaguide Complex (Betic Cordillera–Spain): petrography, geochemistry and geodynamic implications. Earth-Science Reviews, 117: 1–28.
- Petersen, H.I., Lindström, S., Therkalsen, J. & Pedersen, G.K. 2013. Deposition, floral composition and sequence stratigraphy of uppermost Triassic (Rhaetian) coastal coals, southern Sweden. International Journal of Coal Geology, 116–117: 117–134.
- Petti, F.M., Bernardi, M., Kustatscher, E., Renesto, S. & Avanzini, M. 2013. Diversity of terrestrial tetrapods and plants in the Triassic of the Southern Alps: ichnological, paleozoological and paleobotanical evidence. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 448–484.
- Pla, C., Márquez-Aliaga, A. & Botella, H. 2013. The chondrichthyan fauna of the Middle Triassic (Ladinian) of the Iberian Range (Spain). Journal of Vertebrate Paleontology, 33(4): 770–785.
- Plasencia, P., Márquez-Aliaga, A. & Jingeng Sha. 2013. A database of Triassic conodonts from a comprehensive revision of literature. Revista española de paleontología, 28(2): 215–216.
- Pledge, N.S. 2013. First South Australian labyrinthodont: a possible chigutisaurid stereospondyl amphibian from the Late Triassic of Leigh Creek. Transactions of the Royal Society of South Australia, 137(1): 127–134.

- Pledge, N.S. & Baulch, J.C.I. 2013. New fish from the Late Triassic Leigh Creek Coal Measures, South Australia. Transactions of the Royal Society of South Australia, 137(1): 90–96.
- Pomoni, F.A. & Tselelpidis, V. 2013. Lithofacies paleogeography and biostratigraphy of the lowermost horizons of the Middle Triassic Hallstatt Limestones (Argolis Peninsula, Greece). Journal of Palaeogeography, 2(3): 252-274.
- Pöppelreiter, M.C. (ed.). 2013. Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv, 409 pp.
- Pöppelreiter, M.C. & Obermaier, M. 2013. Regional top seal of the Khuff reservoir: the Sudair Formation. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 387–399.
- Poros, Z., Machel, H.G., Mindszenty, A. & Molnár, F. 2013. Cryogenic powderization of Triassic dolostones in the Buda Hills, Hungary. International Journal of Earth Sciences, 102(5): 1513–1539.
- Post, P.J., Elliott, E.T., Klazynski, R.J., Klocek, E.S., Decort, T.M., Riches, T.J. & Kun Li. 2013. US Central Atlantic: new plays and petroleum prospectivity. Geological Society, London, Special Publications, 369: 323–336.
- Pott, C. 2013. *Ctenophyllum* junior synonym of *Pterophyllum* (Bennettitales: Williamsoniaceae). Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 269(3): 271–283.
- Preto, N., Agnini, C., Rigo, M., Sprovieri, M. & Westphal, H. 2013. The calcareous nannofossil *Prinsiosphaera* achieved rock-forming abundances in the latest Triassic of western Tethys: consequences for the δ^{13} C of bulk carbonate. Biogeosciences, 10(9): 6053–6068.
- Preto, N., Willems, H., Guaiumi, C. & Westphal, H. 2013. Onset of significant pelagic carbonate accumulation after the Carnian Pluvial Event (CPE) in the western Tethys. Facies, 59(4): 891–914.
- Previtera, M.J., D'Angelo, J.A. & Mancuso, A.C. 2013. Preliminary chemometric study of bone diagenesis in Early Triassic cynodonts from Mendoza, Argentina. Ameghiniana, 50(4): 460–468.
- Proemse, B.C., Grasby, S.E., Wieser, M.E., Mayer, B. & Beauchamp, B. 2013. Molybdenum isotopic evidence for oxic marine conditions during the latest Permian extinction. Geology, 41(9): 967–970.
- Prokoph, A., El Bilali, H. & Ernst, R. 2013. Periodicities in the emplacement of large igneous provinces through the Phanerozoic: relations to ocean chemistry and marine biodiversity evolution. Geoscience Frontiers, 4(3): 263–276.
- Pugh, A.C., Danise, S. & Twitchett, R.J. 2013. Benthic ecosystem dynamics following the Late Triassic mass extinction event. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 83–84.
- Qin-Qin Xu, Jian-Qing Ji, Lei Zhao, Jun-Feng Gong, Jing Zhou, Guo-Qi He, Da-Li Zhong, Jin-Duo Wang & Griffiths, L. 2013. Tectonic evolution and continental crust growth of northern Xinjiang in northwestern China: remnant ocean model. Earth-Science Reviews, 126: 178–205.

Qing-guo Zhai, Bor-ming Jahn, Li Su, Jun Wang, Xuan-Xue Mo,

Hao-yang Lee, Kuo-lung Wang & Suohan Tang. 2013. Triassic arc magmatism in the Qiangtang area, northern Tibet: zircon U-Pb ages, geochemical and Sr-Nd-Hf isotopic characteristics, and tectonic implications. Journal of Asian Earth Sciences, 63: 162–178.

- QuanFeng Zheng, ChangQun Cao & MingYuan Zheng. 2013. Sedimentary features of the Permian-Triassic boundary sequence of the Meishan section in Changxing County, Zhejiang Province. SCIENCE CHINA Earth Sciences, 56(6): 956–969.
- Quiling Gao, Ning Zhang, Wenchen Xia, Qinglai Feng, Zhong-Qiang Chen, Jianping Zheng, Griffin, W.L., O'Reilly, S.Y., Pearson, N.J., Guoqing Wang, Sha Wu, Wenli Zhong & Xiufeng Sun. 2013. Origin of volcanic ash beds across the Permian-Triassic boundary, Daxiakou, South China: petrology and U-Pb age, trace elements and Hf-isotope composition of zircon. Chemical Geology, 360–361: 41–53.
- Rajabi, A., Rastad, E. & Canet, C. 2013. Metallogeny of Permian-Triassic carbonate-hosted Zn-Pb and F deposits of Iran: a review for future mineral exploration. Australian Journal of Earth Sciences, 60(2): 197–216.
- Raugust, T., Lacerda, M. & Schultz, C.L. 2013. The occurrence of *Chanaresuchus bonapartei* Romer 1971 (Archosauriformes, Proterochampsia) of the Middle Triassic of Brazil from the *Santacruzdon* Assemblage Zone, Santa Maria Formation (Paraná Basin). Geological Society, London, Special Publications, 379: 303–318.
- Reichgelt, T., Parker, W.G., Martz, J.W., Conran, J.G., van Konijnenburg-van Cittert, J.H.A. & Kürschner, W.M. 2013.
 The palynology of the Sonsela member (Late Triassic) Norian) at Petrified Forest National Park, Arizona, USA. Review of Palaeobotany and Palynology, 189: 18–28.
- Renesto, S. & Pieroni, V. 2013. Middle Triassic vertebrate remains from Rasa village (Varse, northern Italy). *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 485-488.
- Retallack, G.J. 2013. Permian and Triassic greenhouse crises. Gondwana Research, 24(1): 90–103.
- Richoz, S., Baud, A., Beauchamp, B., Grasby, S.E., Henderson, C. & Krystyn, L. 2013. Khuff margin: slope to oceanic deposits (Permian-Triassic allochthonous and exotics, Oman). *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 55–76.
- Ridd, M.F. 2013. A Middle Permian-Middle Triassic accretionary complex and a Late Triassic foredeep basin: forerunners of an Indosinian (Late Triassic) thrust complex in the Thailand-Malaysia border area. Journal of Asian Earth Sciences, 76: 99–114.
- Ridd, M.F. & Watkinson, I. 2013. The Phuket-Slate Belt terrane: tectonic evolution and strike-slip emplacement of a major terrane on the Sundaland margin of Thailand and Myanmar. Proceedings of the Geologists' Association, 124(6): 994–1010.
- Riding, J.B. 2013. The literature on Triassic, Jurassic and earliest Cretaceous dinoflagellate cysts: Supplement 1. Palynology, 37(2): 345–354.

- Riel, N., Guillot, S., Jaillard, E., Martelat, J.-E., Paquette, J.-L., Schwartz, S., Goncalves, P., Duclaux, G., Thebaud, N., Lanari, P., Janots, E. & Yuquilema, J. 2013. Metamorphic and geochronological study of the Triassic El Oro metamorphic complex, Ecuador: implications for high-temperature metamorphism in a forearc zone. Lithos, 156–159: 41–68.
- Rigaud, S., Blau, J., Martini, R. & Rettori, R. 2013. Taxonomy and phylogeny of the Trocholinidae (Involutina). Journal of Foraminiferal Research, 43(4): 317–339.
- Rigaud, S., Martini, R. & Rettori, R. 2013. A new genus of Norian involutinid foraminifer: its morphological, biostratigraphic and evolutionary significance. Acta Paleontologica Polonica, 58(2): 391–405.
- Rigo, M., McRoberts, C., Ciarapica, G., Giordana, N. & Bachan,
 A. 2013. New stratigraphic, paleontologic and geochemical data around the Triassic-Jurassic boundary at Portovenere (Northern Apennines, Italy). *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 489–499.
- Rinehart, L.F. & Lucas, S.G. 2013. A population of *Antediplodon dockumensis* (Bivalvia: Unionoida: Hyriidae) from the Revueltian (Upper Triassic: early-mid Norian) Bull Canyon Formation of West Texas: growth, allometry, productivity and ecology. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 500–523.
- Rinehart, L.F. & Lucas, S.G. 2013. The functional morphology of dermal bone ornamentation in temnospondyl amphibians. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 524–532.
- Rinehart, L.F. & Lucas, S.G. 2013. Tooth form and function in temnospondyl amphibians: relationship of shape to applied stress. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 533–542.
- Rippington, S., Cunningham, D., England, R. & Hendriks, B. 2013. The crustal assembly of southern Mongolia: New structural, lithological and geochronological data from the Nemegt and Altan ranges. Gondwana Research, 23(4): 1535–1553.
- Robertson, A.H.F., Parlak, O., Metin, Y., Vergili, Ö., Tasli, K., İnam, N. & Soycan, H. 2013. Late Palaeozoic-Cenozoic tectonic development of carbonate platform, margin and oceanic units in the Eastern Taurides, Turkey. Geological Society, London, Special Publications, 372: 167–218.
- Robertson, A.H.F., Parlak, O. & Ünlügens, U. (eds.). 2013 Geological Development of Anatolia and the Easternmost Mediterranean Region. Geological Society, London, Special Publications, 372: vi+649 pp.
- Robertson, A.H.F., Parlak, O. & Ustaömer, T. 2013. Late Palaeozoic-Early Cenozoic tectonic development of Southern Turkey and the easternmost Mediterranean region: evidence

from the inter-relations of continental and oceanic units. Geological Society, London, Special Publications, 372: 9–48.

- Robertson, A.H.F., Trivić, B., Đerić, N. & Bucur, I.I. 2013. Tectonic development of the Vardar Ocean and its margins: evidence from the Republic of Macedonia and Greek Macedonia. Tectonophysics, 595–596: 25–54.
- Rolland, Y., Alexeiev, D.V., Kröner, A., Corsini, M., Loury, C. & Monié, P. 2013. Late Palaeozoic to Mesozoic kinematic history of the Talas-Ferghana strike-slip fault (Kyrgyz West Tianshan) as revealed by ⁴⁰Ar/³⁹Ar dating of syn-kinematic white mica. Journal of Asian Earth Sciences, 67–68: 76–92.
- Romano, C., Goudemand, N., Vennemann, T.W., Ware, D., Schneebeli-Hermann, E., Hochuli, P.A., Brühwiler, T., Brinkmann, W. & Bucher, H. 2013. Climatic and biotic upheavals following the end-Permian mass extinction. Nature Geoscience, 6(1): 57–60.
- Romano, C. & Koot, M. 2013. Permian-Triassic fishes: extinction and recovery. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 53–54.
- Roniewicz, E. & Stanley, G.D. Jr. 2013. Upper Triassic corals from Nevada, western North America, and the implications for paleoecology and paleogeography. Journal of Paleontology, 87(5): 934–964.
- Rossland, A., Escalona, A. & Rolfsen, R. 2013. Permian-Holocene tectonostratigraphic evolution of the Mandal High, Central Graben, North Sea. AAPG Bulletin, 97(6): 928–957.
- Rožič, B., Gale, L. & Kolar-Jurkovšek, T. 2013. Extent of the Upper Norian-Rhaetian Slatnik Formation in the Tolmin Nappe, eastern Southern Alps. Geologija, 56(2): 175–186.
- Rubin, H., Rubin, K., Witkowski, A.J. & Wróbel, J. 2013. Assessments of the impact of zinc smelter "Miasteczko Śląskie" on groundwater quality of the Triassic carbonate aquifer within Lubliniec – Myszków major groundwater basin. Biuletyn Pánstowowego Instytutu Geologicznego, 456: 525–532.
- Ruta, M., Angielczyk, K.D., Fröbisch, J. & Benton, M.J. 2013. Decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids. Proceedings of the Royal Society, B. 280(1768) 20131971; doi: 10.1098/rspb.2013.1971 (9 pp).
- Ruta, M., Angielczyk, K.D., Fröbisch, J. & Benton, M.J. 2013. Correction: decoupling of morphological disparity and taxic diversity during the adaptive radiation of anomodont therapsids. Proceedings of the Royal Society, B. 280(1768) 20132414; doi: 10.1098/rspb.2013.2414 (2 pp).
- Ruta, M., Botha-Brink, J., Mitchell, S.A. & Benton, M.J. 2013. The radiation of cynodonts and the ground plan of mammalian morphological diversity. Proceedings of the Royal Society, B. 280(1769) 20131865; doi: 10.1098/ rspb.2013.1865 (10 pp).
- Rychliński, T., Ivanova, D.K., Jaglarz, P. & Bucur, I.I. 2013. Benthic foraminifera and calcareous algae from the Anisian-Norian succession in the Tatras (Poland, Slovakia): new data from High-Tatric and Križna units. Studia Universitatis Babeş-Bolyai, Geologia, 58(1), 21–43.
- Sadlok, G. & Wawrzyniak, Z. 2013. Upper Triassic vertebrate tracks from Kraków-Częstochowa Upland, southern Poland.

Annales Societatis Geologorum Poloniae, 83(2): 105–111.

- Saito, R., Kaiho, K., Oba, M., Takahashi, S., Zhong-Qiang Chen & Jinnan Tong. 2013. A terrestrial vegetation turnover in the middle of the Early Triassic. Global and Planetary Change, 105: 152–159.
- Saito, R., Oba, M., Kaiho, K., Maruo, C., Fujibayashi, M., Jing Chen, Zhong-Qiang Chen & Jinnan Tong. 2013. Ether lipids from the Lower and Middle Triassic at Qingyan, Guizhou Province, southern China. Organic Geochemistry, 58: 27–42.
- Saller, F., Renesto, S. & Dalla Vecchia, F.M. 2013. First record of *Langobardisaurus* (Diapsida, Protorosauria) from the Norian (Late Triassic) of Austria, and a revision of the genus. Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen, 268(1): 83–95.
- Sal'nikova, E.B., Kotov, A.B., Kovach, V.P., Velikoslavinskii, S.D., Jahn, B.-M., Sorokin, A.A., Sorokin, A.P., Wang, K.-L., Chan, S.-L., Li, H.-Ya, & Tolmacheva, E.V. 2013. Mesozoic age of the Uril Formation of the Amur Group, Lesser Khingan Terrane of the Central Asian Foldbelt: results of U-Pb and Lu-Hf isotopic studies of detrital zircons. Doklady Earth Sciences, 453(2): 1181–1184.
- Saltzman, M.R. & Sedlacek, A.R.C. 2013. Chemostratigraphy indicates a relatively complete Late Permian to Early Triassic sequence in the western United States. Geology, 41(4): 399–402.
- Sánchez-Beristain, F., López-Esquival Kranksith, L., García-Barrera, P. & Reitner, J. 2013. El primer registro de *Koskinobullina socialis* (Foraminifera) para el Triásico de Europa y sus implicaciones paleoecológicas. Boletin Geológica y Minero, 124: 437–450.
- Sanghoon Kwon, Sung Won Kim & Santosh, M. 2013. Multiple generations of mafic-ultramafic rocks from the Hongseong suture zone, western South Korea: implications for the geodynamic evolution of NE Asia. Lithos, 160–161: 68–83.
- Sanz, T., Lago, M., Galé, C., Ubide, T., Pocovi, A. & Gil, A. 2013. Magmatic rocks in the Tramuntana Chain (N of Mallorca, Spain): stratigraphic context. Geogaceta, 54: 39–42.
- Sanz, T., Lago, M., Gil, A., Galé, C., Ramajo, J., Ubide, T., Pocovi, A., Tierz, P. & Larrea, P. 2013. The Upper Triassic alkaline magmatism in the NW Iberian Chain (Spain). Journal of Iberian Geology, 39(2): 203–222.
- Scheib, C., Appleton, J.D., Miles, J.C.H. & Hodgkinson, E. 2013. Geological controls on radon potential in England. Proceedings of the Geologists' Association, 124(6): 910–928.
- Schintgen, T. & Förster, A. 2013. Geology and basin structure of the Trier–Luxembourg Basin – implications for the existence of a buried Rotliegend Graben. Zeitschrift der Deutschen Gesellschaft für Geowissenschaften, 164(4): 615–637.
- Schneebeli-Hermann, E., Kürschner, W.M., Hochuli, P.A., Ware, D., Weissert, H., Bernasconi, S.M., Roohi, G., ur-Rehman, K., Goudemand, N. & Bucher, H. 2013. Evidence for atmospheric carbon injection during the end-Permian extinction. Geology, 41(5): 579–582.
- Schoch, R.R. 2013. How body size and development biased the direction of evolution in early amphibians. Historical Biology, 25(2): 155–165.

Schoch, R.R. 2013. The evolution of major temnospondyl clades:

An inclusive phylogenetic analysis. Journal of Systematic Palaeontology, 11(6): 673–705.

- Schoch, R.R. 2013. The evolution of life cycles in temnospondyls. Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 20.
- Schoepfer, S.D., Henderson, C.M., Garrison, G.H., Foriel, J., Ward, P.D., Selby, D., Hower, J.C., Algeo, T.J. & Yanan Shen. 2013. Termination of a continent-margin upwelling system at the Permian-Triassic boundary (Opal Creek, Alberta, Canada). Global and Planetary Change, 105: 21–35.
- Schorn, A., Neubauer, F., Genser, J. & Bernroider, M. 2013. The Haselgebirge evaporitic mélange in central Northern Calcareous Alps (Austria): part of the Permian to Lower Triassic rift of the Meliata ocean? Tectonophysics, 583: 28–48.
- Schrag, D.P., Higgins, J.A., Macdonald, F.A. & Johnston, D.T. 2013. Authigenic carbonate and the history of the global carbon cycle. Science, 339(6119): 540–543.
- Schuh, M. 2013. Berichte 2012 über geologische Aufnahmen im mittleren und hinteren Kaisertal (Kaisergebirge) auf Blatt 3213 Kufstein. Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1-4): 430–432.
- Schuster, R. (ed.). 2013. 11th Workshop on Alpine Geological Studies and 7th European Symposium on Fossil Algae, Schladming, Austria, September 2013. Abstracts and Field Guides. Berichte der Geologischen Bundesanstalt, Wien, 99: 286 pp.
- Sebe, O., Crasquin, S. & Gradinaru, E. 2013. Early and Middle Anisian (Triassic) deep-water ostracods (Crustacea) from North Dobrogea (Romania). Revue de Paléobiologie, 32(2): 509–529.
- Seidel, G. 2013. Der Einfluss der Fazies des Zechsteins und der Trias auf den Bau der Thüringer Mülde. Semana (Veröffentlichungen des Naturhistorischen Museums Schleusingen), 28: 69–80.
- Senhu Lin, Xuanjun Yuan, Shizhen Tao & Zhi Yang. 2013. Geochemical characteristics of the source rocks in Mesozoic Yanchang formation, central Ordos Basin. Journal of Earth Science, 24(5): 804–814.
- Senobari-Daryan, B. 2013. *Tubiphytes* Maslov, 1956 and description of similar organisms from Triassic reefs of the Tethys. Facies, 59(1): 75–112.
- Senobari-Daryan, B. & Amirhassankhani, F. 2013. Chambered hexactinellid sponges from Upper Triassic (Norian-Rhaetian) reefs of Nayband Formation in central Iran. Rivista Italiana di Paleontologia e Stratigrafia, 118(2): 247–260.
- Senobari-Daryan, B. & Amirhassankhani, F. 2013. Lovcenipora iranica nov. sp., an unusually large chaetetid sponge from the Upper Triassic (Howz-e Khan member, Nayband Formation) of north east Iran. Zitteliana, A53: 15–22.
- Seward, L.J., Stallebrass, S.E. & Skipper, J. 2013. Remoulding of the Mercia Mudstone Group around CFA pile shafts. Quarterly Journal of Engineering Geology and Hydrogeology, 46(1): 41–51.
- Seyfullah, L.J., Kustatscher, E. & Taylor, W.A. 2013. The first discovery of in situ *Verrucosisporites applanatus* spores from the Middle Triassic flora from Bromsgrove (Worcestershire, UK). Review of Palaeobotany and Palynology, 197: 15–25.

- Shan Li, Tao Wang, Wilde, S.A. & Ying Tong. 2013. Evolution, source and tectonic significance of Early Mesozoic granitoid magmatism in the Central Asian Orogenic Belt (central segment). Earth-Science Reviews, 126: 206–234.
- Shang Qing-hua & Li Chun. 2013. The sexual dimorphism of *Shastasaurus tangae* (Reptilia: Ichthyosauria) from the Triassic Guanling Biota, China. Vertebrata Palasiatica, 51(4): 253–264.
- ShaoFeng Liu & GuoWei Zhang. 2013. Mesozoic basin development and its indication of collisional orogeny in the Dabei orogen. Chinese Science Bulletin, 58(8): 827–852.
- Shaofeng Liu, San Su & Guowei Zhang. 2013. Early Mesozoic basin development in North China: indications of cratonic deformation. Journal of Asian Earth Sciences, 62: 221–236.
- Shen Liu, Ruizhong Hu, Shan Gao, Caixia Feng, Coulson, I.M., Guangying Feng, Youqiang Qi, Yuhong Yang, Chaogui Yang & Liang Tang. 2013. Zircon U–Pb age and Sr–Nd–Hf isotopic constraints on the age and origin of Triassic mafic dikes, Dalian area, Northeast China. International Geology Review, 55(2): 249–262.
- Shi-Yong Liao, Fu-Guang Yin, Zhi-Ming Sun, Dong-Bing Wang, Yuan Tang & Jie Sun. 2013. Early Middle Triassic mafic dikes from the Baoshan subterrane, western Yunnan: implications for the tectonic evolution of the Palaeotethys in southeast Asia. International Geology Review, 55(6): 976–993.
- Shooshtarizadeh, P. & Yazdi, M. 2013. Conodont biostratigraphy of Permian-Triassic boundary deposits in Koh-e-Chah Kaland, north of Abadeh, Iran. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 543–555.
- Shu-zhong Shen, Chang-qun Cao, Hua Zhang, Bowring, S.A., Henderson, C.M., Payne, J.L., Davydov, V.I., Bo Chen, Dong-xun Yuan, Yi-chun Zhang, Wei Wang & Quan-feng Zheng. 2013. High-resolution $\delta^{13}C_{carb}$ chemostratigraphy from latest Guadalupian through earliest Triassic in South China and Iran. Earth and Planetary Science Letters, 375: 156–165.
- Shugen Liu, Bin Deng, Jansa, B., Guozhi Wang, Xianghui Li, Chengshan Wang, Shun Liu, Zhiwu Li, Juchu Li, Wei Sun & Bo Ran. 2013. Late Triassic thickening of the Songpan-Ganzi Triassic flysch at the edge of the northeastern Tibetan Plateau. International Geology Review, 55(16): 2008–2015.
- Simpson, E.L., Hartline, B.W., Szajna, M.J., Fillmore, D.L. & Heness, E.A. 2013. An ensiferan (cricket) fossil from the Late Triassic Lockatong Formation, Newark Basin, Pennsylvania. *In*, Tanner, L.H., Speilmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 556–561.
- Singh, P., Hussain, R., Al-Zuabi, Y., Al-Khaled, O., Rahaman, M., Hameed, M., Ebrahim, M., Haffez, M., Al-Rashid, T., Al-Ghareeb, S., Al-Kandary, A. & Al-Fares, A. 2013. Basement configuration and its impact on Permo-Triassic prospectivity in Kuwait. *In*, Pöppelreiter, M.C. (ed.), Permo-Triassic Sequences of the Arabian Plate. EAGE Publications bv.: 43–54.

- Sinitshenkova, N.D. 2013. New mayflies (Insecta: *Ephemerida* = *Ephemeroptera*) from the intertrappean deposits of the Tunguska Basin, Siberia. Paleontological Journal, 47(1): 84–88.
- Skawina, A. 2013. Population dynamics and taphonomy of the Late Triassic (Carnian) freshwater bivalves from Krasiejón (Poland). Palaeogeography, Palaeoclimatology, Palaeoecology, 379-380: 68–80.
- Small, B.J. & Martz, J.W. 2013. A new aetosaur from the upper Triassic Chinle Formation of the Eagle Basin, Colorado, USA. Geological Society, London, Special Publications, 379: 393–412.
- Smyshlyaeva, O.P. & Zakharov, Y.D. 2013. New members of the Family Flemingitidae (Ammonoidea) from the Lower Triassic of Primorye. Paleontological Journal, 47(3): 247–255.
- Sobolev, N.V., Logvinova, A.M., Nikolenko, E.I. & Lobanov, S.S. 2013. Mineralogical criteria for the diamond potential of Upper Triassic placers on the northeastern margin of the Siberian Platform. Russian Geology and Geophysics, 54(8): 903–916.
- Sookias, R.B. & Butler, R.J. 2013. Euparkeriidae. Geological Society, London, Special Publications, 379: 35–48.
- Sookias, R.B., Senniikov, A.G., Butler, R.J. & Gower, D.J. 2013. Redescription and phylogenetic position of the archosauriform *Dorosuchus neoetus* from the Middle Triassic of Russia. Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 88.
- Sorokin, A.A. & Kudryashov, N.M. 2013. Early Mesozoic magmatism of the Bureinskii Terrane of the Central Asian Foldbelt: age and geodynamic setting. Doklady Earth Sciences, 452(1): 915–921.
- Sowizdzal, A., Papiernik, B., Machowski, G. & Hajto, M. 2013. Petrophysical parameters of the Lower Triassic deposits in prospective location for Enhanced Geothermal System (central Poland). Geological Quarterly, 57(4): 729–744.
- Spielmann, J.A., Lucas, S.G., Heckert, A.B. & Krzyanowski, S.E. 2013. *Trilophosaurus* (Archosauromorpha: Trilophosauridae) postcrania from the Upper Triassic Blue Mesa Member of the Petrified Forest Formation (Carnian: Adananian), Arizona, USA. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 567–572.
- Spielmann, J.A., Lucas, S.G. & Huber, P. 2013. A new record of *Spinosuchus caseanus* (Archosauromorpha: Trilophosauridae) from the Late Triassic (Adamanian) of West Texas. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 573–576.
- Spielmann, J.A., Lucas, S.G. & Hunt, A.P. 2013. The first Norian (Revueltian) rhynchosaur: Bull Canyon Formation, New Mexico, U.S.A. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 562–566.
- Stampfli, G.M., Hochard, C., Vérard, C., Wilhem, C. & von

Raumer, J. 2013. The formation of Pangea. Tectonophysics, 593: 1-19.

- Stanley, G.D. Jr., Yancey, T.E. & Shepherd, H.M.E. 2013. Giant Upper Triassic bivalves of Wrangellia, Vancouver Island, Canada. Canadian Journal of Earth Sciences, 50(2): 142–147.
- Stockar, R., Adatte, T., Baumgartner, P.O. & Föllmi, K.B. 2013. Palaeoenvironmental significance of organic facies and stable isotope signatures: the Ladinian San Giorgio Dolomite and Meride Limestone of Monte San Giorgio (Switzerland, WHL UNESCO). Sedimentology, 60(1): 239–269.
- Stockar, R. & Garassimo, A. 2013. *Meridecaris ladinica*, n.gen. n.sp. (Crustacea, Decapoda, Clytiopsida) from the Middle Triassic (Ladinian) of Monte San Giorgio (Canton Ticino, Switzerland). Neues Jahrbuch für Geologie und Paläontologie Abhandlungen, 270(3): 347–356.
- Stocker, M.R. & Butler, R.J. 2013. Phytosauria. Geological Society, London, Special Publications, 379: 91–117.
- Stubbs, T. & Benton, M.J. 2013. The adaptive radiation of Triassic marine reptiles. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 89.
- Stubbs, T.L., Pierce, S.E., Rayfield, E.J. & Anderson, P.S.L. 2013. Morphological and biomechanical disparity of crocodile-line archosaurs following the end-Triassic extinction. Proceedings of the Royal Society, B. 280(1770) 20131940; doi: 10.1098/ rspb.2013.1940 (9 pp).
- Sudar, M.N., Gawlick, H.-J., Lein, R., Missoni, S., Kovács, S. & Jovanović, D. 2013. Depositional environment, age and facies of the Middle Triassic Bulog and Rid formations in the Inner Dinarides (Zlatibor Mountain, SW Serbia): evidence for the Anisian break-up of the Neotethys Ocean. Neues Jahrbuch für Geologie und Paläontologie - Abhandlungen, 269(3): 291–320.
- Sues, H.-D., Desojo, J.B. & Ezcurra, M.D. 2013. Doswellidae: a clade of unusual armoured archosauriformes from the Lower and Middle Triassic. Geological Society, London, Special Publications, 379: 49–58.
- Sues, H.-D., Hook, R.W. & Olsen, P.E. 2013. Donald Baird and his discoveries of Carboniferous and early Mesozoic vertebrates in Nova Scotia. Atlantic Geology, 49: 90–103.
- Sues, H.-D. & Schoch, R.R. 2013. First record of *Colognathus* (?Amniota) from the Middle Triassic of Europe. Journal of Vertebrate Paleontology, 33(4): 998–1002.
- Sues, H.-D. & Schoch, R.R. 2013. Anatomy and phylogenetic relationships of *Calamops paludosus* (Temnospondyli, Stereospondyli) from the Triassic of the Newark Basin, Pennsylvania. Journal of Vertebrate Paleontology, 33(5): 1061–1070.
- Sues, H.-D. & Schoch, R.R. 2013. Reassessment of cf. *Halticosaurus orbitoangulatus* from the Upper Triassic (Norian) of Germany – a pseudosuchian, not a dinosaur. Zoological Journal of the Linnean Society, 168(4): 859–872.
- Sulej, T. & Niedźwiedski, G. 2013. A new large temnospondyl amphibian from the Early Triassic of Poland. Acta Palaeontologica Polonica, 58(1): 65–75.
- Sun, Z., Tintori, A., Jiang, D. & Motani, R. 2013. A new perleidid from the Spathian (Olenekian, Early Triassic)

of Chaohu, Anhui Province, China. Rivista Italiana di Paleontologia e Stratigrafia, 119(3): 275–285.

- Sung Kwun Chough, Dong Jin Lee & Jin-Han Ree. 2013. Whereabouts of the collision belt between the Sino-Korean and South China blocks in the northeast Asian margin. Geosciences Journal, 17(4): 397–401.
- Szurlies, M. 2013. Late Permian (Zechstein) magnetostratigraphy in Western and Central Europe. Geological Society, London, Special Publications, 376: 73–85.
- Taborda, J.R.A., Cerda, I.A. & Desojo, J.B. 2013. Growth curve of *Aetosauroides scagliai* Casamiguela 1960 (Pseudosuchia: Aetosauria) inferred from osteoderm histology. Geological Society, London, Special Publications, 379: 413–440.
- Takahashi, S., Kaiho, K., Hori, R.S., Gorjan, P., Watanabe, T., Yamakita, S., Aita, Y., Takemura, A., Spörli, K.B., Kakegawa, T. & Oba, M. 2013. Sulfur isotope profiles in the pelagic Panthalassic deep sea during the Permian-Triassic transition. Global and Planetary Change, 105: 68–78.
- Tan Kai & Jin Fan. 2013. Re-study of *Gymnoichthys inopinatus* from Middle Triassic of Luoping, Yunnan, China. Vertebrata Palasiatica, 51(1): 1–16.
- Tanner, L.H. 2013. The enigmatic Quaco cobbles, Upper Triassic, Canadian Maritimes: deformation by tectonics or seismic shock? *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 577–581.
- Tanner, L.H. & Lucas, S.G. 2013. Degraded wood in the Upper Triassic Petrified Forest Formation (Chinle Group), northern Arizona: differentiating fungal rot from arthropod boring. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 582–588.
- Tanner, L.H. & Lucas, S.G. 2013. Isotopic composition of carbonate facies in formations of the Upper Triassic Chinle Group, Four Corners area, southwestern United States. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 589–596.
- Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.). 2013. The Triassic System: new developments in stratigraphy and paleontology. New Mexico Museum of Natural History and Science Bulletin, 61: iv+612 pp.
- Tao Chongzhi, Bai Guoping, Liu Junlan, Deng Chao, Lu Xiaoxin, Liu Houwu & Wang Dapeng. 2013. Palaeogeography and mineral resources, Mesozoic lithofacies, palaeogeography and petroleum prospectivity in North Carnarvon Basin, Australia. Journal of Palaeogeography, 2(1): 66–80.
- Tao Yang, Laimin Zhu, Fei Wang, Hujun Gong & Rukui Lu. 2013. Geochemistry, petrogenesis and tectonic implications of granitic plutons at the Lizizuan orogenic goldfield in the Western Qinlong Orogen, central China. Geological Magazine, 150(1): 50–71.
- Tari, G. & Jabour, H. 2013. Salt tectonics along the Atlantic margin of Morocco. Geological Society, London, Special

Publications, 369: 337-353.

- Tassi, L.V., Monti, M., Gallego, O.F., Zavattieri, A.M. & Lara, M.B. 2013. The first spinicaudatan (Crustacea: Diplostraca) from Permo-Triassic continental sequences of South America and its palaeoecological context. Alcheringa, 37(2): 189–201.
- Tatzreiter, F. & Zorn, I. 2013. The cephalopods of the Hallstatt Limestone described in the monographs by Edmund von Mojsisovics 1873, 1875 and 1902 stored in the Palaeontological Collection of the Geological Survey of Austria. Jahrbuch der Geologischen Bundesanstalt, Wien, 153(1–4): 107–190.
- Thuy, B., Ishida, Y., Doi, E. & Kroh, A. 2013. New ophiacanthid brittle stars (Echinodermata: Ophiuroidea) from the Upper Triassic of Japan: first insights into the origin and evolution of an extant deep-sea group. Journal of Systematic Palaeontology, 11(5): 515–530.
- Tintori, A. 2013. A new species of *Saurichthys* (Actinopterygii) from the Middle Triassic (Early Ladinian) of the Northern Grigna Mountain (Lombardy, Italy). Rivista Italiana di Paleontologia e Stratigrafia, 119(3): 287–302.
- Tohver, E., Cawood, P.A., Riccomini, C., Lana, C. & Trindade, R.I.F. 2013. Shaking a methane fizz: seismicity from the Araguainha impact event and the Permian-Triassic global carbon isotope record. Palaeogeography, Palaeoclimatology, Palaeoecology, 387: 66–75.
- Toljagić, O. & Butler, R.J. 2013. Triassic-Jurassic mass extinction as trigger for the Mesozoic radiation of crocodylomorphs. Biology Letters, 9(3), 20130095, doi: 10.1098/rsbl.2013.0095 (4 pp).
- Torsvik, T.H. & Cocks, L.R.M. 2013. Gondwana from top to base in space and time. Gondwana Research, 24(3-4): 999–1030.
- Touping Peng, Wilde, S.A., Yuejun Wang, Weiming Fan & Bingxia Peng. 2013. Mid-Triassic felsic igneous rocks from the southern Lancangjiang Zone, SW China: petrogenesis and implications for the evolution of Paleo-Tethys. Lithos, 168-169: 15–32.
- Tremblay, A., Roden-Tice, M.K., Brandt, J.A. & Megan, T.W. 2013. Mesozoic fault reactivation along the St. Lawrence rift system, eastern Canada: thermochronologic evidence from apatite fission-track dating. Geological Society of America Bulletin, 125(5/6): 794–810.
- Trendell, A.M., Nordt, L.C., Atchley, S.C., Leblanc, S.L. & Dworkin, S.I. 2013. Determining floodplain plant distributions and populations using paleopedology and fossil root traces: Upper Triassic Sonsela Member of the Chinle Formation at Petrified Forest National Park, Arizona. Palaios, 28(7): 471–490.
- Trippetta, F., Collettini, C., Meredith, P.G. & Vinciguerra, S. 2013. Evolution of the elastic moduli of seismogenic Triassic evaporites subjected to cyclic stressing. Tectonophysics, 592: 67–79.
- Trotteyn, M.J., Arcucci, A.B. & Raugust, T. 2013. Proterochampsia: an endemic archosauriform clade from South America. Geological Society, London, Special Publications, 379: 59–90.
- Trotteyn, M.J. & Martinez, R.N. 2013. First record of

coxofemoral dysplasia in a Late Triassic rhynchosaur. Ameghiniana, 50(2): 217–226.

- Tsuji, L.A., Sobral, G. & Müller, J. 2013. *Ruhuhuaria reiszi*, a new procolophonoid reptile from the Triassic Ruhuhu Basin of Tanzania. Comptes Rendus Palevol, 12(7-8): 487–494.
- Turner, A.H. & Nesbitt, S.J. 2013. Body size evolution during the Triassic archosauriform radiation. Geological Society, London, Special Publications, 379: 573-597.
- Vajda, V., Calner, M. & Ahlberg, A. 2013. Palynostratigraphy of dinosaur footprint-bearing deposits from the Triassic-Jurassic boundary interval of Sweden. GFF, 135(1): 120-130.
- Van de Schootbrugge, B., Bachan, A., Suan, G., Richoz, S. & Payne, J.L. 2013. Microbes, mud and methane: cause and consequence of recurrent Early Jurassic anoxia following the end-Triassic mass extinction. Palaeontology, 56(4): 685–709.
- Veevers, J.J. 2013. Pangea: geochronological correlation of successive environments and strati-tectonic phases in Europe and Australia. Earth-Science Reviews, 127: 48–95.
- Vérard, C. & Stampfli, G.M. 2013. Geodynamic reconstructions of the Australides – 2: Mesozoic-Cainozoic. Geosciences 3(2): 331–353.
- Viglietti, P.A., Smith, R.M.H. & Compton, J.S. 2013. Origin and palaeoenvironmental significance of *Lystrosaurus* bonebeds in the earliest Triassic Karoo Basin, South Africa. Palaeogeography, Palaeoclimatology, Palaeoecology, 392: 9–21.
- Vilhena, D.A. & Smith, A.B. 2013. Spatial bias in the marine fossil record. PLoS ONE, 8(10): e74470. doi:10.1371/ journal.pone.0074470.
- Vitte, L.V. & Vasilevskiy, A.N. 2013. Geologic nature of regional magnetic and gravity anomalies in the Mongol-Transbaikalian province of the Central Asian Fold Belt. Russian Geology and Geophysics, 54(12): 1442–1449.
- Volkov, A.N. 2013. New species of Triaplidae from the Babii Kamen' locality (Kuznetsk Basin). Paleontological Journal, 47(1): 94–97.
- Walther, M. & Fröbisch, J. 2013. The quality of the fossil record of anomodonts (Synapsida, Therapsida). Comptes Rendus Palevol, 12(7-8): 495–504.
- Wang Feng Qin, Wang Xiang Zeng, Zhang Li Xia, Liu Hong Jun & Lei Yu. 2013. The shale gas resources estimation: an example from Mesozoic Triassic Yanchang Formation Member Chang 7, Ordos Basin. Earth Science Frontiers, 20(3): 240–246.
- Wang, H., Rahn, M., Zhou, J. & Tao, X. 2013. Tectonothermal evolution of the Triassic flysch in the Songpan-Garzê orogen, eastern Tibetan plateau. Tectonophysics, 608: 505–516.
- Wang Jianpo, Cheng Long & Zeng Xiongwei. 2013. Microfacies and depositional environment evolution of the Middle-Upper Triassic Zhuganpo Formation in Zhenfeng, southwestern Guizhou. Geological Journal of China Universities, 19(3): 513–519.
- Wang Runfu, Xu Shichao, Wu Xiaochun, Li Chun & Wang Suozhu. 2013. A new specimen of *Shansisuchus shansisuchus* Young, 1964 (Diapsida, Archosauriformes) from the Triassic of Shanxi, China. Acta Geologica Sinica, 87(5): 1185–1197.
- Wang Zecheng, Huang Shipeng, Gong Deyu, Wu Wei & Yu Chong. 2013. Geochemical characteristics of natural gases

in the Upper Triassic Xujiahe Formation in the southern Sichuan Basin, SW China. International Journal of Coal Geology, 120: 15–23.

- Ware, D. & Bucher, H. 2013. High resolution biostratigraphy and biodiversity dynamics of Dienerian (Early Triassic) ammonoids from the Northern Indian Margin. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 59–60.
- Warrington, G. 2013. Tale of a boundary (Abstract). Geoscience in South-West England, 13(2): 247.
- Warrington, G. 2013. Inside a Mid Triassic fluvial deposit in Cheshire (Abstract). Geoscience in South-West England, 13(2): 247.
- Waters, C.N., Bridge, D.M., Humpage, A.J., Barclay, W.J. & Smith, N.J.P. 2013. Geology of the Dudley district– a brief explanation of the geological map. Sheet Explanation of the British Geological Survey, 1:50 000 Sheet 167 Dudley (England and Wales). Keyworth, Nottingham: British Geological Survey, ii+46 pp.
- Wawrzyniak, Z., Filipiak, P. & Kenrick, P. 2013. New Triassic floras from Southern Poland. The Palaeontological Association Newsletter, 84; Annual Meeting Programme and Abstracts: 93.
- Wei Shi, Shu-Wen Dong, Ratschbacher, L., Mi Tian, Jian-Hua Li & Guo-Li Wu. 2013. Meso-Cenozoic tectonic evolution of the Dangyang Basin, north-central Yangtze craton, central China. International Geology Review, 55(3): 382–396.
- Wei Yang, Jolivet, M., Dupont-Nivet, G., Zhaojie Guo, Zhicheng Zhang & Chaodong Wu. 2013. Source to sink relations between the Tian Shan and Junggar Basin (northwest China) from Late Palaeozoic to Quaternary: evidence from detrital U-Pb zircon geochronology. Basin Research, 25(2): 219–240.
- Wei-Jen Chen, Lavoué, S. & Mayden, R.L. 2013. Evolutionary origin and early biogeography of otophysan fishes (Ostariophysi: Teleostei). Evolution, 67(8): 2218–2239.
- Weinbaum, J.C. 2013. Posrcranial skeleton of *Postosuchus kirkpatricki* (Archosauria: Paracrocodylomorpha), from the Upper Triassic of the United States. Geological Society, London, Special Publications, 379: 525–553.
- Weissert, H. 2013. C-isotope geochemistry tool for chemostratigraphy and carbon-cycle history. Ciências da Terra, 18: 19–22.
- Wen Wen, Qi-yue Zhang, Shi-xue Hu, Benton, M.J., Changyong Zhou, Xie Tao, Jin-yuan Huang & Zhong-qiang Chen. 2013. Coelacanths from the Middle Triassic Luoping biota, Yunnan, South China, with the earliest evidence of ovoviviparity. Acta Palaeontologica Polonica 58(1): 175–193.
- Wen-Liang Xu, Fu-Ping Pei, Feng Wang, En Meng, Wei-Qiang Ji, De-Bin Yang & Wei Wang. 2013. Spatial-temporal relationships of Mesozoic volcanic rocks in NE China: constraints on tectonic overprinting and transformations between multiple tectonic regimes. Journal of Asian Earth Sciences, 74: 167–193.
- Wendruff, A.J. & Wilson, M.V.H. 2013. New Early Triassic coelacanth in the family Laugiidae (Sarcopterygii: Actinistia) from the Sulphur Mountain Formation near Wapiti Lake, British Columbia, Canada. Canadian Journal of Earth

Sciences, 50(9): 904-910.

- Wenjiao Xiao, Windley, B.F., Allen, M.B. & Chunming Han. 2013. Paleozoic multiple accretionary and collisional tectonics of the Chinese Tianshan orogenic collage. Gondwana Research, 23(4): 1316–1341.
- Wilson, L.A.B., Furrer, H., Stockar, R. & Sánchez-Villagra, M.R. 2013. A quantitative evaluation of evolutionary patterns in opercle bone shape in *Saurichthys* (Actinopterygii: Saurichthyidae). Palaeontology, 56(4): 901–915.
- Winguth, A. & Winguth, C. 2013. Precession-driven monsoon variability at the Permian-Triassic boundary – implications for anoxia and the mass extinction. Global and Planetary Change, 105: 160–170.
- Withjack, M.O., Schlische, R.W., Malinconico, M.L. & Olsen, P.E. 2013. Rift-basin development: lessons from the Triassic-Jurassic Newark Basin of eastern North America. Geological Society, London, Special Publications, 369: 301–321.
- Witzmann, F. 2013. Phylogenetic patterns of character evolution in the hyobrachial apparatus of early tetrapods. Earth and Environmental Science Transactions of the Royal Society of Edinburgh, 104(2): 145–167.
- Woods, A.D. 2013. Microbial ooids and cortoids from the Lower Triassic (Spathian) Virgin Limestone, Nevada, USA: evidence for an Early Triassic microbial bloom in shallow depositional environments. Global and Planetary Change, 105: 91–101.
- Wopfner, H. 2013. Late Palaeozoic Early Triassic deposition and climates between Samfrau and Tethys: a review. Geological Society, London, Special Publications, 376: 5-32.
- Wright, N., Zahirovic, S., Müller, R.D.M. & Seton, M. 2013. Towards community-driven paleogeographic reconstructions: integrating open-access paleogeographic and paleobiology data with plate tectonics. Biogeosciences, 10(3): 1529–1541.
- Xiang Bai, Shuwen Liu, Wei Wang, Pengtao Yang & Qiugen Li. 2013. U-Pb geochronology and Lu-Hf isotopes of zircons from newly identified Permian–Early Triassic plutons in western Liaoning province along the northern margin of the North China Craton: constraints on petrogenesis and tectonic setting. International Journal of Earth Sciences, 102(3): 671–685.
- Xiangyang Xie & Heller, P.L. 2013. U-Pb detrital zircon geochronology and its implications: the early Late Triassic Yanchang Formation, south Ordos Basin, China. Journal of Asian Earth Sciences, 63: 86–98.
- Xiao-Wei Li, Xuan-Xue Mo, Xue-Hui Yu, Xiong-Fei Huang, Ping Wei & Wen-Yan He. 2013. Petrology and geochemistry of the early Mesozoic pyroxene andesites in the Maixiu area, west Qinling, China: products of subduction or syn-collision? Lithos, 172-173: 158–174.
- XiaoChun Liu, BorMing Jahn & SanZhong Li. 2013. Diachroneity of continental subduction and exhumation: constraints from the Permian-Triassic HP metamorphic terrane in the Tongbai orogen, central China. Chinese Science Bulletin, 58(35): 4397–4404.
- Xiaoxia Wang, Tao Wang & Chengli Zhang. 2013. Neoproterozoic, Paleozoic, and Mesozoic granitoid magmatism in the Qinling Orogen, China: constraints on orogenic process. Journal of Asian Earth Sciences, 72: 129–151.

- Xin Bushe, Yang Hua, Fu Jinhua, Wang Duoyun, Yao Jingli, Zhang Yu & Zuo Bo. 2013. SHRIMP U-Pb zircon age and its stratigraphic significance of tuff at the bottom of the Upper Triassic Nanying' er Group, Mount Baoji area, Jingyuan County, Gansu Province. Geological Review, 59(2): 267–273.
- Xin-chun Liu, Wei Wang, Shu-zhong Shen, Gorgij, M.N., Fa-cheng Ye, Yi-chun Zhang, Furuyama, G., Kano, A. & Xiao-zheng Chen. 2013. Late Guadalupian to Lopingian (Permian) carbon and strontium isotopic chemostratigraphy in the Abadeh section, central Iran. Gondwana Research, 24(1): 222–232.
- Xing, L., Klein, H., Lockley, M.G., Jianjun Li, Jianping Zhang, Matsukawa, M. & Jiafei Xiao. 2013. *Chirotherium* trackways from the Middle Triassic of Guizhou, China. Ichnos, 20(2): 99-107.
- Xing Li-da, Klein, H., Lockley, M. G., Wang Shi-li, Chen Wei, Ye Yong, Matsukawa, M. & Zhang Jian-ping. 2013. Earliest records of theropod and mammal-like theropod footprints in the upper Triassic of Sichuan Basin, China. Vertebrata Palasiatica, 51(3): 184–198.
- Xiucheng Tan, Qingsong Xia, Jingshan Chen, Ling Li, Hong Liu, Bing Luo, Jiwen Xia & Jiajing Yang. 2013. Basin-scale sand deposition in the Upper Triassic Xujiahe Formation of the Sichuan Basin, southwest China: sedimentary framework and conceptual model. Journal of Earth Sciences, 24(1): 89–103.
- Xuefeng Zhang, Tonglou Guo, Bo Liu, Xiaoyue Fu & Shuanglin Wu. 2013. Porosity formation and evolution of the deeply buried Lower Triassic Feixianguan Formation, Puguang gas field, NE Sichuan basin, China. Open Journal of Geology, 3(4): 300–312.
- Yang Lirong, Song Haijun, Tong Jinnan, Chu Daoliang & Tian Li. 2013. The extinction pattern of fusulinids during the Permian-Triassic crisis at the Kaijiaping section, Cili, Hunan Province. Acta Micropalaeontologica Sinica, 30(4): 353–366.
- Yanlong Chen, Twitchett, R.J., Haishui Jiang, Richoz, S., Xulong Lai, Chunbo Yan, Yadong Sun, Xiaodan Liu & Wang, L. 2013. Size variation of conodonts during the Smithian-Spathian (Early Triassic) global warming event. Geology, 41(8): 823-826.
- Yanqiu Zhang, Krause, M. & Mutti, M. 2013. The formation and structure evolution of Zechstein (Upper Permian) salt in Northeast German Basin: a review. Open Journal of Geology, 3(8): 411–426.
- Yan-hao Lin, Ze-ming Zhang, Xin Dong, Hua Xiang & Rong Yan. 2013. Early Mesozoic metamorphism and tectonic significance of the eastern segment of the Lhasa terrane, south Tibet. Journal of Asian Earth Sciences, 79: 160–183.
- Yao-Hui Jiang, Ru-Ya Jia, Zheng Liu, Shi-Yong Liao, Peng Zhao & Qing Zhou. 2013. Origin of Middle Triassic high-K calcalkaline granitoids and their potassic microgranular enclaves from the western Kunlun orogen, northwest China: a record of the closure of Paleo-Tethys. Lithos, 156–159: 13–30.
- Yapaskurt, O.V. & Shikhanov, S.E. 2013. Stages of mineral formation in early Triassic to Quaternary deposits with implications for the geodynamic evolution of the Koltogor-Urengoy trough system, West Siberian Plate. Moscow University Geology Bulletin, 68(6): 329–338.

- Yarmolyuk, V.V., Kuzmin, M.I., Vorontsov, A.A. & Khomutova, M.Yu. 2013. West Pacific-type convergent boundaries: role in the crust growth history of the central-Asian orogen. Journal of Asian Earth Sciences, 62: 67–78.
- Yarmolyuk, V.V., Kuz'min, M.I. & Vorontsov, A.A. 2013. West Pacific-type convergent boundaries and their role in the formation of the Central Asian Fold Belt. Russian Geology and Geophysics, 54(12): 1427–1441.
- Yi Duan, Yiding Yuan & Raorong Qian. 2013. Migration features of crude oil in fluvial deposits of Maling oilfield in Ordos Basin, China. Organic Geochemistry, 58: 78–85.
- Yin HongFu & Song HaiJun. 2013. Mass extinction and Pangea integration during the Paleozoic-Mesozoic transition. Science China Earth Sciences, 56(1): 1–13.
- Yong-Fei Zheng, Wen-Jiao Xiao & Guochun Zhao. 2013. Introduction to tectonics of China. Gondwana Research, 23(4): 1189–1206.
- Yousefirad, M. 2013. Using conodont elements to distinguish Permian-Triassic boundary disconformity near Haftad Gholleh, central Iran. Earth Sciences Research Journal, 17(1): 61–65.
- Yuan-Bao Wu & Yong-Fei Zhang. 2013. Tectonic evolution of a composite collision orogen: an overview on the Qinling-Tongbai-Hong'an-Dabie-Sulu orogenic belt in central China. Gondwana Research, 23(4): 1402–1428.
- Yueyun Wang, Weiming Fan, Guowei Zhang & Yanhua Zhang. 2013. Phanerozoic tectonics of the South China Block: key observations and controversies. Gondwana Research, 23(4): 1273-1305.
- Yümün, Z.Ü., Kiliç, A.M., Martini, R., Metzger, J. & Tunç, M. 2013. Late Triassic and Lower Jurassic foraminifera of the carbonate platform of the Beyaz Aladağ Group (Eastern Taurus, Turkey): new stratigraphic implications. Geobios, 46(5): 447–459.
- Yunpeng Wang, Changyi Zhao, Hongjun Wang, Zhaoyun Wang & Zecheng Wang. 2013. Origins of natural gases from marine strata in northeastern Sichuan Basin (China) from carbon molecular moieties and isotopic data. Journal of Asian Earth Sciences, 65: 13–20.
- Yunyan Ni, Jinxing Dai, Guangyu Zhu, Shuichang Zhang, Dijia Zhang, Jin Su, Xiaowan Tao, Fengrong Liao, Wei Wu, Deyu Cong & Quanyou Liu. 2013. Stable hydrogen and carbon isotopic ratios of coal-derived and oil-derived gases: a case study in the Tarim basin, NW China. International Journal of Coal Geology, 116–117: 302–313.
- Zakharov, Y.D. & Abnavi, N.M. 2013. The ammonoid recovery after the end-Permian mass extinction: evidence from the Iran-Transcaucasia area, Siberia, Primorye, and Kazakhstan. Acta Palaeontologica Polonica, 58(1): 127–147.
- Zakharov, Y.D., Bondarenko, L.G., Smyshylyaeva, O.P & Popov, A.M. 2013. Late Smithian (Early Triassic) ammonoids from the *Anasibirites nevolini* Zone of South Pimorye, Russian Far East. *In*, Tanner, L.H., Spielmann, J.A. & Lucas, S.G. (eds.), The Triassic System: New Developments in Stratigraphy and Paleontology. New Mexico Museum of Natural History and Science, Bulletin 61: 597–612.
- Zamannejad, A., Jahani, D., Lotfpour, M. & Movahed, B. 2013.

Mixed evaporite/carbonate characteristics of the Triassic Kangan Formation, offshore area, Persian Gulf. Revista Mexicana de Ciencias Geologicas, 30(3): 540–551.

- Zanchetta, S., Berra, F., Zanchi, A., Bergomi, M., Caridroit, M., Nicora, A. & Heidarzadeh, G. 2013. The record of the Late Palaeozoic active margin of the Palaeotethys in NE Iran: constraints on the Cimmerian orogeny. Gondwana Research, 24(3-4): 1237–1266.
- Zatoń, M., Taylor, P.D. & Vinn, O. 2013. Early Triassic (Spathian) post-extinction microconchids from western Pangea. Journal of Paleontology, 87(1): 159-165.
- Zeng, Y.H., Fu, X.G., Zeng, S.Q. & Du, G. 2013. Upper Triassic potential source rocks in the Qiangtang Basin, Tibet: organic geochemical characteristics. Journal of Petroleum Geology, 36(3): 237–255.
- Zhang Cai-li, Zhang Lei, Chen Tiao-sheng & Zhang Yan. 2013. Provenance and parent rock types of Member 7 of Yanchang Formation (Triassic), Ordos Basin. Acta Sedimentologica Sinica, 31(3): 430–439.
- Zhang Feng-bo, Yang Yun-xiang, Cao Yan-ping, Bai Feng, Xu Ning, Jiang Hong-yan, Wang Le & Song Fu-ming. 2013. The compatability experiment research on Chang 8 reservoir of Liuluoyu area in Xiasiwan oilfield. Northwestern Geology, 46(3): 197–202.
- Zhang Qingchun, Wang Kaiming, Luo Shunshe & Wuxiao Zhi. 2013. Study on the characteristics and origin of the oil shale in the Chang 7 Member, Yanchang Formation in Ordos Basin. Advances in Geosciences, 3(4): 197–209.
- Zhang, W., Wu, T.R., Feng, J.C., Zheng, R.G. & He, Y.K. 2013. Time constraints for the closing of the Paleo-Asian Ocean in the northern Alxa region: evidence from Wuliji granites. SCIENCE CHINA Earth Sciences, 56(1): 153–164.
- Zhao Li-Jun, Liu Jun, Li Chun & He Tao. 2013. A new thalattosaur, *Concavispina biseridens* gen. et sp. nov. from Guanling, Guizhou, China. Vertebrata Palasiatica, 51(1): 24–28.
- Zhao Yang, Ratschbacher, L., Jonckheere, R., Enkelman, E., Yunpeng Dong, Chuanbo Shen, Wiesinger, M. & Qian Zhang. 2013. Late-stage foreland growth of China's largest orogens (Qinling, Tibet): evidence from the Hannan-Micang crystalline massifs and the northern Sichuan Basin, central China. Lithosphere, 5(4): 420–37.
- Zhao Yun-xiang, Chen Jing-shan, Ding Xiong, Yao Jing-li, Li Shi-xiang & Li Ning. 2013. Vertical structure and dominating factors of Chang 9 sandbody from Yanchang Formation in Ordos Basin. Acta Sedimentologica Sinica, 31(1): 77–88.
- Zheng Yue-juan, Su Fei, Chen Shu-wang, Zhang Jian, Huang Xin & Gong Fan-hao. 2013. Inner happiness Balinyouqi Lower Triassic fossil discoveries of the road group. Geological Bulletin of China, 32(9): 1423–1435.
- Zhonghe Zhou & Antunes, M.T. 2013. Terrestrial Mesozoic stratigraphy. Ciências da Terra, 18: 95–98.
- Zhu, G., Su, J., Yang, H., Wang, Y., Fei, A., Liu, K., Zhu, Y., Hu, J. & Zhang, B. 2013. Formation mechanisms of secondary hydrocarbon pools in the Triassic reservoirs of the northern Tarim Basin. Marine and Petroleum Geology, 46: 51–66.
- Zhu Hai-hua, Zhong Da-kang, Li Qi-rong, Du Ben-qiang,

Long Hui, Ma Yan-liang, Wu Lian-bo & Jiang Nan. 2013. Characteristics and controlling factors of Upper Triassic Xujiahe tight sandstone reservoir in southern Sichuan Basin. Acta Sedimentologica Sinica, 31(1): 167–175.

- Zhu Ru-kai, Bai Bin, Yuan Xuan-jun, Luo Zhang, Wang Pei, Gao Zhi-yong, Su Ling & Li Ting-ting. 2013. A new approach for outcrop characterization and geostatistical analysis of meandering channel sandbodies within delta plain setting using digital outcrop models: Upper Triassic Yanchang tight sandstone formation, Yanhe outcrop, Ordos Basin. Acta Sedimentologica Sinica, 31(5): 867–877.
- Zhu Tongxing, Feng Xintao, Wang Xiaofei & Zhou Mingkui. 2013. Reconstruction of the Triassic tectonic lithofacies paleogeography in Qiangtang region, northern Qinghai-Tibet Plateau, China. Acta Geologica Sinica, 87(2): 378–394.
- Zonneveld, J.-P. & Bistran, R. 2013. Bored bivalves in Upper Triassic (Norian) event beds, northeastern British Columbia, Canada. Ichnos, 20(2): 88–98.
- Zou Guangfu, Pan Zhongxi, Zhuang Zhonghai, Zhu Tongxing, Li Jianzhong & Feng Xintao. 2013. Phanerozoic paleomagnetism characteristics of the Qomolangma area in Tibet. Acta Geologica Sinica, 87(2): 517–527.
- Zulauf, G., Blau, J., Dörr, W., Klein, T., Krahl, J., Kustatscher, E., Petschick, R. & van de Schootbrugge, B. 2013. New U-Pb zircon and biostratigraphic data of the Tyros Unit, eastern Crete: constraints on Triassic palaeogeography and depositional environments of the eastern Mediterranean, Zeitschrift der Deutschen Gesellschaft für Geowissenschaften, 164(2): 337–352.
- Żyła, D., Wegierek, P., Owocki, K. & Niedźwiedzki, G. 2013. Insects and crustaceans from the latest Early-early Middle Triassic of Poland. Palaeogeography, Palaeoclimatology, Palaeoecology, 371: 136–144.

Addendum to Warrington, G. 2014. New Triassic Litertaure. Albertiana, 41: 50–81.

- Axsmith, B.J. 2009. A new *Cynepteris* from the Upper Triassic of Arizona: potential implications for the early diversification of Schizaealean ferns. International Journal of Plant Sciences, 170(5): 657–665.
- Benson, R.B.J., Evans, M. & Druckenmiller, P.S. 2012. High diversity, low disparity and small body size in plesiosaurs (Reptilia, Sauropterygia) from the Triassic-Jurassic boundary. PLoS ONE, 7(3): e31838. doi:10.1371/journal. pone.0031838.
- Bomfleur, B., Taylor, E.L., Taylor, T.N., Serbet, R., Krings, M. & Kerp. H. 2011. Systematics and paleoecology of a new peltaspermalean seed fern from the Triassic polar vegetation of Gondwana. International Journal of Plant Sciences, 172(6): 807–835.
- Burliga, S., Koyi, H.A. & Chemia, Z. 2012. Analogue and numerical modelling of salt supply to a diapiric structure rising above an active basement fault. Geological Society, London, Special Publications, 363: 395-408.

- Butler, R.J., Barrett, P.M. & Gower, D.J. 2012. Ressessment of the evidence for postcranial skeletal pneumaticity in Triassic archosaurs, and the early evolution of the avian respiratory system. PLoS ONE, 7(3): e34094. doi:10.1371/journal. pone.0034094.
- Butler, R.J., Brusatte, S.L., Reich, M., Nesbitt, S.J., Schoch, R.R. & Hornung, J.J. 2011. The sail-backed reptile *Ctenosauriscus* from the latest Early Triassic of Germany and the timing and biogeography of the early archosaur radiation. PLoS ONE, 6(10): e25693. doi:10.1371/journal.pone.0025693.
- Callot, J.-P., Trocmé, V., Letouzey, J., Albouy, E., Jahani, S. & Sherkati, S. 2012. Pre-existing salt structures and the folding of the Zagros mountains. Geological Society, London, Special Publications, 363: 545–561.
- Chen Fei, Hu Guangyi, Sun Lichun, Fan Tailiang, Gao Zhiqian, Yu Xitong, Pang Zhenglian & Li Rangbin. 2012. Characteristics of sedimentary facies and evolution in sequence stratigraphic framework of the Upper Triassic Yanchang Formation in southern Ordos Basin. Journal of Palaeogeography, 14(3): 321–330.
- Cisneros, J.C., Cabral, U.G., de Beer, F., Damiani, R. & Fortier, D.C. 2010. Spondarthritis in the Triassic. PLoS ONE, 5(10): e13425. doi:10.1371/journal.pone.0013425.
- Colombi, C.E., Fernández, E., Currie, B.S., Alcober, O.A., Martínez, R. & Correa, G. 2012. Large-diameter burrows of the Triassic Ischigualasto Basin, NW Argentina: paleoecological and paleoenvironmental implications. PLoS ONE, 7(12): e50662. doi:10.1371/journal.pone.0050662.
- Deng Xiuqin, Fu Jinhua, Yao Jingli, Pang Jinlian & Sun Bo. 2011. Sedimentary facies of the Middle-Upper Triassic Yanchang Formation in Ordos Basin and breakthrough in petroleum exploration. Journal of Palaeogeography, 13(4): 443–455.
- Escapa, I.H., Decombeix, A.-L., Taylor, E.L. & Taylor, T.N. 2010. Evolution and relationships of the conifer seed cone *Telemachus*: evidence from the Triassic of Antarctica. International Journal of Plant Sciences, 171(5): 560–573.
- Feldman-Olszewska, A., Adamczak-Biały, T. & Becker, A. 2012. Characterization of the Jurassic and Triassic reservoirs and seals from north Mazovia as a candidate site for CO_2 storage based on data from deep boreholes. Biuletyn Państwowy Instytut Geologiczny, 448(1): 27–46.
- Feng Juanping, Li Wenhou & Ouyang Zhengjian. 2012. Sedimentary characters and geological implication of turbidite of the Chang 6 and Chang 7 intervals of the Upper Triassic Yanchang Formation in Huangling area, Ordos Basin. Journal of Palaeogeography, 14(3): 295–302.
- Ferrer, O., Jackson, M.P.A., Roca, E. & Rubinat, M. 2012. Evolution of salt structures during extension and inversion of the Offshore Parentis Basin (eastern Bay of Biscay). Geological Society, London, Special Publications, 363: 361–379.
- Fröbisch, J. 2008. Global taxonomic diversity of anomodonts (Tetrapoda, Therapsida) and the terrestrial rock record across the Permian-Triassic boundary. PLoS ONE, 3(11): e3733. doi:10.1371/journal.pone.003733.
- Fu Jinhua, Li Shixiang, Liu Xianyang & Deng Xiuqin. 2012. Sedimentary facies and its evolution of the Chang 9 interval of Upper Triassic Yanchang Formation in Ordos Basin. Journal

of Palaeogeography, 14(3): 269-84.

- Gallois, R.W. 2009. The lithostratigraphy of the Penarth Group (Late Triassic) of the Severn Estuary area. Geoscience in South-West England, 12(2): 71–84.
- Garcia, M. de A., Ribeiro, H.J.S., de Souza, A.S. & Triguis, J.A. 2011. Correlation between faciology and organic geochemistry of the Pirambóia Formation tar sands, Betumita Farm, Triassic of the Paraná Basin, São Paulo State. Geociências (Sao Paulo), 30(3): 357–369.
- González-León, C.M., Lawton, T.F. & Weber, R. 2011. Estratigrafia del Triásico y el Jurásico inferior de Sonora, México. Boletín del Instituto de Geologia, 118: 57–80.
- Grasby, S.E., Sanei, H. & Beauchamp, B. 2011. Catastrophic dispersion of coal fly ash into oceans during the latest Permian extinction. Nature Geoscience, 4: 104–107.
- Guinot, G., Adnet, S. & Cappetta, H. 2012. An analytical approach for estimating fossil record and diversification events in sharks, skates and rays. PLoS ONE, 7(9): e44632. doi:10.1371/journal.pone.0044632.
- Holliday, C.M., Ridgely, R.C., Sedlmayr, J.C. & Witmer, R.M. 2010. Cartilaginous epiphyses in extant archosaurs and their implications for reconstructing limb function in dinosaurs. PLoS ONE, 5(9): e13120. doi:10.1371/journal. pone.0013120.
- Jakóbczk, S. & Kowalczyk, A. 2010. Groundwater chemistry of the Triassic carbonate aquifer in the Bibiela well field area in view of geochemical modelling. Biuletyn Pánstwowego Instytutu Geologicznego, 442: 79–84.
- Kitahara, M.V., Cairns, S.D., Stolarski, J., Blair, D. & Miller, D.J. 2010. A comprehensive phylogenetic analysis of the Scleractinia (Cnidaria, Anthozoa) based on mitochondrial CO1 sequence data. PLoS ONE, 5(7): e11490. doi:10.1371/ journal.pone.0011490.
- Klein, N. 2010. Long bone histology of Sauropterygia from the Lower Muschelkalk of the Germanic Basin provides unexpected implications for phylogeny. PLoS ONE, 5(7): e11613. doi:10.1371/journal.pone.0011613.
- Kowal-Linka, M. & Bodzioch, A. 2012. Bonebeds with *Dactylosaurus* (Reptilia, Sauropterygia) from the Röt (Lower Triassic, Olenekian) in the Opole Silesia region (S Poland). Przegląd Geologiczny, 60(12): 646–649.
- Lin Yu, Wu Shenghe, Xu Zhangyou & Chen Ce. 2012. Applying multi-scale data to study diagenetic reservoir facies of the Upper Triassic Xujiahe Formation in Dayi Structure, western Sichuan Province. Journal of Palaeogeography, 14(4): 522–532.
- Liu Haowei, Wang Jian, Liu Qunming & Zhao Xin. 2012. Favorable reservoir distribution and its controlling factors in the Chang 8 interval of Upper Triassic Yanchang Formation in Jiyuan area, Ordos Basin. Journal of Palaeogeography, 14(3): 285–294.
- Mah, C.L. & Blake, D.B. 2012. Global diversity and phylogeny of the Asteroidea (Echinodermata). PLoS ONE, 7(4): e35644. doi:10.1371/journal.pone.0035644.
- Mander, L., Collinson, M.E., Chaloner, W.G., Brain, A.P.R. & Long, D.G. 2012. The ultrastructure and botanical affinity of the problematic mid-Mesozoic palynomorph *Ricciisporites*

tuberculatus Lundblad. International Journal of Plant Sciences, 173(4): 429–440.

- Mander, L., Wesseln, C.J., McElwain, J.C. & Punyasena, S.W. 2012. Tracking taphonomic regimes using chemical and mechanical damage of pollen and spores: an example from the Triassic-Jurassic mass extinction. PLoS ONE, 7(11): e49153. doi:10.1371/journal.pone.0049153.
- Martinez, R.N. & Alcober, O.A. 2009. A basal sauropodomorph (Dinosauria: Saurischia) from the Ischigualasto Formation (Triassic, Carnian) and the early evolution of Sauropodomorpha. PLoS ONE, 4(2): e4397. doi:10.1371/ journal.pone.004397.
- Martz, J.W. & Parker, W.G. 2010. Revised lithostratigraphy of the Sonsela Member (Chinle Formation, Upper Triassic) in the southern part of Petrified Forest National Park, Arizona. PLoS ONE, 5(2): e9329. doi:10.1371/journal.pone.009329.
- Min Li, Ying Tian, Ying Zhao & Wenjun Bu. 2012. Higher level phylogeny and the first divergence time estimation of *Heteroptera* (Insecta: Hemiptera) based on multiple genes. PLoS ONE, 7(2): e32152. doi:10.1371/journal. pone.0032152.
- Mueller, C. & Reinhold, K. (eds.). 2011. Geological characterisation of deep-seated reservoir and barrier horizons in Germany. Schriftenreihe der Deutschen Gesellschaft fur Geowissenschaften, 74: 244 pp.
- Onoue, T., Sato, H., Nakamura, T., Noguchi, T., Hidaka, Y., Shirai, N., Ebihara, M., Osawa, T., Hatsukawa, Y., Toh, Y., Koizumi, M., Harada. H., Orchard, M.J. & Nedachi, M. 2012. Deep-sea record of impact apparently unrelated to mass extinction in the Late Triassic. Proceedings of the National Academy of Sciences, 109(47): 19134–19139.
- Pan Liyin, Liu Zhanguo, Li Chang, Shou Jianfeng, Zhang Jianyong, Zhang Jie, Shen Anjiang, Zhou Jingao & Zheng Xingping. 2012. Dolomitization and its relationship with reservoir development of the Lower Triassic Feixianguan Formation in eastern Sichuan Basin. Journal of Palaeogeography, 14(2): 176–186.
- Petersen, H.I. & Lindström, S. 2012. Synchronous wildfire activity rise and mire deforestation at the Triassic-Jurassic boundary. PLoS ONE, 7(10): e47236. doi:10.1371/journal. pone.0047236.
- Qiang Zitong, Zeng Deming, Wang Xingzhi & Wu Shijiu. 2012. Isotopic geochemical characteristics of oolitic bank dolostones in the Lower Triassic Feixianguan Formation in northeastern Sichuan Province. Journal of Palaeogeography, 14(1): 13–20.
- Qiao Yanguo, Shi Zhiqiang, Wang Yanyan & Zhang Biao. 2012. The Late Permian- Early Triassic cyclostratigraphy in Shangsi section of Guangyuan area, Sichuan province: implications for P-T geological event based on wavelet analysis. Journal of Palaeogeography, 14(3): 403–410.
- Quintà, A., Tavani, S. & Roca, E. 2012. Fracture pattern analysis as a tool for constraining the interaction between regional and diapir-related stress fields: Poza de la Sal Diapir (Basque Pyrenees, Spain). Geological Society, London, Special Publications, 363: 521–532.
- Salamon, M.A., Niedźwiedzki, R., Lach, R., Brachaniec, T. & Gorzelak, P. 2012. Ophiuroids discovered in the Middle

Triassic hypersaline environment. PLoS ONE, 7(11): e49798. doi:10.1371/journal.pone.0049798.

- Sander, P.M., Xiaohong Chen, Long Cheng & Xiaofeng Wang. 2011. Short-snouted toothless ichthyosaur from China suggests Late Triassic diversification of suction feeding ichthyosaurs. PLoS ONE, 6(5): e19480. doi:10.1371/journal. pone.0019480.
- Shi Zhensheng, Zhao Zhengwang, Jin Hui & Song Haijing. 2012. Depositional characteristics and its geological significance of the Upper Triassic Xiaotangzi Formation in Sichuan Basin. Journal of Palaeogeography, 14(4): 477–486.
- Shi Zhiqiang, An Hongyan, Yi Haisheng, Zhang Hua & Zeng Deyong. 2011. Classification and characters of the Early Triassic anomalous carbonate rocks in Upper Yangtze area. Journal of Palaeogeography, 13(1): 1–10.
- Soares Júnior, A.V., Hasui, Y., Costa, J.B.S. & Machado, F.B. 2011. Rifting evolution and paleogeography of the Brazilian Equatorial Atlantic Margin: from Triassic to Holocene. Geociências (Sao Paulo), 30(4): 669–692.
- Tapanila, L. & Roberts, E.M. 2012. The earliest evidence of holometabolan insect pupation in conifer wood. PLoS ONE, 7(2): e31668. doi:10.1371/journal.pone.0031668.
- Tian Jingchun, Wang Wenzhi, Wang Feng, Wang Weihong & Zhang Jinquan. 2012. Palaeoearthquake deformation structures of the Chang 6 interval of Upper Triassic Yanchang Formation and superimposed relationship between sandbodies in Huachi-Qingyang area of Ordos Basin. Journal of Palaeogeography, 14(3): 303–310.
- Vanschoenwinkel, B., Pinceel, T., Vanhove, M.P.M., Denis, C., Jocque, M., Timms, B.V. & Brendonck, L. 2012. Toward a global phylogeny of the "living fossil" crustacean order of the Notostraca. PLoS ONE, 7(4): e34998. doi:10.1371/journal. pone.0034998.
- Wang Lan, Li Wenhou, Lin Tong & Wang Ruogu. 2012. Diagenesis and its influencing factors of the Chang 6 interval of Upper Triassic Yanchang Formation in Ordos Basin. Journal of Palaeogeography, 14(3): 311–320.
- Wang Yanyan, Zhang Biao, Shi Zhiqiang & Yi Haisheng. 2012. Oxygen and carbon isotopic records of the Late Triassic Carnian pluvial event deposits in northwestern Sichuan Province. Journal of Palaeogeography, 14(3): 375–382.
- Wu Shenghe, Fan Zheng, Xu Changfu, Yue Dali, Zheng Zhan, Peng Shouchang & Wang Wei. 2012. Internal architecture in alluvial fan in the Triassic Lower Karamay Formation in Karamay Oilfield, Xinjiang. Journal of Palaeogeography, 14(3): 331–340.
- Yang Wentao, Wang Xiaofeng, Yang Jianghai & Du Yuansheng. 2011. Soft-sediment deformation structures caused by palaeoearthquake in the Middle-Late Triassic in Yima area, western Henan Province. Journal of Palaeogeography, 13(6): 635–644.
- Yates, A.M., Neumann, F.H. & Hancox, P.J. 2012. The earliest post-Paleozoic freshwater bivalves preserved in coprolites from the Karoo Basin, South Africa. PLoS ONE, 7(2): e30228. doi:10.1371/journal.pone.0030228.
- Yukawa, H., Terada, K., Ge Sun & Suzuki, S. 2012. The oldest fossil forest in Japan discovered from the Upper

Triassic Nariwa Group, Okayama Prefecture, SW Japan – implication for reconstruction of depositional environment and paleovegetation. Okayama University Earth Science Reports, 19(1): 25–37.

Zhang Tianfu, Shou Jianfeng, Zheng Xingping, She Min & Pan Liyin. 2012. Spatial distribution and characterization of pore and throat of oolitic dolostone of the Lower Triassic Feixianguan Formation in northeastern Sichuan Province. Journal of Palaeogeography, 14(2): 187–196.



Albertiana Forum

Towards an evolutionary sound definition of 'recovery'?

In the Early Triassic community, 'biotic recovery' is one of the most commonly used terms and a recurring debate is whether the biotic recovery was prolonged in the aftermath of the end-Permian crisis. The problem with this debate is that 'recovery' is understood in different ways by different authors. The latest review about the pattern and tempo of the biotic recovery following the end-Permian mass extinction (EPME) was written by Chen and Benton (2012). Although Chen and Benton are well aware of definition issues with the term 'recovery' (Chen and Benton, 2012, p. 4), their paper still implicitly and alternatively adopts various definitions of 'recovery'. In these regards, their paper illustrates well the wandering of the current literature and this is the reason I am discussing it here. My intention is not to criticize Chen and Benton's paper per se but to draw our attention towards this pervading conceptual issue.

In their 2012 paper, Chen and Benton reviewed the latest results about the biotic recovery following the end-Permian mass extinction (EPME) and they concluded that Triassic ecosystems were rebuilt stepwise from low to high trophic levels. Many of us reacted strongly to that paper, especially to their figure 4 (Chen and Benton, 2012, p. 7) and their claim that "immediate postextinction ecosystems in the Griesbachian-Dienerian show only the lowest trophic level" (Chen and Benton, 2012, p.7), that is, primary producers, which they equate to microbes. It is obvious (as mentioned several times by Chen and Benton) that many more groups and other trophic levels were represented during this earliest Triassic interval [e.g. ammonoids (Brayard et al., 2009) and conodonts (Orchard, 2007), some being predators presumably (Purnell, 1995); see also Scheyer et al., 2014].

Then, how can we reconcile their figure and claim with the asserted presence of other organisms during that interval? I shall assume that Chen and Benton (2012) implicitly considered a given trophic level to be present only if it has achieved full recovery. This, in turn, raises the issue of defining the term 'recovery' for a trophic level. One recurrent definition used by Chen and Benton (2012) for a group or a community is a low abundance, high diversity and high evenness of that community. Let us take over this definition and have a look again at the Griesbachian data. Where is the evidence for a high diversity and low evenness of microbial communities in the Griesbachian? How do we know that the lowest trophic level had already recovered? The short-lived dominance of microbes in the earliest Griesbachian could also suggest that the corresponding microbes were ,disaster taxa'. The same question applies to the other trophic levels, which, following Chen and Benton, were added sequentially on top of one another. Chen and Benton's figure 4 demonstrates that they did not apply the same definition of 'recovery' consistently throughout their paper.

In fact, like Chen and Benton (2012) we all implicitly and alternatively adopt various definitions of 'recovery'. My intention

here is to stress out our need for either a consensual definition of ,recovery⁶ or the global acceptance by our community that discussions about recovery should be better contextualized. Recovery of a species, of a supra-generic clade, of a community, of an ecosystem or of a trophic level may be defined and quantified in different and possibly contradicting ways. The granularity (spatial, temporal or ecological resolution) we choose for our studies may influence our definition and hence our conclusions about the tempo of recovery. Moreover, although the recovery of a particular taxonomic group can be easily defined in terms of, for instance, its specific diversity, defining the recovery of a complex system with many interacting constituents can be challenging.

A generic, maybe largely approved definition of 'recovery' is "the return to a previous state" (Chen and Benton, 2012, p.3). Yet, this definition does not specify what the 'state' of a clade or of an ecosystem should be. For a given taxon, the most used metric of its state of recovery is its morphological diversity. One could argue that the state of recovery is better characterized by a set of parameters such as diversity, size, and community evenness, as well as the spatial distribution thereof, that is, whether the corresponding recovery signature is observed locally or globally.



Figure 1 – Fictitious temporal evolution of an ecosystem's trophic pyramid during a recovery.

Yet, what is a good metric for the recovery of an ecosystem? Individual taxonomic groups show major differences in their response to the EPME: ammonoids thrived less than 2 myr after the EPME (Brayard et al., 2009) but corals are unknown from the fossil record until the Middle Triassic. Shall we compute an average of the groups' recoveries? Remember that some groups like the conodonts were positively affected (immediate recovery) and some groups like the trilobites never recovered. How can we appropriately describe the 'state' of an ecosystem?

The familiar way to understand recovery of a given system implies a relative stability (unity) of that system. If the system evolves too fast, we are not able to identify (a) normal, healthy state(s) in comparison with which we can define abnormal, crisis periods. The faster the system evolves in normal conditions, the shorter and tougher the crisis has to be in order for us to detect it and to define its boundaries. In other words, defining ecosystem descriptors that are less prone to variation is key for the identification of healthy steady states and crisis transitional phases.

Recently, Foster and Twitchett (2014) showed that there had been no loss of functional diversity of benthics through the EPME at the global scale. This is not surprising if one considers that the global structure of the network of ecological interactions within an ecosystem is likely to evolve much slower than its individual taxonomic groups: a given group of organisms may disappear but its ecological position or niche will eventually be filled by another group. Even if one expects the network of ecological interactions to co-evolve with the individual taxa, the relative stability of functional ecology makes it a better descriptor of the ecosystem than its ever-changing constitutive organisms. In these regards, Chen and Benton's suggestion (Chen and Benton, 2012) to using a trophic model of recovery for the EPME, a crisis that possibly lasted millions of years, is praiseworthy.

Let us assume that the shape of a particular trophic level is represented in three dimensions by, for instance, the abundance, diversity and evenness of the corresponding community. The state of the ecosystem is then described by a trophic pyramid, whose shape in this three- dimensional space is not necessarily a geometric pyramid (Fig. 1). Using this ecosystem descriptor, recovery could be defined as the restoration of a pre-crisis pyramid shape. As mentioned above, Chen and Benton's bottom-up restoration model is not supported by the data and it calls for a critical reappraisal. Alternative scenarios should be explored. For instance, the recovery trajectory may have instead involved a lateral expansion of the proposed trophic pyramid through increase of both diversity and evenness at all ecological levels (Fig. 1). 87

Besides the practical issues of identifying the ecological role of past organisms, in particular the role of taxa without extant analogs such as conodonts and ammonoids, one may argue that the term ,recovery' sensu stricto implies that the post-crisis steady state (recovery state) is essentially the same as the pre-crisis state. Since the biosphere is likely to evolve drastically during major crises such as the EPME, it is not excluded that, when enough data is gathered, we have to recognize that the ecosystem did not, strictly speaking, recover from the EPME but went through what is called a critical transition: a transition from one equilibrium of the ecosystem to a new, distinct equilibrium.

This work was supported by the Swiss SNF fellowship PA00P1-145314.

Nicolas Goudemand

Paleontological Institute and Museum, University of Zurich, nicolas.goudemand@pim.uzh.ch

References Cited

- Brayard, A., Escarguel, G., Bucher, H., Monnet, C., Brühwiler, T., Goudemand, N., Galfetti, T. & Guex, J. 2009. Good genes and good luck: ammonoid diversity and the end-Permian mass extinction. Science, 325: 1118–1125.
- Chen, Z.Q. & Benton, M.J. 2012. The timing and pattern of biotic recovery following the end-Permian mass extinction. Nature Geoscience 5: 375–383.
- Foster, W.J. & Twitchett, R.J. 2014. Functional diversity of marine ecosystems after the Late Permian mass extinction event. Nature Geoscience, 7(3): 233–238.
- Orchard, M.J. 2007. Conodont diversity and evolution through the latest Permian and Early Triassic upheavals. Palaeogeography, Palaeoclimatology, Palaeoecology, 252: 93–117.
- Payne, J.L., Summers, M., Rego, B.L., Altiner, D., Yu, Y., Wei, J. & Lehrmann, D.J. 2011. Early and Middle Triassic trends in diversity, evenness, and size of foraminifers on a carbonate platform in south China: implications for tempo and mode of biotic recovery from the end-Permian mass extinction. Paleobiology, 37: 409–425.
- Purnell, M.A. 1995. Microwear on conodont elements and macrophagy in the first vertebrates. Nature, 374: 798–800.
- Scheyer, T.M., Romano, C., Jenks, J. & Bucher, H. 2014. Early Triassic marine biotic recovery: the predators' perspective. PLoS ONE, 9(3):e8898. doi: 10.1371/journal. pone.0088987.

Albertiana publishes letters to the STS community including comments on past articles. Editors will solicit a reply from original authors if the comment is approved

Publication Announcement

Stratigraphie von Deutschland XI: Buntsandstein. J. Lepper und H.-G. Röhling, Editors. 2013. 657 pp., 275 figs., 45 tab., 1 pl. Schriftenreihe der Deutschen Gesellschaft für Geowissenchaften. vol 69. ISBN 978-3-510-49229-9; € 109,00

As the 11th volume of the series 'Stratigraphy of Germany', the Buntsandstein Monograph has been issued with scientific contributions from 27 authors, mainly from the federal Geological State Surveys, The central theme of the monograph covers the stratigraphy of the German Buntsandstein (Lower Triassic) in the Germanic Basin. It is accompanied by a geoscientific historical review on Buntsandstein research since the 18th Century, followed by a palaeogeographical overview.

The main chapter on the Buntsandstein stratigraphy explains the dual stratigraphy agreed by the German Stratigraphic Commission in 2000. The lithostratigraphic component of the historical units is retained in the seven formations Calvörde, Bernburg, Volpriehausen, Detfurth, Hardegsen, Solling and Röt, whereas the regional geochronological component determines the seven Folgen s1 to s7, which have quasi-isochronous boundaries. In wide areas of the Buntsandstein, the boundaries of these formations and Folgen coincide. The synopsis reflects the combined outcome of several working groups and panels, including those of the German State Geological Surveys.

The dual classification scheme of the Buntsandstein can be correlated throughout the major part of the German section of the Buntsandstein depository. It comprises the following units that are partly separated by unconformities with variable characters:

(Muschelkalk)	
Röt-Formation	s7-Folge 6-Folge (incl. the S-Unconformity)
H-Unconformity	
Hardegsen-Formation	s5-Folge
Detfurth-Formation	s4-Folge
D-Unconformity	
Volpriehausen-Formation	s3-Folge
V-Unconformity	
Bernburg-Formation	s2-Folge
Calvörde-Formation	s1-Folge

(Zechstein)

The lower boundary of the Buntsandstein Group is in the uppermost Permian Changhsingian stage. Thus a very small portion of the Buntsandstein belongs to the Permian. The upper boundary of the Buntsandstein between the Myophorien beds of the Buntsandstein and the Grenzgelbkalk of the Muschelkalk is of early Anisian age.

The "Bröckelschiefer", which was previously attributed to the lowest Buntsandstein, in fact represents a transitional facies with a saliniferous component. It has been now assigned to the Zechstein Group.

The siliclastic, fining-upward Calvörde Formation (Folge s1) begins in central basin areas with sandy strata, represented by the Heigenbrücken Sandstone or the lower Eck Conglomerate. The base of the Bernburg Formation (Folge s2) begins with a distinct increase in the proportion of sandstone, as well as oolitic layers. In the coarser clastic proximal-marginal facies this lower boundary can not always be detected.

The Middle Buntsandstein consists of four formations. It starts with the coarse-grained Volpriehausen Sandstone, a member of the Volpriehausen Formation (Folge s3), which is recognizable towards the southern basin margin by a distinct conglomeratic sandstone. A comparable coarse-clastic base and fining upward sequence also characterizes the Detfurth and Hardegsen Formations. Epeirogenetic/taphrogenetic movements were active during sedimentation of the Folge s5 (Hardegsen Formation) and reached a climax towards its end, which lead to differential and widespread erosion of previously-deposited strata. Above this major erosional unconformity, coarse basal sediments of the Solling Formation (Folge s6) were accumulated. In contrast to older layer-cake deposits of the Middle and Lower Buntsandstein, this formation is characterized by local facies changes. It thins out towards the southern basin margin into coarse clastic sediments with multiple pedogenic intervals and extensive hiatuses.

Above the siliciclastic Solling Formation (Folge s6) of the Middle Buntsandstein Subgroup there follows the partiallyevaporitic Upper Buntsandstein Subgroup. Stratigraphically it is also classified as Röt Formation and Folge s7. It is subdivided into the six members Vitzenburg, Göschwitz, Glockenseck, Karsdorf,

Gleina and Dornburg, and in some areas also into the four Subfolgen s7-1 to s7-4. In contrast to the basinal evaporitic facies, the marginal fluvial-lacustrine facies is dominated by a condensed siliciclastic succession interrupted proximally by hiatuses and overprinted by stacked palaeosols, which prevent any detailed assignment of lithostratigraphic units.

These central achievements are accompanied by explanations



of the biostratigraphic subdivision, of the magneto- and cyclostratigraphy and of the geological time scale, of the climate of the Buntsandstein time, of palaeosols in the Buntsandstein and their stratigraphic relevance, and by a status report on the sequence-, base level- and cyclostratigraphy.

Special insights on regional characteristics of the German Buntsandstein are given by 10 individual contributions. Finally, the Buntsandstein Monograph encompasses further 5 contributions on the lithology and stratigraphy in the adjacent countries of Netherlands and NE-Belgium, Denmark, Poland, NE-France, and eastern Alps.

Meeting annoucements





Have a news item or meeting annoucement you would lilke published in *Albertiana*? Please contact the editor.