

DEȘLI CAIRA (ROMANIA) OLENEKIAN-ANISIAN BOUNDARY SECTION: REVISED STRATIGRAPHY AND NEW FINDINGS

**Micha Horacek¹, Marco Balini^{2, 3}, Marius Stoica⁴, Valentin Răuță⁴,
Alexandra-Daniela Lăcătuș⁴, and Iuliana Lazăr^{3, 4}**

¹Department of Lithospheric Research, Vienna University, Josef-Holaubek-Platz 2, 1090 Vienna, Austria.
untertrias@googlemail.com

²Dipartimento di Scienze della Terra “Ardito Desio”, Università degli studi di Milano, Via Mangagalli 34, 2033, Milano, Italy. *marco.balini@unimi.it*

³Research Institute of the University of Bucharest, 90-92, Panduri Road, 5th District, Bucharest, Romania

⁴University of Bucharest, Faculty of Geology and Geophysics, Doctoral School of Geology, 1 N. Balcescu Bd, 010041 Bucharest, Romania

Abstract – We provide an update on the ongoing investigation of the Deșli Caira section, Dobrogea, Romania, candidate section for the Anisian stage GSSP. We report on a discussion about a fault in the section, which resulted in a comment in a different paper, and demonstrate that the fault is present, cutting the section almost perpendicular and continuing from the lower part to the top of the section. However, as long as the fault’s position is recognized, it does not affect research in the section, because there is sufficient room to work on both sides; nevertheless, we recommend that all work be conducted on the same (left) side of the fault throughout the section just to rule out any potential mistakes, even though, when taking it into account accordingly, one can move from one side of the fault to the other without problems. Furthermore, we suspect the occurrence of *Pseudacrochordiceras*, that in Nevada is limited to the uppermost part of the *Neopopanoceras haugi* ammonoid zone, just below the boundary with the overlying *Courtilloticeris stevensi* ammonoid zone are present in the Deșli Caira section. If confirmed, this would enable long-range correlation with data from Nevada (USA) and thus, the Panthalassa realm.

INTRODUCTION

The Deșli Caira section of North Dobrogea, Romania has been under investigation for more than 100 years, beginning with the work of Kittl (1908) and Simionescu (1910, 1911), and later continued by Mirăuță (1974) and Mirăuță and Gheorghian (1978). Then, in the 1990’s, when the Chios section became unsuitable for further consideration as a GSSP for the Anisian stage due to the presence of a fault, the Deșli Caira section became a candidate for the GSSP of the Anisian stage, which resulted in increased scientific activities, mainly lead by Eugen Grădinaru. In 2000 a field workshop conducted at the section resulted in several publications as well as a formal proposal for Deșli Caira as a GSSP candidate (Grădinaru et al., 2006, 2007). However, Grădinaru failed to publish the ammonoid record of the section, which was deemed crucial for the definition of the boundary, and

for the next ~10 years little to no progress was made, even though Grădinaru did publish a few additional articles.

Then, in the early 2020’s, two more potential candidates were proposed for the GSSP of the Anisian stage, i.e., the Kçira section in Albania and the Wantou section in China, both of which were accompanied by supporting publications (Muttoni et al., 2019, Chen et al., 2020). Meanwhile, a few additional articles were published to further support Deșli Caira’s candidate status, e.g., Golding (2021), Grădinaru (2022), Horacek and Grădinaru (2023), and, most recently, Balini et al., (2024) and Golding, (2025). While the work of Balini et al., (2024) was intended to be a continuation of Grădinaru’s ammonoid research, which unfortunately was never published, that of Golding (2025) was the continuation of earlier work on Deșli Caira conodonts (Orchard et al., 2007; Golding, 2021). However, both of these latest contributions require further attention. The

Published online: 4 June, 2026




Figure 1– Geology of North Dobrogea, south-eastern Romania, with the position of Deșli Caira section, part of the Tulcea Unit (=T.U., after Grădinaru et al., 2007). The inset shows the position of the Tulcea Unit in the North Dobrogean Orogen.

lithology, which was published in Balini et al., (2024), needs a few minor revisions in the lower part even though this part of the stratigraphic column had not been the focus of the Balini et al., (2024) article. These revisions are in progress and will be published in an article currently in preparation. The work of Golding (2025) contains some serious ambiguities concerning a fault in the Deșli Caira section and the exact positions from which samples were taken. Even though two of the present authors reviewed the Golding manuscript and both independently requested the exact sampling positions with respect to the fault's location, the author and editors, for whatever reason, apparently did not think these requests had merit and the manuscript was published without this crucial information. This situation seemed unacceptable to the present authors, because it could lead to differing views on the presence and position of the fault and the accuracy of the results provided by Golding (2025). Therefore, we have written an article (Balini et al., 2026) commenting on our concerns with Golding (2025) and identified the problems arising from the absence of a description of the fault's position with respect to their sampling locations. Obviously, it must be stressed that in any stratigraphic investigation it is essential that the section be screened for the presence of faults in order to avoid incorrect results (Krystyn et al., 2019, Balini et al., 2026).

In the present article we want to make certain that everyone understands the relevance of the fault running more or less perpendicular to the bedding, resulting in an offset of ca. 0.8 to 1m between the right and left sides of the fault, which causes a serious problem when not identified (or recognized) if samples were collected on both sides of the fault. Also, we want to make clear that the fault in no way disqualifies Deșli Caira as a GSSP candidate (and in our opinion, the best), because there is ample space on both sides of the fault, thus enabling sampling without the need to cross it. Furthermore, as stated by Balini et al., (2026), if the fault is recognized and accounted for appropriately, it is possible to work and sample on both sides. While we have only included published material and data in our comments on Golding 2025 (Balini et al., 2026), we now also

present additional evidence on and photographs of the fault in the Deșli Caira section.

LOCALITY

The Deșli Caira section is positioned south of the Danube Delta in North Dobrogea Orogen, in south-eastern Romania, close to the Black Sea coast (Fig. 1). GPS coordinates for the section are as follows: 28°48'08"E, 45°04'27"N (Grădinaru et al., 2007). Paleontological investigations of the locality were first conducted by Kittl (1908) and Simionescu (1910, 1911), and later continued by Mirăuță (1974), and Mirăuță and Gheorghian (1978). Grădinaru et al. (2006, 2007) then published numerous articles in collaboration with an earlier working group, of which the latter work included a partial report on some of the ammonoids as well a report on the Deșli Caira foraminiferas and conodonts. Golding (2021, 2025) recently updated the conodont record, Grădinaru & Gaetani (2019) reported on the brachiopod record and Forel and Grădinaru (2021) reported on the ostracod fauna. The paleomag signal was also investigated by Grădinaru et al., (2007), as was the carbonate $\delta^{13}\text{C}$ evolution of the section by Atudorei (1998) and Horacek & Grădinaru (2023).

The Deșli Caira section is composed of the Agighiol Limestone, a medium to thick massive-bedded limestone named after a neighboring hill and nearby village (Grădinaru et al., 2007). It was deposited on a carbonate platform developed in the Tulcea Unit (Fig. 1), which is interpreted as having been located on the distal north-western rim of the Paleotethys (Stampfli & Kozur, 2006). Its lower part, which consists mainly of shallow-water carbonates almost devoid of macrofossils (no ammonoids), is only present in the eastern A-section (Grădinaru et al., 2007).

The fault was recognized in the western Deșli Caira B-section about 150 meters west of section A (Balini et al., 2024; Horacek & Grădinaru 2023: Section A equals section 100/500, section B equals section 810). The first appearance datum (FAD) of *Chiosella timorensis* (Nogami, 1968) was reported between GR6 and GR7 by Grădinaru et al., (2007), Orchard et al., (2007), and later was moved down to level 817 of the Grădinaru numbering by Golding (2021). The first occurrence datum (FOD) of *Stenopopanoceras transiens* Tozer, 1972, the ammonoid that could potentially serve to mark the lowermost Anisian, was identified at the base of bed 822A2 (Balini et al., 2024).

RESULTS AND DISCUSSION

The fault, once its presence is recognized, is quite prominent and easy to follow, as it runs more or less perpendicular to the bedding (Fig. 2A-C, 3, 4), with an offset of 80 to 100 cm. As excavation and cleaning of the outcrop (an old quarry) progressed (see Fig. 4), we have provided new photos that show the improved accessibility (due to excavation and cleaning) of the individual beds in the section, and thus, also the visibility of the fault, which now can be followed more or less continuously from the top of the section (bed 836) down to bed 812. It is now obvious that several beds were labeled incorrectly by Grădinaru (as also




Figure 2 – Overview of middle and upper part of Deșli Caira B-section, showing the uppermost Olenekian – lowermost Anisian succession and the two faults originally identified in Balini et al., (2024 see fig. 4), and shown again in Balini et al., (2026). A) General view of the upper part of the succession; B) Detail of the proposed boundary interval (Balini et al., (2024, 2026) of the section; C) Detail of the uppermost part of the section. Note that the direction of movement of the (assumed) right/eastern fault (dashed red line) has been changed. White ellipse encircles Alexandra Lăcătuș who provides scale for the photo (ca. 160 cm). See text and Balini et al. (2024, 2026) for additional explanations. Photo taken by a drone.




Figure 3 – A, B: Detail of the boundary interval, as proposed in Balini et al., (2024, 2026). A: as indicated in Balini et al., (2026), B: with the displacement by the fault indicated in the section by the horizontal lines representing the top of 822A2. The two red arrows in the photograph on both sides of the fault indicate the relative movement with respect to the other side of the fault.

mentioned in Balini et al., 2024, Balini et al., 2026) e.g. right side (old) 830 is left side 829, right side (old) 826 is left side 825, right side (old) 824 is left side 823, right side (old) 819 is (approximately) left side 817 (see Fig. 4 and Fig. 5). Below bed 817 (see Fig. 4), the situation is not completely clear and still under investigation. These results will be published in a future article. As seen in the lower part of the B-section after cleaning and excavating (Fig. 4), correlation between both sides of the

fault is still under investigation, and until complete correlation is achieved, we recommend that ongoing work be conducted only along the left (western) side of the section to rule out any errors due to the fault. As can be seen on the photos, there is more than sufficient space on both sides of the fault. To provide an impression of the good fit of both sides of the fault when the offset is taken into account, we show for a part of the section the logs from both sides (Fig. 5).

Meanwhile, ammonoid investigations are progressing and the latest results suggest that *Pseudacrochordiceras*, whose range is restricted to the uppermost part of the *Neopopanoceras haugi* Zone and the *Courtilloticeris stevensi* Zone of the North American scale, both latest Olenekian, may be present in bed GR7. If this occurrence is confirmed, the late Olenekian of Deşli Caira will be correlatable with the Nevada ammonoid scale, and thus, with the Panthalassa realm.

CONCLUSIONS

A fault runs more or less perpendicular to the bedding in the center of section B at Deşli Caira, resulting in a displacement of 80 to 100 cm between both sides of the fault. However, about 30 m of room exists on each side of the fault, leaving sufficient space to work, thus enabling a sampling study of the section without having to cross the fault. In most parts of the section (except the lower part) one can even move from one side of the fault to the other without problem, because the fault and its displacement are easily recognized, and bed thicknesses on both sides of the fault show a very good fit when the offset is taken into account. For the sake of clearness, we propose to only work on the left side of the fault, to avoid any problem of correct correlation. The Deşli Caira section possesses rich ammonoid and conodont faunas and has been investigated for several other fossil groups, as well as paleomag and carbon isotopes. Furthermore, a detailed lithologic accounting is available and will be extended further downwards in the near future, thus enabling very detailed investigations of the entire succession and most importantly, the boundary interval. Ongoing ammonoid investigations suggest the occurrence of the ammonoid *Pseudacrochordiceras* in the late Olenekian of Deşli Caira section. This taxon is crucial for long range correlation with the succession of the late Olenekian *Neopopanoceras haugi* and *Courtilloticeris stevensi* ammonoid zones of Nevada and beyond. Finally, in our opinion, the Deşli Caira section is best suited for eventual selection as the GSSP for the Olenekian-Anisian Boundary.

ACKNOWLEDGMENTS

This work was financially supported by the Research Institute of the University of Bucharest (ICUB), grant CPI 835/2023-2025 to Iuliana Lazăr. We thank J.F. Jenks and S. Lucas for very constructive and timely reviews that significantly improved the manuscript, and editor C.A. McRoberts for very skillful handling. J. F. Jenks is especially thanked for significantly improving the language.




Figure 4 – Photograph of the lower (excluding the lowermost) to middle part of the B-section, which was excavated and cleaned in 2025. Red line represents the fault. Red ovals on right side of fault indicate numbers by E. Grădinaru, which need to be corrected with respect to the left side of the fault. Violet ovals represent E. Grădinaru’s numbers to the left side of the fault, which are regarded as references in the section. White ovals show E. Grădinaru’s numbers on the right side, which are presently in use and have been correlated accordingly to the left side of the fault. Blue oval indicates one of Grădinaru’s numbers on the left side, which has been abandoned for the number on the right side of the fault. Other red numbers are of recent age, taking the fault into account. Thin dashed yellow lines represent individual beds. The black chisel (red arrow) next to the black-and-orange hammer in the lower left center (left of the fault) is ~30cm long. The two red arrows in the center of the photograph on both sides of the fault represent the relative movement with respect to the other side of the fault.

REFERENCES

- Atudorei, N.-V. 1999. Constraints on the Upper Permian to Upper Triassic marine carbon isotope curve. Case studies from the Tethys. PhD Thesis. Faculty of Sciences, University of Lausanne, Switzerland. 160 p.
- Balini, M., Lăcătuș, A., Grădinaru, E. & Lazăr, I. 2024. Aegean ammonoids from the Anisian GSSP candidate section Deșli Caira (Northern Dobrogea, Romania, Riv. It. Paleont. Strat., 130, 711-759, doi: 10.54103/2039-4942/24683.
- Balini, M., Horacek, M., & Lazăr, I. 2026. Comment on the paper “Detailed conodont data from the Olenekian-Anisian boundary interval of the GSSP candidate section at Deșli Caira, Romania”, by Martyn L. Golding. *J. Micropalaeontology*, 45, 27–31. <https://doi.org/10.5194/jm-45-27-2026>.
- Chen, Y., Jiang, H., Ogg, J.G., Zhang, Y., Gong, Y., & Yan, C. 2020. Early-Middle Triassic boundary interval: Integrated chemo-bio-magneto-stratigraphy of potential GSSPs for the base of the Anisian Stage in South China. *Earth Planet. Sci. Lett.* 530, 115863. <https://doi.org/10.1016/j.epsl.2019.115863>
- Forel, M.B., & Grădinaru, E. 2021. A unique diversity hotspot for Polycopidae (Ostracoda) in the Triassic of North Dobrogea. *Historical Biology* 34(6), 995–1018. <https://doi.org/10.1080/08912963.2021.19>.
- Golding, M.L. 2021. Early Anisian (Middle Triassic) conodonts from Romania and China, with comments on their role in the recognition and correlation of the base of the Anisian. *Journal of Earth Science* 32 (3), 573–591. <https://doi.org/10.1007/s12583-020-1392-9>.
- Golding, M.L. 2025. Detailed conodont data from the Olenekian-Anisian boundary interval of the GSSP

Section B1 on the left side of the fault




Figure 5 – Lithologs from both sides of the fault with the offset (0.8-1.0m) taken into account, demonstrating a good comparability of thicknesses. Note that only the upper part (Bed 823 to 836) of the section is shown here, both logs continue downwards. The logs were measured ca. 0.5–2m away on each side of the fault. Note that on the left side Bed 836 has been added with respect to earlier logs. Red amonionid symbol shows presence of Aegean amonionids, white amonionid symbol identifies layers containing amonionids undetermined up to now. Asterisk on the east side log identifies bed 823, which is not exposed directly at the fault but in the “eastern trench” (Balini et al., 2024), ca. 20m east of the fault.

candidate section at Deșli Caira, Romania, Journal of Micropalaeontology 44, 1–17.

- Grădinaru, E. 2000. Introduction to the Triassic geology of North Dobrogea Orogene—an overview of the Triassic System in the Tulcea Unit and the amonionid biostratigraphy. In: Grădinaru, E. (Ed.), Workshop on the Lower-Middle Triassic (Olenekian-Anisian) Boundary, 7-10 June, Tulcea, Romania. Field Trip Section, pp. 5–37.
- Grădinaru, E., Kozur, H.W., Nicora, A., & Orchard, M.J. 2006. The *Chiosella timorensis* and correlation of the amonionids and conodonts around the base of the Anisian in the GSSP candidate at Deșli Caira (North Dobrogea, Romania). Albertiana 34, 34–38.
- Grădinaru, E., Orchard, J.J., Nicora, A., Gallet, Y., Besse, J.,

Krystyn, L., Sobolev, E.S., Atudorei, N.-V., & Ivanova, D. 2007. The Global Boundary Stratotype Section and Point (GSSP) for the base of the Anisian Stage: Deșli Caira Hill, North Dobrogea, Romania. Albertiana 36, 54–71.

- Grădinaru, E., & Gaetani, M. 2019. Upper Spathian to Bithynian (Lower to Middle Triassic) brachiopods from North Dobrogea (Romania). Riv. Ital. Paleontol. Stratigr. 125 (1), 91–123. <https://doi.org/10.13130/2039-4942/11182>.
- Grădinaru, E. 2022. the Olenekian-Anisian/early-middle Triassic boundary, and assessment of the potential of conodonts for chronostratigraphic calibration of the Triassic timescale. ACTA Palaeontol. Roman. 18 (2), 3–5. <https://doi.org/1035463/j.apr.2022.02.01>.

- Horacek, M., Grădinaru, E. 2023. The Spathian Anisian (Lower-Middle Triassic) Boundary in the candidate GSSP section at Deșli Caira, Romania: Review of existing data, new findings, and comparison with other candidates. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 111407. <https://doi.org/10.1016/j.palaeo.2023.111407>.
- Kittl, E. 1908. Beiträge zur Kenntnis der Triasbildungen der nordöstlichen Dobrudscha. *Denkschr.-Österr. Akad. Wiss. Math.Nat. Kl.* 81, 447–532.
- Krystyn, L., Horacek, M., Brandner, R., Bhat, G., G. 2019. News from the Permian-Triassic Guryul Ravine Section (Kashmir/India): A fault causing biostratigraphic correlations and remarks on the Dinerian conodont UAZ's. *Albertiana*, vol. 45, p. 1–4.
- Mirăuță, E. 1974. Über die Conodontenfauna des oberen Werfens des tieferen Anis der nördlichen Dobrudscha/Rumänien. *Geol. Paleontol.* 8, 149–158.
- Mirăuță, E., & Gheorghian, D. 1978. In: Étude microfaunique des formations triasiques (transylvaines, bucoviniennes et getiques) des Carpates Orientales, 643/3. *Dări de seamă ale Sedintelor, Institutul de Geologie si Geofizică*, pp. 109–162.
- Muttoni, G., Nicora, A., Balini, M., Katz, M., Schaller, M., Kent, D.V., Maron, M., Meco, S., Rettori, R., Doda, V., & Nazaj, S. 2019. A candidate GSSP for the base of the Anisian from Kcira, Albania. *Albertiana* 45, 39–49.
- Nogami, Y. 1968. Trias-Conodonten von Timor, Malaysien und Japan, in: *Memoirs of the Faculty of Science, Series Geology and Mineralogy, Kyoto University*, 34, 115–136.
- Orchard, M. J., Grădinaru, E., and Nicora, A. 2007. A summary of the conodont succession around the Olenekian-Anisian boundary at Deșli Caira, North Dobrogea, Romania, in: *The Global Triassic*, edited by: Lucas, S. G. and Spielmann, J. A., *New Mexico Museum Nat. Hist. Sci. Bulletin*, 41, 341–346.
- Simionescu, I. 1910. Studii geologice si palaontologice din Dobrogea. III. Fauna triasică dela Delsi-Caira. La faune triasique de Deșli-Caira (Dobrogea). *Acad. Roman. Publ. Fond. V. Adamachi* 5 (26), 495–524. [in Romanian, French summary].
- Simionescu, I. 1911. Studii geologice si paleontologice din Dobrogea. V. Fauna triasică inferioară din Dobrogea (La faune du Trias inférieur de Dobrogea). *Acad. Roman. Publ. Fond. V. Adamachi* 5 (29), 63–79. [In Romanian, French summary].
- Stampfli, G.M. & Kozur, H. 2006. Europe from the Variscan to the Alpine cycles. In: *Gee, D.G. & Stephenson, R. (eds.) European Lithosphere Dynamics*. Geological Society, London, *Memoirs*, 32, 57–82.
- Tozer, E.T. 1972. The Earliest Marine Triassic Rocks: their Definition, Ammonoid fauna, Distribution and relationship to underlying formations. *Bull. Can. Petrol. Geol.*, 20 (4), p. 643–650.

